


Lecture 4

Multivariable linear regression

Sung Kim <hunkim+ml@gmail.com>
<http://hunkim.github.io/ml/>

Video (Korean): <https://youtu.be/kPxpJY6fRkY>



Recap

- Hypothesis
- Cost function
- Gradient descent algorithm

Recap

- Hypothesis

$$H(x) = Wx + b$$

- Cost function

$$cost(W, b) = \frac{1}{m} \sum_{i=1}^m (H(x^{(i)}) - y^{(i)})^2$$

- Gradient descent algorithm

Predicting exam score: regression using one input (x)

one-variable
one-feature

x (hours)	y (score)
10	90
9	80
3	50
2	60
11	40

Predicting exam score: regression using three inputs (x_1 , x_2 , x_3)

multi-variable/feature

x_1 (quiz 1)	x_2 (quiz 2)	x_3 (midterm 1)	Y (final)
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142

Test Scores for General Psychology

Hypothesis

$$H(x) = Wx + b$$

Hypothesis

$$H(x) = Wx + b$$

$$H(x_1, x_2, x_3) = w_1x_1 + w_2x_2 + w_3x_3 + b$$

Cost function

$$H(x_1, x_2, x_3) = w_1x_1 + w_2x_2 + w_3x_3 + b$$

$$cost(W, b) = \frac{1}{m} \sum_{I=1}^m (H(x_1^{(i)}, x_2^{(i)}, x_3^{(i)}) - y^{(i)})^2$$

Multi-variable

$$H(x_1, x_2, x_3) = w_1x_1 + w_2x_2 + w_3x_3 + b$$

$$H(x_1, x_2, x_3, \dots, x_n) = w_1x_1 + w_2x_2 + w_3x_3 + \dots + w_nx_n + b$$

Matrix

$$w_1x_1 + w_2x_2 + w_3x_3 + \dots + w_nx_n$$

Matrix multiplication

The diagram shows the multiplication of two matrices. The first matrix is $\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$ and the second matrix is $\begin{bmatrix} 7 & 8 \\ 9 & 10 \\ 11 & 12 \end{bmatrix}$. A yellow curved arrow labeled "Dot Product" connects the first row of the first matrix (1, 2, 3) to the first column of the second matrix (7, 9, 11). The result is shown as $= \begin{bmatrix} 58 & \end{bmatrix}$, with the value 58 highlighted in a yellow circle.

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \times \begin{bmatrix} 7 & 8 \\ 9 & 10 \\ 11 & 12 \end{bmatrix} = \begin{bmatrix} 58 & \end{bmatrix}$$

Hypothesis using matrix

$$w_1x_1 + w_2x_2 + w_3x_3 + \dots + w_nx_n$$

$$(x_1 \quad x_2 \quad x_3) \cdot \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} = (x_1w_1 + x_2w_2 + x_3w_3)$$

$$H(X) = XW$$

Hypothesis using matrix

$$H(x_1, x_2, x_3) = x_1w_1 + x_2w_2 + x_3w_3$$

x_1	x_2	x_3	Y
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142

Test Scores for General Psychology

Hypothesis using matrix

$$H(x_1, x_2, x_3) = x_1w_1 + x_2w_2 + x_3w_3$$

x_1	x_2	x_3	Y
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142

Test Scores for General Psychology

$$(x_1 \quad x_2 \quad x_3) \cdot \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} = (x_1w_1 + x_2w_2 + x_3w_3)$$

$$H(X) = XW$$

Many x instances

\mathbf{x}_1	\mathbf{x}_2	\mathbf{x}_3	\mathbf{Y}
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142

Test Scores for General Psychology

$$(x_1 \quad x_2 \quad x_3) \cdot \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} = (x_1 w_1 + x_2 w_2 + x_3 w_3)$$

x_1	x_2	x_3	Y
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142

Hypothesis using matrix

$$w_1x_1 + w_2x_2 + w_3x_3 + \dots + w_nx_n$$

$$\begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \\ x_{41} & x_{42} & x_{43} \\ x_{51} & x_{52} & x_{53} \end{pmatrix} \cdot \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} = \begin{pmatrix} x_{11}w_1 + x_{12}w_2 + x_{13}w_3 \\ x_{21}w_1 + x_{22}w_2 + x_{23}w_3 \\ x_{31}w_1 + x_{32}w_2 + x_{33}w_3 \\ x_{41}w_1 + x_{42}w_2 + x_{43}w_3 \\ x_{51}w_1 + x_{52}w_2 + x_{53}w_3 \end{pmatrix}$$

$$H(X) = XW$$

Hypothesis using matrix

$$\begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \\ x_{41} & x_{42} & x_{43} \\ x_{51} & x_{52} & x_{53} \end{pmatrix} \cdot \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} = \begin{pmatrix} x_{11}w_1 + x_{12}w_2 + x_{13}w_3 \\ x_{21}w_1 + x_{22}w_2 + x_{23}w_3 \\ x_{31}w_1 + x_{32}w_2 + x_{33}w_3 \\ x_{41}w_1 + x_{42}w_2 + x_{43}w_3 \\ x_{51}w_1 + x_{52}w_2 + x_{53}w_3 \end{pmatrix}$$

[5, 3]

[3, 1]

[5, 1]

$$H(X) = XW$$

Hypothesis using matrix

$$\begin{array}{ccc} \left(\begin{array}{|c|} \hline \mathbf{X} \\ \hline \end{array} \right) & \times & \left(\begin{array}{|c|} \hline \mathbf{W} \\ \hline \end{array} \right) = \left(\begin{array}{|c|} \hline \mathbf{H(X)} \\ \hline \end{array} \right) \\ [5, 3] & [?, ?] & [5, 1] \end{array}$$

$$H(X) = XW$$

Hypothesis using matrix

$$\begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \\ x_{41} & x_{42} & x_{43} \\ x_{51} & x_{52} & x_{53} \end{pmatrix} \cdot \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} = \begin{pmatrix} x_{11}w_1 + x_{12}w_2 + x_{13}w_3 \\ x_{21}w_1 + x_{22}w_2 + x_{23}w_3 \\ x_{31}w_1 + x_{32}w_2 + x_{33}w_3 \\ x_{41}w_1 + x_{42}w_2 + x_{43}w_3 \\ x_{51}w_1 + x_{52}w_2 + x_{53}w_3 \end{pmatrix}$$

$[n, 3] \qquad [3, 1] \qquad [n, 1]$

$$H(X) = XW$$

Hypothesis using matrix (n output)

$$\begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \\ x_{41} & x_{42} & x_{43} \\ x_{51} & x_{52} & x_{53} \end{pmatrix} \cdot \begin{matrix} ? \\ ? \\ ? \\ ? \\ ? \end{matrix} = \begin{pmatrix} x_{11}w_{11} + x_{12}w_{21} + x_{13}w_{31} & x_{11}w_{12} + x_{12}w_{22} + x_{13}w_{32} \\ x_{21}w_{11} + x_{22}w_{21} + x_{23}w_{31} & x_{21}w_{12} + x_{22}w_{22} + x_{23}w_{32} \\ x_{31}w_{11} + x_{32}w_{21} + x_{33}w_{31} & x_{31}w_{12} + x_{32}w_{22} + x_{33}w_{32} \\ x_{41}w_{11} + x_{42}w_{21} + x_{43}w_{31} & x_{41}w_{12} + x_{42}w_{22} + x_{43}w_{32} \\ x_{51}w_{11} + x_{52}w_{21} + x_{53}w_{31} & x_{51}w_{12} + x_{52}w_{22} + x_{53}w_{32} \end{pmatrix}$$

$[n, 3] \quad [?, ?] \quad [n, 2]$

$$H(X) = XW$$

Hypothesis using matrix (n output)

$$\begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \\ x_{41} & x_{42} & x_{43} \\ x_{51} & x_{52} & x_{53} \end{pmatrix} \cdot \begin{pmatrix} w_{11} & w_{12} \\ w_{21} & w_{22} \\ w_{31} & w_{32} \end{pmatrix} = \begin{pmatrix} x_{11}w_{11} + x_{12}w_{21} + x_{13}w_{31} & x_{11}w_{12} + x_{12}w_{22} + x_{13}w_{32} \\ x_{21}w_{11} + x_{22}w_{21} + x_{23}w_{31} & x_{21}w_{12} + x_{22}w_{22} + x_{23}w_{32} \\ x_{31}w_{11} + x_{32}w_{21} + x_{33}w_{31} & x_{31}w_{12} + x_{32}w_{22} + x_{33}w_{32} \\ x_{41}w_{11} + x_{42}w_{21} + x_{43}w_{31} & x_{41}w_{12} + x_{42}w_{22} + x_{43}w_{32} \\ x_{51}w_{11} + x_{52}w_{21} + x_{53}w_{31} & x_{51}w_{12} + x_{52}w_{22} + x_{53}w_{32} \end{pmatrix}$$

[n, 3]

[3, 2]

[n, 2]

$$H(X) = XW$$

WX vs XW

- Lecture (theory):

$$H(x) = Wx + b$$

- Implementation (TensorFlow)

$$H(X) = XW$$

Next **Logistic Regression** **(Classification)**

