

Chipsplitting Games: A Combinatorial Approach to Classifying One-Dimensional Discrete
Statistical Models with Rational Maximum Likelihood Estimator

by

Viet-Duc Nguyen

A thesis submitted in partial satisfaction of the

requirements for the degree of

Master of Science

in

Mathematics

in the

Mathematics Department

of

Technische Universität Berlin

Supervisors:

Professor Carlos Améndola

Professor Christian Haase

Winter 2024

Eidesstattliche Erklärung

Hiermit versichere ich, dass ich die vorliegende Arbeit eigenständig ohne Hilfe Dritter und ausschließlich unter Verwendung der aufgeführten Quellen und Hilfsmittel angefertigt habe. Alle Stellen die den benutzten Quellen und Hilfsmitteln unverändert oder sinngemäß entnommen sind, habe ich als solche kenntlich gemacht.

Sofern generative KI-Tools verwendet wurden, habe ich Produktnamen, Hersteller, die jeweils verwendete Softwareversion und die jeweiligen Einsatzzwecke (z.B. sprachliche Überprüfung und Verbesserung der Texte, systematische Recherche) benannt. Ich verantworte die Auswahl, die Übernahme und sämtliche Ergebnisse des von mir verwendeten KI-generierten Outputs vollumfänglich selbst.

Die Satzung zur Sicherung guter wissenschaftlicher Praxis an der TU Berlin vom 8. März 2017 habe ich zur Kenntnis genommen.

Ich erkläre weiterhin, dass ich die Arbeit in gleicher oder ähnlicher Form noch keiner anderen Prüfungsbehörde vorgelegt habe.

Berlin, den

Zusammenfassung in deutscher Sprache

Die vorliegende Ausarbeitung setzt die Forschung von Arthur Bik und Orlando Marigliano zur Klassifizierung ein-dimensionaler diskreter statistischer Modelle mit rationalen Maximum Likelihood Schätzern unter Verwendung fundamentaler Modelle fort. Wir erzielen bedeutende Fortschritte beim Beweis zur endlichen Anzahl der fundamentalen Modelle im Wahrscheinlichkeitssimplex Δ_5 . Zudem bestimmen wir die Anzahl der fundamentalen Modelle im Simplex Δ_6 mit einem maximalen Grad von 11.

Abstract

Chipsplitting Games: A Combinatorial Approach to Classifying One-Dimensional Discrete Statistical Models with Rational Maximum Likelihood Estimator

by

Viet-Duc Nguyen

Technische Universität Berlin

This paper continues the research of Arthur Bik and Orlando Marigliano on the classification of one-dimensional discrete statistical models with rational maximum likelihood estimators using fundamental models. We present a missing proof of an algorithm from their work. Furthermore, we make significant progress in proving the finite number of fundamental models in the probability simplex Δ_5 . We also determine the number of fundamental models in the simplex Δ_6 with a maximum degree of 11.

Contents

Contents	i
1 Introduction	1
2 Classification with Fundamental Models	3
2.1 Parametrization	3
2.2 Reduced Models	7
2.3 Fundamental Models	8
2.4 On the Finiteness of Fundamental Models	11
3 Chipsplitting Games	14
3.1 Basic Definitions	14
3.2 Pascal Equations	17
3.3 Valid Outcomes and Reduced Statistical Models	22
4 Supports of Valid Outcomes	29
4.1 Invertibility Criterion	29
4.2 Divide and Conquer	31
4.3 Symmetry	33
4.4 Impossible Supports	36
5 Valid Outcomes of Positive Support Size One, Two, and Three	39
6 Constraints on Supports of Valid Outcomes	43
6.1 Hyperfield Criterion	43
6.2 Solving Homogeneous Hyperfield Linear Systems	49
6.3 Implementation of the Hyperfield Criterion	53
6.4 Contractions	54
7 Valid Outcomes of Positive Support Size Four	63
Bibliography	66

Chapter 1

Introduction

In statistics we come across various collections of probability distributions, such as the normal distribution, Poisson distribution, and binomial distribution. These distributions are used to model random variables in applications, and are referred to as *statistical models*. Precisely, a statistical model is just a set of probability distributions. If the set contains only discrete distributions, we call it a *discrete statistical model*. In this case, discrete statistical models are just subsets of the probability simplex $\Delta_n := \{p \in \mathbb{R}^{n+1} \mid \sum p_i = 1\}$.

A discrete distribution $p \in \mathcal{M} \subset \Delta_n$ from a discrete statistical model encapsulates the probabilities of observing the states $0, \dots, n$, i.e. if $X \in \{0, \dots, n\}$ is a discrete random variable, then the state $X = i$ occurs with probability p_i for all $i = 0, \dots, n$. Say we have a binomial random variable X with $n + 1$ states, then $p_i = \binom{n}{i} \theta^i (1 - \theta)^{n-i}$ computes the probability of observing i successes in n trials with success probability $\theta \in [0, 1]$. The set \mathcal{M} of all probability distributions of that form, i.e. $\mathcal{M} = \{(\binom{n}{i} \theta^i (1 - \theta)^{n-i})_{i=0}^n \mid \theta \in [0, 1]\}$, is our first example of a discrete statistical model, and is known as the *binomial model*.




Figure 1.1: This figure shows the probability simplex Δ_2 with the binomial model (red curve). Every point on the curve is a binomial distribution.

Given a statistical model $\mathcal{M} \subset \Delta_n$ and data $u \in \mathbb{N}^{n+1}$, a typical problem in statistics is to find a distribution from a statistical model that best describes the data. “Best” can mean a lot of things, but in *maximum likelihood estimation* it means finding the distribution that maximizes the probability of observing the data; the map $\Phi : \Delta_n \rightarrow \mathcal{M}, u \mapsto \hat{p}$ that assigns the data u to a distribution $\hat{p} \in \mathcal{M}$ from the statistical model is called the *maximum likelihood estimator (MLE)*. This map is characterized by the property that \hat{p} maximizes the log-likelihood function $\ell(p) = \sum u_i \log p_i$ for all $p \in \mathcal{M}$.

We focus on **one-dimensional discrete statistical models with rational MLE**. These are models \mathcal{M} satisfying

- $\mathcal{M} = \text{image}(p)$ for some rational map $p = (p_0, \dots, p_n) : I \rightarrow \Delta_n$ where p_i is rational, $I \subset \mathbb{R}$ is a union of closed intervals and $p(\partial I) \subset \partial \Delta_n$,
- all the $n + 1$ coordinates of the maximum likelihood estimator Φ are rational functions in the data u .

There are two intriguing questions to ask about statistical models with rational MLE: the first one is about which *form* do the maximum likelihood estimators take; the second one is more concerned with the *classification* of the statistical models, i.e. can we divide these models into easier to understand classes? An answer to the first question was given by June Huh. He showed that if Φ is rational, then each of its coordinates is an alternative product of linear forms with numerator and denominator of the same degree, see [4, 3]. For the second question, Arthur Bik and Orlando Marigliano classified all one-dimensional discrete statistical models with rational MLE using *fundamental models* [2].

This thesis continues the work of Arthur Bik and Orlando Marigliano. In the first half, we present their classification results on how fundamental models serve as the building blocks of one-dimensional discrete models with rational MLE. In the second half, we establish and extend their finding that there are only finitely many fundamental models within the probability simplices Δ_n for $n \leq 4$. Due to the complexity of the problem, the cases $n \geq 5$ were left open. For the first time, we make significant progress for $n = 5$. Other contributions of this thesis include a new result on the number of fundamental models in Δ_6 with a maximum degree of 11, and a missing proof of a key algorithm from their work.

Chapter 2

Classification with Fundamental Models

In this chapter we present the classification of one-dimensional discrete statistical models with rational maximum likelihood estimator (MLE) using fundamental models. The classification is due to Arthur Bik and Orlando Marigliano [2].

Problem statement: Can we find a class of easy to understand models that serve as building blocks for all one-dimensional discrete statistical models with rational MLE?

The answer to this question are *reduced* and *fundamental models*.

2.1 Parametrization

It turns out that one-dimensional discrete statistical models with rational MLE admit the following parametrization.

Proposition 2.1. *Let \mathcal{M} be a one-dimensional discrete statistical models with rational maximum likelihood estimator. Then, there exists a map of the form*

$$p : [0, 1] \rightarrow \Delta_n, \quad \theta \mapsto (w_k \theta^{i_k} (1 - \theta)^{j_k})_{k=0}^n$$

$$i_k, j_k \in \mathbb{Z}_{\geq 0}, \quad w_k \in \mathbb{R}_{>0} \quad \forall k = 0, \dots, n$$

such that $\mathcal{M} = \text{image}(p)$.

We introduce some notation to simplify the proof of Proposition 2.1. Let $\mathcal{M} \subset \Delta_n$ be a one-dimensional discrete statistical model parametrized by rational functions $p_0 = \frac{g_0}{h_0}, \dots, p_n = \frac{g_n}{h_n}$. Define b to be the least common multiple of h_0, \dots, h_n and $a_i := bp_i$. Since $\sum p_k = 1$, we can multiply by b to obtain $\sum a_k = b$. We see that the polynomials a_0, \dots, a_n, b

determine the statistical model \mathcal{M} , and have no common factors. The log-likelihood function is then given by

$$\begin{aligned}\ell(p) &= \sum u_i \log p_i \\ &= \sum u_i \log \frac{a_i}{b} \\ &= \sum u_i \log a_i - \sum u_i \log b.\end{aligned}$$

To find the maximum likelihood estimator, we need find all critical points of the log-likelihood function. This is equivalent to finding the roots of the gradient of the log-likelihood function

$$\ell(p(\theta))' = \sum u_k \frac{a'_k}{a_k} - \sum u_k \frac{b'}{b} = 0. \quad (2.1)$$

These equations are called the *score equations* in algebraic statistics, and the number of complex solutions to these equations for general data $u \in \mathbb{C}^{n+1}$ is called the *maximum likelihood degree* of the statistical model. This ML degree has an important meaning in algebraic statistics, as it determines the complexity of the model. We have the following relationship between the ML estimator and the ML degree.

Proposition 2.2. *Having rational maximum likelihood estimator can be expressed equivalently by saying that the maximum likelihood degree of the statistical model is one.*

Proof. Refer to [3] for a proof. □

To prove Proposition 2.1, we need the following lemma.

Lemma 2.3. *If \mathcal{M} has rational MLE, then there are exactly two distinct complex linear factors in a_0, \dots, a_n , and b .*

Proof. We prove the lemma in three steps:

- Let f be the product of all distinct complex linear factors in a_0, \dots, a_n, b . If we multiply the score equations (2.1) by f , we get

$$f \cdot \ell(p(\theta))' = \sum u_k f \frac{a'_k}{a_k} - \sum u_k f \frac{b'}{b} = 0.$$

Note that every linear factor of a_k with multiplicity m occurs in a'_k with multiplicity $m - 1$; thus every summand of $\frac{a'_k}{a_k}$ is of the form $\frac{\lambda}{(x-\xi)}$, where $\lambda \in \mathbb{R}$ and $x - \xi$ is some linear factor of a_k ; hence $f \cdot \frac{\lambda}{(x-\xi)}$ is of degree $\deg(f) - 1$, and therefore $f \cdot \ell(p(\theta))'$ is of degree $\deg(f) - 1$.

- We claim that the roots of $\ell(p(\theta))'$ are the same as the roots of $f \cdot \ell(p(\theta))'$. Assume we have shown this claim. By Proposition 2.2 the ML degree is one. So, $\ell(p(\theta))'$ has one root. Thus, $f \cdot \ell(p(\theta))'$ has one root, and therefore $f \cdot \ell(p(\theta))'$ is of degree one. This implies that $\deg(f) = 2$ with the previous step. Thus, there are exactly two distinct complex linear factors in a_0, \dots, a_n , and b .
- It remains to show that the roots stay the same. Clearly, every root of $\ell(p(\theta))'$ is a root of $f \cdot \ell(p(\theta))'$. Conversely, we want to show that no new roots are introduced when multiplying by f , i.e. roots of f are not roots of $f \cdot \ell(p(\theta))'$. To do so, we rewrite

$$\begin{aligned} f \cdot \ell(p(\theta))' &= \sum_{k=0}^n u_k f \frac{a'_k}{a_k} - \sum_{k=0}^n u_k f \frac{b'}{b} = \sum_{k=0}^{n+1} v_k f \frac{c'_k}{c_k} \\ v_k &= u_k, \quad c_k = a_k \quad \text{for } k = 0, \dots, n, \\ v_{n+1} &= -\sum_{k=0}^n u_k, \quad c_{n+1} = b. \end{aligned}$$

Let q be a complex linear factor of f . We define polynomials r_0, \dots, r_{n+1} and r such that $c_k = q^{l_k} r_k$, $f = qr$, and r_0, \dots, r_{n+1}, r do not have q as a factor. Then, we have for $k = 0, \dots, n+1$ that

$$f \frac{c'_k}{c_k} = qr \cdot \frac{l_k q^{l_k-1} q' r_k + q^{l_k} r'_k}{q^{l_k} r_k} = qr \frac{l_k q'}{q} + qr \frac{r'_k}{r_k} \equiv r l_k q' \pmod{q}.$$

Thus, we obtain

$$f \cdot \ell(p(\theta))' \equiv r q' \sum_{k=0}^{n+1} v_k l_k \equiv r q' \sum_{k=0}^n v_k (l_k - l_{n+1}) \pmod{q}.$$

Note that by definition of l_k , a value of $l_k = 0$ means that q is not a factor of c_k . By definition of f , at least one $l_k > 0$. On the other hand, not all l_k can be positive since a_0, \dots, a_n, b share no common factors. Hence, not all $l_k - l_{n+1} = 0$ vanish. Hence, for generic data u we assume $\sum_{k=0}^n v_k (l_k - l_{n+1}) \neq 0$. This with $q'r \not\equiv 0 \pmod{q}$ implies that q is not a complex linear factor of $f \cdot \ell(p(\theta))'$. We showed that the roots of f are not roots of $f \cdot \ell(p(\theta))'$.

□

Equipped with the lemma, we can now prove Proposition 2.1.

Proof. We want to show the following parametrization of \mathcal{M} :

$$p : [0, 1] \rightarrow \Delta_n, \quad \theta \mapsto (w_k \theta^{i_k} (1 - \theta)^{j_k})_{k=0}^n$$

First, we show that I is a single closed real interval and not a union of closed intervals. For the sake of contradiction assume that $I = \bigcup_k I_k$ is a union of closed disjoint intervals. By definition of \mathcal{M} we know that $p(\partial I) \subset \partial \Delta_n$. Thus, there exist $\theta_1, \theta_2 \in \partial I_0$ and $\theta_3, \theta_4 \in \partial I_1$ with $p_i(\theta_1) = p_i(\theta_2) = 0$ and $p_j(\theta_3) = p_j(\theta_4) = 0$ for some $i, j = 0, \dots, n$. Note that θ_1, θ_2 are roots of $\frac{a_i}{b}$ and θ_3, θ_4 are roots of $\frac{a_j}{b}$. By Lemma 2.3 exactly two distinct complex linear factors occur in a_0, \dots, a_n, b . Hence, $\theta_3 = \theta_1$ or $\theta_3 = \theta_2$. Contradiction for I_0 and I_1 are disjoint.

The previous argument shows that $I = [\alpha, \beta]$ is a real single closed interval. Thus, the roots of a_0, \dots, a_n, b are real and take values in $\partial I = \{\alpha, \beta\}$. By a suitable parametrization, we can assume without loss of generality that $I = [0, 1]$. We can now write the polynomials a_0, \dots, a_n, b as

$$\begin{aligned} a_k(\theta) &= w_k \theta^{i_k} (1 - \theta)^{j_k} \\ b(\theta) &= w \theta^i (1 - \theta)^j \end{aligned}$$

with $w_k, w \in \mathbb{R}_{>0}$, and $i_k, j_k, i, j \in \mathbb{Z}_{\geq 0}$ for all $k = 0, \dots, n$. Since a_0, \dots, a_n, b share no common factors, there exists some $i_k = 0$ if $i > 0$; however this would contradict $0 < w_k \leq a_0(0) + \dots + a_n(0) = b(0) = 0$. So $i = 0$. Similarly, $j = 0$. Finally, we divide p by w to obtain $b \equiv 1$. \square

Corollary 2.4. *Any one-dimensional discrete statistical models with rational MLE can be represented by $(w_k, i_k, j_k)_{k=0}^n$ for $w_k \in \mathbb{R}_{>0}$ and $i_k, j_k \in \mathbb{Z}_{\geq 0}$.*

From now on, we only consider one-dimensional discrete statistical models with rational MLE; we call them *models* for short.

Definition 2.5. The degree $\deg(\mathcal{M})$ of a model \mathcal{M} represented by $(w_k, i_k, j_k)_{k=0}^n$ is defined as $\max \{i_k + j_k : k = 0, \dots, n\}$.

Remark 2.6. We view two models $(w_k, i_k, j_k)_{k=0}^n$ and $(w'_k, i'_k, j'_k)_{k=0}^n$ as the same model if they are equal up to a permutation of the coordinates.

Example 2.7. The sequence $((1, 0, 2), (2, 1, 1), (1, 2, 0))$ represents the binomial model with two trials. It has degree two. Its parametrization is given by $\theta \mapsto ((1 - \theta)^2, 2\theta(1 - \theta), \theta^2)$. Also see Figure 1.1 for a visualization of the binomial model within the probability simplex Δ_2 .

Note that we view the sequences $((1, 0, 2), (2, 1, 1), (1, 2, 0))$ or $((2, 1, 1), (1, 0, 2), (1, 2, 0))$ as the same model as $((2, 1, 1), (1, 2, 0), (1, 0, 2))$ since the order of the coordinates does not matter.

Definition 2.8. Let \mathcal{M} be a model represented by $(w_k, i_k, j_k)_{k=0}^n$. The set of exponent pairs $(i_k, j_k)_{k=0}^n$ is called the support of \mathcal{M} , denoted by $\text{supp}(\mathcal{M})$.

This was our first step towards understanding the structure of models. The next step is to introduce the concept of reduced models.

2.2 Reduced Models

Models in this section refer to one-dimensional discrete statistical models with rational MLE.

Definition 2.9. We call a model represented by $(w_k, i_k, j_k)_{k=0}^n$ *reduced* if $(i_k, j_k) \neq \mathbf{0}$ for all $k = 0, \dots, n$, and $(i_k, j_k) \neq (i_l, j_l)$ for all $k \neq l$.

Due to $(i_k, j_k) \neq (i_l, j_l)$, we can use functions to represent reduced models.

Remark 2.10. A reduced model \mathcal{M} represented by $(w_k, i_k, j_k)_{k=0}^n$ can also be identified by a function $f : \mathbb{Z}^2 \rightarrow \mathbb{R}_{\geq 0}$, $(i, j) \mapsto w$, where $w = w_k$ if $(i_k, j_k) = (i, j)$ and $w = 0$ otherwise. The support of f is the set of all pairs (i, j) with $f(i, j) > 0$. It coincides with the support of \mathcal{M} .

Reduced models are our first building blocks for the classification of models. This statement is justified by the following two propositions. They show that every non-reduced model can be transformed into a reduced model by a sequence of linear embeddings.

Proposition 2.11. Let $n \in \mathbb{N}_{>0}$. Let \mathcal{M} be a model represented by $(w_k, i_k, j_k)_{k=0}^n$. If $(i_l, j_l) = \mathbf{0}$ for some index l , then there exist a model \mathcal{M}' , $\lambda \in [0, 1]$ and $k = 0, \dots, n$ such that

$$\mathcal{M} = \Psi_{\lambda, k}(\mathcal{M}'),$$

where $\Psi_{\lambda, k} : \Delta_{n-1} \rightarrow \Delta_n$ is defined as $p_i \mapsto \begin{cases} \lambda p_i & \text{if } k \neq i, \\ 1 - \lambda & \text{if } k = i. \end{cases}$

Proof. Let $(i_l, j_l) = \mathbf{0}$ for some index l . If $w_l = 1$, then $w_m = 0$ for all $m \neq l$; this contradicts $w_m > 0$ by Proposition 2.1. Set $\lambda = 1 - w_l > 0$ and $k = l$. Define the model \mathcal{M}' represented by

$$\left(\frac{w_h}{1 - w_l}, i_h, j_h \right)_{h=0, h \neq l}^n.$$

Then, $\mathcal{M} = \Psi_{\lambda, k}(\mathcal{M}')$. □

Proposition 2.12. Let $n \in \mathbb{N}_{>0}$. Let \mathcal{M} be model represented by $(w_k, i_k, j_k)_{k=0}^n$. If $(i_m, j_m) = (i_l, j_l)$ for $m \neq l$, then there exist a model \mathcal{M}' , $\lambda \in [0, 1]$ and $k, h = 0, \dots, n$ such that

$$\mathcal{M} = \Psi_{\lambda, k, h}(\mathcal{M}'),$$

where $\Psi_{\lambda, k, h} : \Delta_{n-1} \rightarrow \Delta_n$ is defined as $p_i \mapsto \begin{cases} p_i & \text{if } i \notin \{k, h\}, \\ \lambda p_k & \text{if } k = i, \\ (1 - \lambda)p_h & \text{if } h = i. \end{cases}$

Proof. Define $\lambda = \frac{w_m}{w_m + w_l}$, $k = m$, and $h = l$. Define the model \mathcal{M}' represented by

$$(w_g + \delta_{gm} w_l, i_g, j_g)_{g=0, g \neq l}^n.$$

Then, $\mathcal{M} = \Psi_{\lambda, k}(\mathcal{M}')$. □

By repeatedly applying the two propositions, we can transform any model into a reduced model.

Corollary 2.13. *If Δ_n contains a model of degree d , then there also exists a reduced model of degree d in Δ_m for some $m \leq n$.*

2.3 Fundamental Models

As before, models refer to one-dimensional discrete statistical models with rational MLE. The main building blocks for the classification of models are *fundamental models*; we will see that reduced models come from fundamental models.

Definition 2.14. We call a model represented by $(w_k, i_k, j_k)_{k=0}^n$ *fundamental* if it is reduced and the equation $p_0 + \dots + p_n \equiv 1$ for given $(i_k, j_k)_{k=0}^n$ uniquely determines the weights $(w_k)_{k=0}^n$.

Example 2.15. The binomial model with two trials is fundamental. Given $(i_0, j_0) = (0, 2)$, $(i_1, j_1) = (1, 1)$, and $(i_2, j_2) = (2, 0)$, the equation $p_0 + p_1 + p_2 = w_0 \theta^2 + w_1 \theta(1 - \theta) + w_2 (1 - \theta)^2 \equiv 1$ uniquely determines the weights $w_0 = 1, w_1 = 2, w_2 = 1$. To see this observe that this equation is equivalent to $w_0 \theta^2 + w_1 \theta - w_1 \theta^2 + w_2 - w_2 2\theta + w_2 \theta^2 = 1$ which is equivalent to solving $w_2 - 1 + \theta(w_1 - 2w_2) + \theta^2(w_0 - w_1 + w_2) = 0$ for all $\theta \in \mathbb{R}$.

Example 2.16. Consider the probability simplex Δ_0 . It only contains the model 1 which is fundamental.

Example 2.17. Now, consider the probability simplex Δ_1 . It only contains the models $\theta \mapsto (\theta, 1 - \theta)$ and $\theta \mapsto (1 - \theta, \theta)$ which are equivalent. They are fundamental.

We will see that fundamental models like the ones above are building blocks for all reduced models by *composition*.

Definition 2.18. Let \mathcal{M} and \mathcal{M}' be reduced models which are represented by functions $f, g : \mathbb{Z}^2 \rightarrow \mathbb{R}_{\geq 0}$, see Remark 2.10. Let $\mu \in (0, 1)$. The *composite* $\mathcal{M} *_{\mu} \mathcal{M}'$ of \mathcal{M} and \mathcal{M}' is the reduced model represented by the function

$$(i, j) \mapsto \mu f(i, j) + (1 - \mu)g(i, j).$$

We are about to show that every reduced model is the composite of finitely many fundamental models.

Proposition 2.19. *Let \mathcal{M} be a reduced model. Then \mathcal{M} is the composite of finitely many fundamental models.*

Proof. For Δ_0 and Δ_1 we know that they only contain fundamental models, see Examples 2.16 and 2.17.

Assume we are given Δ_n with $n \geq 2$, and let \mathcal{M} be a model that is not fundamental. We aim to show that \mathcal{M} can be expressed as a composite of two models, \mathcal{M}' and \mathcal{M}'' , whose supports are proper subsets of $\text{supp}(\mathcal{M})$. Assume this is indeed the case. Then, by applying the same argument to \mathcal{M}' and \mathcal{M}'' , we can recursively decompose each non-fundamental model into models with smaller supports. Since $\text{supp}(\mathcal{M})$ is finite, this recursive decomposition must eventually terminate, yielding a decomposition of \mathcal{M} into fundamental models. Thus, we have shown that any reduced model is the composite of a finite number of fundamental models.

Let us prove that \mathcal{M} is the composite of two models whose supports are proper subsets of $\text{supp}(\mathcal{M})$. Since \mathcal{M} is not fundamental, the equation $p_0 + \dots + p_n = 1$ has distinct solutions $\mathbf{w}, \mathbf{w}' \in \mathbb{R}_{>0}^{n+1}$. Define $\mathbf{v} := \mathbf{w} - \mathbf{w}' \neq \mathbf{0}$. Then,

$$\sum_{k=0}^n v_k \theta^{i_k} (1 - \theta)^{j_k} = 0 \quad \forall \theta \in (0, 1).$$

Observe that there exist strictly positive and negative coefficients v_k . Define

$$\begin{aligned} \lambda &:= \min \left\{ \frac{w_k}{|v_k|} : k = 0, \dots, n, v_k < 0 \right\}, \\ u_k &:= w_k + \lambda v_k \quad \text{for } k = 0, \dots, n, \\ S_1 &:= \{(i_k, j_k) : k = 0, \dots, n, u_k \neq 0\}. \end{aligned}$$

Note that $\lambda > 0$ since all the coefficients w_k are strictly positive by definition. Also observe that $u_k \geq 0$ if $v_k \geq 0$. Moreover, by definition $\frac{w_k}{|v_k|} \geq \lambda$ for all $k \geq 0$. Hence, if $v_k < 0$, we also have $\frac{u_k}{v_k} = \frac{w_k}{v_k} + \lambda \leq 0$. Multiplying by $v_k < 0$ we obtain $u_k \geq 0$. All in all, we have $u_k \geq 0$ for all $k = 0, \dots, n$. Moreover, $u_k = 0$ if and only if $v_k < 0$ and $\lambda = \frac{w_k}{|v_k|}$. This shows that $S_1 \subsetneq \text{supp}(\mathcal{M})$. Since $u_0 + \dots + u_n = 1$, we have found a reduced model \mathcal{M}' represented by $(u_k, i_k, j_k)_{(i_k, j_k) \in S_1}$.

For the second model, we define

$$\begin{aligned} \mu &:= \min \left\{ \frac{w_k}{u_k} : k = 0, \dots, n, u_k \neq 0 \right\}, \\ t_k &:= \frac{w_k - \mu u_k}{1 - \mu} \quad \text{for } k = 0, \dots, n, \\ S_2 &:= \{(i_k, j_k) : k = 0, \dots, n, t_k \neq 0\}. \end{aligned}$$

Similarly, $\mu > 0$. We have $\mu < 1$ because some v_k is positive implying $u_k > w_k$. By definition, we have $t_k \geq 0$, and $t_k = 0$ if and only if $u_k \neq 0$ and $\mu = \frac{w_k}{u_k}$. This shows that $S_2 \subsetneq \text{supp}(\mathcal{M})$.

and $S_1 \cup S_2 = \text{supp}(\mathcal{M})$. Since $t_0 + \cdots + t_n = 1$, we have found a reduced model \mathcal{M}'' represented by $(t_k, i_k, j_k)_{(i_k, j_k) \in S_2}$.

Finally, we see that $w_k = \mu u_k + (1 - \mu)t_k$. This shows that $\mathcal{M} = \mathcal{M}' *_{\mu} \mathcal{M}''$. \square

By applying the previous proposition with Corollary 2.13, we obtain the following corollary.

Corollary 2.20. *If Δ_n contains a non-fundamental model of degree d , then there exists a fundamental model of degree d in Δ_m for some $m < n$.*

Example 2.21. For the two-dimensional probability simplex Δ_2 , we can classify all models. Again, models refer to one-dimensional discrete statistical models with rational MLE. Note that the model \mathcal{M} parametrized by $\theta \mapsto (\theta, 1 - \theta)$ satisfies $\mathcal{M} *_{\mu} \mathcal{M} = \mathcal{M}$ for all μ . Since Δ_1 only contains the model $\theta \mapsto (\theta, 1 - \theta)$, we can conclude that Δ_2 only contains fundamental models or models that are not reduced.

To find all the fundamental models in Δ_2 , we need to check for all sets $S = \{(i_k, j_k)\}_{k=0}^2 \subset \mathbb{Z}_{>0}^2$ of size three if the equation $p_0 + p_1 + p_2 = \sum_{k=0}^2 w_k \theta^{i_k} (1 - \theta)^{j_k} = 1$ has a unique solution (w_0, w_1, w_2) . As we can see, a priori infinitely many sets S need to be checked. However, as we will see in the next section, only those sets S with $\max\{i + j : (i, j) \in S\} \leq 2n - 1 = 3$ need to be considered. Clearly, this reduces the number of sets S to be checked to a finite number.

We compute that only the following supports uniquely determine the weights (w_0, w_1, w_2) :

$$\{(0, 3), (1, 1), (3, 0)\}, \{(0, 2), (1, 1), (2, 0)\}, \{(0, 1), (1, 1), (2, 0)\}, \{(0, 2), (1, 0), (1, 1)\}.$$

They correspond to the fundamental models $((1 - \theta)^3, 3\theta(1 - \theta), \theta^3)$, $((1 - \theta)^2, 2\theta(1 - \theta), \theta^2)$, $(1 - \theta, \theta(1 - \theta), \theta^2)$, and $((1 - \theta)^2, \theta, \theta(1 - \theta))$. The last model is equivalent to the second last model by a parametrization $\theta \mapsto 1 - \theta$ and permutation of the coordinates.



Figure 2.1: From left to right the illustration depicts the models parametrized $((1 - \theta)^3, 3\theta(1 - \theta), \theta^3)$, $((1 - \theta)^2, 2\theta(1 - \theta), \theta^2)$, $(1 - \theta, \theta(1 - \theta), \theta^2)$, and $((1 - \theta)^2, \theta, \theta(1 - \theta))$. The illustration is taken from [2].

We just computed all fundamental models of degree three or less in Δ_2 . We will see shortly that these are all models in the probability simplex Δ_2 . Of course, Δ_2 contains non-reduced models, too. These are models that come from linear embeddings $\Psi_{\lambda,k}$ and $\Psi_{\lambda,k,h}$, see Proposition 2.11 and Proposition 2.12. There are infinitely many of them, and for $\lambda = \frac{1}{3}$ we obtain the models $\theta \mapsto (\frac{2}{3}\theta, \frac{1}{3}, \frac{2}{3}(1-\theta))$ and $\theta \mapsto (1-\theta, \frac{1}{3}\theta, \frac{2}{3}\theta)$.



Figure 2.2: This illustration depicts two non-reduced models in Δ_2 for $\lambda = \frac{1}{3}$. They are parametrized by $\theta \mapsto (\frac{2}{3}\theta, \frac{1}{3}, \frac{2}{3}(1-\theta))$ and $\theta \mapsto (1-\theta, \frac{1}{3}\theta, \frac{2}{3}\theta)$. All other non-reduced models can be obtained by varying λ . The illustration is taken from [2].

Let us summarize the results of this section. It is the first part of our classification theorem.

Theorem 2.22. *Every one-dimensional discrete statistical model with rational MLE in Δ_n is the image of a reduced model in Δ_m under a linear embedding $\Delta_m \rightarrow \Delta_n$ for some $m \leq n$.*

Moreover, every reduced model $\mathcal{M} \subset \Delta$ can be written as a composite of finitely many fundamental models

$$\mathcal{M} = \mathcal{M}_1 *_{\mu_1} (\cdots *_{\mu_{m-2}} (\mathcal{M}_{m-1} *_{\mu_{m-1}} \mathcal{M}_m))$$

for some $m < n$ and $\mu_1, \dots, \mu_m \in (0, 1)$.

Proof. See Proposition 2.19, Proposition 2.11, and Proposition 2.12. □

2.4 On the Finiteness of Fundamental Models

We have established the first part of our classification theorem, namely that fundamental models are building blocks for all models. The second part is showing that there are only finitely many fundamental models in Δ_n given $n \in \mathbb{N}$. Artuhr Bik and Orlando Marigliano proved that there are only finitely many fundamental models in Δ_n for $n \leq 4$ [2]. We will

later make significant progress towards proving the case $n = 5$. For $n \geq 6$ no attempt has been made yet to the best of our knowledge.

Arthur Bik and Orlando Marigliano first proved the following proposition.

Theorem 2.23. *Let \mathcal{M} be a one dimensional discrete statistical model with rational MLE in Δ_n . For $n \leq 4$ we have $\deg(\mathcal{M}) \leq 2n - 1$.*

Given Theorem 2.23 it is easy to show the second part of our classification.

Theorem 2.24. *There are only finitely many fundamental models in Δ_n for all $n \leq 4$.*

Proof. By Theorem 2.23 we know that the degree of a fundamental model is at most $2n - 1$. Since the number of supports of a fundamental model of degree $2n - 1$ is finite, there are only finitely many fundamental models in Δ_n for all $n \leq 4$. \square

We will now spend the rest of this thesis on proving Theorem 2.23. The idea is to use the building blocks of fundamental models that we have established so far. Namely, it suffices to show the theorem for fundamental models.

Theorem 2.25. *Let $N \in \mathbb{N}$. If for all $n \leq N$ and for all fundamental models $\mathcal{M} \in \Delta_n$ the upper bound $\deg(\mathcal{M}) \leq 2n - 1$ holds, then the upper bound also holds for all statistical models in $\Delta_{n'}$ for all $n' \leq N$.*

Proof. Let $N \in \mathbb{N}$ and $n \leq N$. Assume there is some non-fundamental model \mathcal{M}' in Δ_n of degree greater than $2n - 1$. By Corollary 2.20 there exists a fundamental model \mathcal{M} in Δ_m for some $m < n$ of degree greater than $2m - 1$. This contradicts the assumption that the degree of fundamental models is at most $2n' - 1$ for all $n' \leq N$. \square

Counting all *fundamental* models in Δ_n for $n \leq 4$ is our guiding objective. As a first step, we introduce a combinatorial game that aids in counting fundamental models. We know that every reduced model can be represented by the sequence of triples $(w_k, i_k, j_k)_{k=0}^n$, where $w_k \in \mathbb{R}_{>0}$ and $i_k, j_k \in \mathbb{Z}_{\geq 0}$. The model can be visualized in a directed graph with vertices in \mathbb{Z}^2 , where we can place values w_k on vertices (i_k, j_k) . Each vertex (i, j) is connected by directed edges to $(i + 1, j)$ and $(i, j + 1)$.

Surprisingly, we can derive a combinatorial game from this graph by defining a specific set of rules. This game, called the *chipsplitting game*, will be rigorously introduced in the next chapter. After that, we will explore the game's properties and show how it can be used to count fundamental models in Δ_n for $n \leq 4$.




Figure 2.3: The binomial model with two trials visualized in a directed graph with vertices in $\{0, 1, 2, 3\}^2$.

Chapter 3

Chipsplitting Games

The notion of a chipsplitting game was introduced by [2] as a combinatorial approach to classifying one-dimensional discrete statistical models with rational maximum likelihood estimator. It was inspired by *chipfiring games* and for a subset of chipfiring games, the chipsplitting game is equivalent to the chipfiring game. We refer to [5] for a comprehensive introduction to chipfiring games.

3.1 Basic Definitions

Let us define the notion of a chipsplitting game.

Definition 3.1. Let (V, E) be a directed graph without loops.

1. A *chip configuration* is a vector $\mathbf{w} = (w_v)_{v \in V} \in \mathbb{Z}^V$ such that there are only finitely many nonzero components w_k .
2. The *initial configuration* is the chip configuration $\mathbf{0} \in \mathbb{Z}^V$.
3. A *splitting move* at $u \in V$ maps a chip configuration \mathbf{w} to some chip configuration \mathbf{w}' defined by

$$w'_v := \begin{cases} w_v - 1 & \text{if } v = u, \\ w_v + 1 & \text{if } (u, v) \in E \\ w_v & \text{otherwise.} \end{cases}$$

This map is denoted by split_u .

4. An *unsplitting move* at $u \in V$ maps \mathbf{w}' back to \mathbf{w} . This map is denoted by unsplit_u .
5. A *chipsplitting game* is a finite sequence of splitting and unsplitting moves.

6. An *outcome* of a chipsplitting game is the chip configuration obtained from applying the sequence of splitting and unsplitting moves defined by the game at the initial configuration.
7. Any outcome of a chipsplitting game is called an *outcome*.

Proposition 3.2. *The order of the moves in a chipsplitting game does not affect the outcome.*

Proof. This follows from commutativity of addition. \square

Note that all moves are reversible. Thus, we obtain the following corollary with Proposition 3.2.

Corollary 3.3. *Let \mathbf{w} be an outcome. Then, there exists a chipsplitting game whose outcome is \mathbf{w} and where at no point both a splitting and an unsplitting move are applied at the same vertex.*

Games that satisfy the condition in the corollary are called *reduced*. We will only consider reduced games in this thesis for simplicity. The map

$$\begin{aligned} \{\text{reduced games on } (V, E)\} / \sim &\rightarrow \{g : V' \rightarrow \mathbb{Z} : \#\{p \in V' : g(p) \neq 0\} < \infty\} \\ f &\mapsto (p \mapsto \text{number of moves at } p \text{ in game } f) \end{aligned}$$

is a bijection, where $V' \subset V$ is the subset of vertices with at least one outgoing edge. The equivalence relation \sim is defined by $f \sim g$ if f and g are the same up to reordering. Unsplitting moves are counted negatively by $p \mapsto \text{number of moves at } p \text{ in game } f$. Using the map above we identify a chipsplitting game with its corresponding function $V' \rightarrow \mathbb{Z}$. For every outcome $\mathbf{w} = (w_v)_{v \in V}$ we have

$$w_v = -f(v) + \sum_{u \in V', (u,v) \in E} f(u),$$

where we define $f(v) = 0$ for $v \notin V'$.

Now, we define the directed graphs that we will consider in this thesis. For $d \in \mathbb{N} \cup \{\infty\}$ we write

$$\begin{aligned} V_d &:= \{(i, j) \in \mathbb{Z}_{\geq 0}^2 \mid i + j \leq d\}, \\ E_d &:= \{(v, v + e) \mid v \in V_{d-1}, e \in \{(1, 0), (0, 1)\}\}. \end{aligned}$$

Definition 3.4. The degree $\deg(\mathbf{v})$ of a vertex $\mathbf{v} = (i, j)$ is defined as $i + j$.

Example 3.5. A chip configuration $\mathbf{w} = (w_{i,j})_{(i,j) \in V_d} \in \mathbb{Z}^{V_d}$ can be illustrated as a triangle of numbers where $w_{i,j}$ is placed at the position (i, j) in the triangle. For example, $w_{2,4} = 4$ means that the value 4 is placed in the second column and fourth row of the triangle. The following is an example of a sequence of chip configurations for $d = 3$:

.	.	.	1	1	1
. .	. .	1 .	. 1	. 1	. .
. . .	1 . .	. 2 .	. 2 .	. 2 1	. 3 .
0 . . .	-1 1 . .	-1 . 1 .	-1 . 1 .	-1 . . 1	-1 . . 1

When $w_{i,j} = 0$, we omit the value in the triangle and write a dot instead. The sequence above starts with the initial configuration and then applies a splitting move at the vertex $(0, 0)$, $(1, 0)$, $(0, 1)$, $(0, 2)$ and $(2, 0)$. Finally, we apply an unsplitting move at the vertex $(1, 1)$ to obtain the final configuration. Coming back to figure 2.4, we see that it is represented as the third configuration of the triangle above.

Let us define some more terminology.

Definition 3.6. Let $\mathbf{w} = (w_{i,j})_{(i,j) \in V_d}$ be a chip configuration.

1. The *positive support* of \mathbf{w} is defined as $\text{supp}^+(\mathbf{w}) := \{(i, j) \in V_d \mid w_{i,j} > 0\}$.
2. The *negative support* of \mathbf{w} is defined as $\text{supp}^-(\mathbf{w}) := \{(i, j) \in V_d \mid w_{i,j} < 0\}$.
3. The *support* of \mathbf{w} is defined as the union of the positive and negative support.
4. The *degree* of \mathbf{w} is defined as $\deg(\mathbf{w}) := \max \{i + j \mid (i, j) \in \text{supp}(\mathbf{w})\}$.
5. We say \mathbf{w} is *valid* if its negative support is empty or only contains $(0, 0)$.

We are interested in *outcomes* that are *valid* since they will correspond to reduced models as we will see later. For that reason, it would be convenient to have a criterion for when a chip configuration is an outcome. The next section will provide such a criterion with the help of *Pascal equations*.

Example 3.7. Consider the following chip configuration:

.
.	1
1	.	5
.	5	.	2	.	.	.
1	.	.	5	.	.	.
.	.	8	.	.	2	.
-2	2	.

We clearly see that this configuration is valid, but is it also an outcome of a chipsplitting game? Currently, the only way to answer this question is to apply all possible sequences of splitting and unsplitting moves to the initial configuration and check if the outcome is the given configuration. In the next section, we present an easily computable characterization to answer this question.

3.2 Pascal Equations

In this chapter we will establish that outcomes are roots of Pascal equations. So let us first define Pascal equations which are special cases of *linear forms*.

Definition 3.8. A *linear form* on \mathbb{Z}^{V_d} is a map of the form

$$\mathbb{Z}^{V_d} \rightarrow \mathbb{Z}, \quad \mathbf{w} \mapsto \sum_{(i,j) \in V_d} c_{i,j} w_{i,j}.$$

It is denoted by $\sum_{(i,j) \in V_d} c_{i,j} x_{i,j}$.

Definition 3.9. A *Pascal form* on \mathbb{Z}^{V_d} is a linear form $\sum_{(i,j) \in V_d} c_{i,j} x_{i,j}$ on \mathbb{Z}^{V_d} satisfying

$$c_{i,j} = c_{i+1,j} + c_{i,j+1} \quad \text{for all } (i,j) \in V_{d-1}.$$

Example 3.10. We can visualize a Pascal form as a triangle of numbers where $c_{i,j}$ is placed at the position (i,j) in the triangle. Here are examples of Pascal forms for $d = 2$:

0	1	0	0
1 1	1 0	0 0	1 1
2 1 0	1 0 0	1 1 1	0 -1 -2

Evaluating Pascal equations is invariant under splitting and unsplitting moves.

Proposition 3.11. Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be a chip configuration. Let $p = \sum c_{i,j} x_{i,j}$ be a Pascal equation on \mathbb{Z}^{V_d} . Then, we have $p(\mathbf{w}) = p(\text{split}_u(\mathbf{w})) = p(\text{unsplit}_v(\mathbf{w}))$ for all $u, v \in V_{d-1}$.

Proof. Let $u := (i', j') \in V_{d-1}$. By the Pascal property, we have

$$c_{i'+1, j'} + c_{i', j'+1} - c_{i', j'} = 0.$$

Thus, we have

$$\begin{aligned} p(\text{split}_u(\mathbf{w})) &= \sum_{(i,j) \in V_d} c_{i,j} (\text{split}_u(\mathbf{w}))_{i,j} \\ &= \sum_{(i,j) \in V_d} c_{i,j} \begin{cases} w_{i,j} - 1 & \text{if } (i,j) = u, \\ w_{i,j} + 1 & \text{if } (i,j) \in \{(i'+1, j'), (i', j'+1)\} \\ w_{i,j} & \text{otherwise} \end{cases} \\ &= \sum_{(i,j) \in V_d} c_{i,j} w_{i,j} = p(\mathbf{w}). \end{aligned}$$

Similarly, we can show that $p(\text{unsplit}_v(\mathbf{w})) = p(\mathbf{w})$ for all $v \in V_{d-1}$. □

Corollary 3.12. *Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be an outcome. Let $p = \sum c_{i,j}x_{i,j}$ be a Pascal equation on \mathbb{Z}^{V_d} . Then, $p(\mathbf{w}) = 0$.*

Proof. Clearly, we have $p(\mathbf{0}) = 0$. Then, we use Proposition 3.11 and the fact that \mathbf{w} is obtained from the initial configuration $\mathbf{0}$ by a sequence of splitting and unsplitting moves. \square

This demonstrates that outcomes are roots of Pascal equations. The converse is also true as we will see now. This is one of the most important results; so let us state it now.

Theorem 3.13. *Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be a chip configuration. Then, \mathbf{w} is an outcome if and only if \mathbf{w} is a root of all Pascal equations on \mathbb{Z}^{V_d} .*

The direction left to right is the content of the previous corollary. For the other direction life would be easier if we had not to deal with infinitely many Pascal equations. So let us fix this first by introducing a basis from which we can generate all Pascal equations through linear combinations.

Example 3.14. Fix the degree $d = 2$. We later claim that the following set of Pascal forms is a basis:

$$\begin{array}{ccc} 0 & 0 & 1 \\ 0 & 0 & 0 \end{array} \quad \begin{array}{ccc} 0 & 1 & 1 \\ 1 & 1 & 1 \end{array}, \quad \begin{array}{ccc} 0 & -1 & -2 \\ 0 & 0 & 1 \end{array}, \quad \begin{array}{ccc} 1 & -1 & 1 \\ 0 & 0 & 1 \end{array}.$$

Note that the first column of each Pascal form is a unit vector in \mathbb{R}^3 . We can also fix the first row of each Pascal form to be a unit vector in \mathbb{R}^3 :

$$\begin{array}{ccc} 1 & -2 & 1 \\ 1 & 0 & 0 \end{array}, \quad \begin{array}{ccc} -1 & 1 & 0 \\ 0 & 1 & 0 \end{array}, \quad \begin{array}{ccc} 0 & -1 & 1 \\ 0 & 0 & 1 \end{array}.$$

We will denote the first set of Pascal forms by $\{\text{col}(0), \text{col}(1), \text{col}(2)\}$ and the second set by $\{\text{row}(0), \text{row}(1), \text{row}(2)\}$.

To generalize the example above to an arbitrary degree $d \in \mathbb{N}$ and to vectors beyond unit vectors, we assert that there exists a unique Pascal form whose first column is any chosen vector.

Proposition 3.15. *Let $\mathbf{a} = (a_0, \dots, a_d)$ be any vector with integer entries. Then, the following two statements hold:*

1. *There exists a unique Pascal form $\sum c_{i,j}x_{i,j}$ such that $c_{0,\cdot} = \mathbf{a}$.*
2. *There exists a unique Pascal form $\sum c_{i,j}x_{i,j}$ such that $c_{\cdot,0} = \mathbf{a}$.*

Proof. Set $c_{0,\cdot} := \mathbf{a}$. Define $c_{i+1,j} := c_{i,j} - c_{i,j+1}$ for all $(i,j) \in V_d$ with $i = 0$. Then, we use the same formula to define $c_{i+1,j}$ for all $(i,j) \in V_d$ with $i = 1$. We repeat this process until we have defined all $c_{i,j}$ for $(i,j) \in V_d$.

For the second statement, we set $c_{\cdot,0} := \mathbf{a}$. Define $c_{i,j+1} := c_{i,j} - c_{i+1,j}$ for all $(i,j) \in V_d$ with $j = 0$. Then, we use the same formula to define $c_{i,j+1}$ for all $(i,j) \in V_d$ with $j = 1$. We repeat this process until we have defined all $c_{i,j}$ for $(i,j) \in V_d$. \square

Let us define our first two Pascal form bases.

Definition 3.16. Let $k = 0, \dots, d$ and $\mathbf{e}_k \in \mathbb{R}^{d+1}$ be the k -th unit vector.

- We define $\text{col}(k)$ to be the unique Pascal form $\sum c_{i,j} x_{i,j}$ such that $c_{0,\cdot} = \mathbf{e}_k$.
- We define $\text{row}(k)$ to be the unique Pascal form $\sum c_{i,j} x_{i,j}$ such that $c_{\cdot,0} = \mathbf{e}_k$.

For examples of the Pascal forms $\text{col}(k)$ and $\text{row}(k)$ for $d = 2$ see Example 3.14. We provide another example for $d = 7$.

Example 3.17. Let us consider the Pascal form $\text{col}(3)$ for $d = 7$. We visualize this Pascal form as follows:

$$\begin{array}{ccccccc}
 & & & & & & \\
 & & & & & & \\
 & & & & & & \\
 & & & & & & \\
 1 & 1 & 1 & 1 & 1 & & \\
 \cdot & -1 & -2 & -3 & -4 & -5 & \\
 \cdot & \cdot & 1 & 3 & 6 & 10 & 15 \\
 \cdot & \cdot & \cdot & -1 & -4 & -10 & -20 & -35.
 \end{array}$$

The Pascal form $\text{row}(3)$ is visualized as follows:

$$\begin{array}{ccccccc}
 -35 & & & & & & \\
 -20 & 15 & & & & & \\
 -10 & 10 & -4 & & & & \\
 -4 & 6 & -4 & 1 & & & \\
 -1 & 3 & -3 & 1 & \cdot & & \\
 \cdot & 1 & -2 & 1 & \cdot & \cdot & \\
 \cdot & \cdot & -1 & 1 & \cdot & \cdot & \cdot \\
 \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot & \cdot
 \end{array}$$

Proposition 3.18. *For all integers $k = 0, \dots, d$ the following formulas hold:*

$$\begin{aligned} \text{col}(k) &= (-1)^k \sum_{(i,j) \in V_d} (-1)^j \binom{i}{k-j} x_{i,j}, \\ \text{row}(k) &= (-1)^k \sum_{(i,j) \in V_d} (-1)^i \binom{j}{k-i} x_{i,j}. \end{aligned}$$

Note that $\binom{a}{b} = 0$ for $b < 0$ or $b > a$.

Proof. We claim that $(-1)^k \sum_{(i,j) \in V_d} (-1)^j \binom{i}{k-j} x_{i,j}$ is a Pascal equation. To see that observe

$$(-1)^j \binom{i+1}{k-j} + (-1)^{j+1} \binom{i}{k-j-1} = (-1)^j \binom{i}{k-j}$$

for all $(i,j) \in V_d$ due to $\binom{a}{b+1} + \binom{a}{b} = \binom{a+1}{b}$ where we set $a = i$ and $b = k - j - 1$. Next, we see that $(-1)^{k+j} \binom{0}{k-j} = \delta_{jk}$. Thus, $(-1)^k \sum_{(i,j) \in V_d} (-1)^j \binom{i}{k-j} x_{i,j}$ is indeed $\text{col}(k)$.

By symmetry of the binomial coefficients, we can use the same argument to show the second formula. \square

We now show that $\{\text{col}(k)\}_{k=0}^d$ is indeed a basis for all Pascal forms on \mathbb{Z}^{V_d} .

Proposition 3.19. *Let p be a Pascal form on \mathbb{Z}^{V_d} . The following statements hold:*

1. *There exist unique coefficients $\mu_0, \dots, \mu_d \in \mathbb{Z}$ such that $p = \mu_0 \text{col}(0) + \dots + \mu_d \text{col}(d)$.*
2. *There exist unique coefficients $\lambda_0, \dots, \lambda_d \in \mathbb{Z}$ such that $p = \mu_0 \text{row}(0) + \dots + \mu_d \text{row}(d)$.*

Proof. Let $p = \sum c_{i,j} x_{i,j}$ be a Pascal form on \mathbb{Z}^{V_d} . If we try to solve the equation

$$\sum_{(i,j) \in V_d} c_{i,j} x_{i,j} = \lambda_0 \text{col}(0) + \dots + \lambda_d \text{col}(d) \tag{3.1}$$

for $\lambda_0, \dots, \lambda_d$, then due to Proposition 3.18 we see for all $(i,j) \in V_d$ that we have

$$\begin{aligned} c_{i,j} &= \lambda_0 (-1)^{0+j} \binom{i}{0-j} + \lambda_1 (-1)^{1+j} \binom{i}{1-j} + \dots + \lambda_d (-1)^{d+j} \binom{i}{d-j} \\ &= \lambda_j (-1)^{2j} \binom{i}{0} + \lambda_{j+1} (-1)^{2j+1} \binom{i}{1} + \dots + \lambda_{i+j} (-1)^{2j+i} \binom{i}{i}. \end{aligned}$$

We see $c_{0,\cdot} = (\lambda_0, \dots, \lambda_d)$. Thus we set the coefficients $\boldsymbol{\mu} := c_{0,\cdot}$ and by Proposition 3.15 we see that $\sum_{(i,j) \in V_d} c_{i,j} x_{i,j} = \mu_0 \text{col}(0) + \dots + \mu_d \text{col}(d)$. Moreover, the same proposition shows that the coefficients $\lambda_0, \dots, \lambda_d$ in Equation 3.1 are uniquely determined.

For the second statement we use the same argument. \square

Corollary 3.20. *The set $\{\text{col}(k)\}_{k=0}^d$ is a basis for all Pascal forms on \mathbb{Z}^{V_d} . The same holds for $\{\text{row}(k)\}_{k=0}^d$.*

Proof. This follows from the previous proposition. \square

Let us come back to Theorem 3.13. We can now prove the other direction; namely that roots of all Pascal equations on \mathbb{Z}^{V_d} are outcomes.

Proposition 3.21. *Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be a chip configuration. If for all Pascal equations p on \mathbb{Z}^{V_d} we have $p(\mathbf{w}) = 0$, then \mathbf{w} is an outcome.*

Proof. Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be a chip configuration. By assumption

$$\text{col}(\deg(\mathbf{w}))(\mathbf{w}) = 0. \quad (3.2)$$


Note that by Proposition 3.18 for $\text{col}(\deg(\mathbf{w})) = \sum c_{i,j} x_{i,j}$ we have $c_{i,\deg(\mathbf{w})-i} = (-1)^i$ for all $i = 0, \dots, \deg(\mathbf{w})$. Moreover, we have

$$c_{i,j} = 0 \quad \text{for all } i + j < \deg(\mathbf{w}) \quad (3.3)$$

by Proposition 3.18. Together with Equation 3.2 and 3.3 we obtain

$$\sum_{i=0}^{\deg(\mathbf{w})} (-1)^i w_{i,\deg(\mathbf{w})-i} = 0. \quad (3.4)$$

Furthermore, we know that there exists a unique minimal set of splitting or unsplitting moves at vertices (i, j) of degree $\deg(\mathbf{w}) - 1$ such that applied to \mathbf{w} we obtain a chip configuration \mathbf{w}' with $w'_{i,j} = 0$ for all $i = 0, \dots, \deg(\mathbf{w})$. We call applying these set of moves to \mathbf{w} *retraction*.



```

.
.  .
.  .  .
.  .  .  .
*  .  .  .  .
*  *  .  .  .  .
*  *  *  .  .  .  .
*  *  *  *  .  .  .  .
*  *  *  *  *  .  .  .  .

```

Thus, \mathbf{w}' has degree less than $\deg(\mathbf{w})$. By Proposition 3.11 \mathbf{w}' is also a root of all Pascal equations. We repeat the retraction process $\deg(\mathbf{w})$ many times until we obtain some chip configuration of degree 0. This chip configuration is the initial configuration due to Equation 3.4. Thus, \mathbf{w} is an outcome. \square

We have shown Theorem 3.13. Characterizing outcomes as roots of Pascal equations is a powerful tool to determine if a chip configuration is an outcome.

Algorithm 1 Validating outcomes

Require: chipsplitting configuration $\mathbf{w} \in \mathbb{Z}^{V_d}$

Ensure: True if \mathbf{w} is an outcome, False otherwise

```

1: function ISOUTCOME( $A, n$ )
2:   initialize set  $S = \{\text{col}(0), \dots, \text{col}(\deg(\mathbf{w}))\}$ 
3:   for  $p$  of  $S$  do
4:     if  $p(\mathbf{w}) \neq 0$  then
5:       return False
6:     end if
7:   end for
8:   return True
9: end function

```

Proof of correctness of Algorithm 1. This follows from Theorem 3.13. \square

Example 3.22. Returning to Example 3.7, we see that the chip configuration is a root of all Pascal equations $\text{col}(0), \dots, \text{col}(6)$ using Algorithm 1. Thus, the chip configuration is an outcome.

3.3 Valid Outcomes and Reduced Statistical Models

In the previous sections, we have established that outcomes are roots of Pascal forms. Now, we will demonstrate that a subset of *valid outcomes* are in one-to-one correspondence with

reduced statistical models. Thus, we obtain not only a combinatorial characterization of reduced statistical models through chip-splitting games but also an algebraic characterization through Pascal equations. As before, statistical models mean one-dimensional discrete statistical models with rational maximum likelihood estimator.

We remind that valid chipsplitting configurations are those where the negative support is empty or only contains the vertex $(0, 0)$. Hence, valid outcomes are roots of Pascal equations whose negative supports are empty or only contain the vertex $(0, 0)$.

The function $\mathbf{w}(\mathcal{M})$ maps reduced models $\mathcal{M} = (w_k, i_k, j_k)_{k=0}^n$ to chip configurations $\mathbf{w}(\mathcal{M}) = (w_{i,j})_{(i,j) \in V_\infty}$ by

$$w_{i,j} := \begin{cases} -1 & \text{if } (i, j) = (0, 0), \\ w_k & \text{if } (i, j) = (i_k, j_k) \text{ for some } k, \\ 0 & \text{otherwise.} \end{cases}$$

Note that the map $\mathbf{w}(\mathcal{M})$ defines a *real* chipsplitting games; the rules of the game are the same as for integer chipsplitting games.

Example 3.23. The binomial model $((1, 3, 0), (3, 2, 1), (3, 1, 2), (1, 0, 3))$ with three trials is mapped to the chip configuration below:

$$\begin{array}{cccc} 1 & & & \\ \cdot & 3 & & \\ \cdot & \cdot & 3 & \\ -1 & \cdot & \cdot & 1. \end{array}$$

Example 3.24. Does the following valid real outcome from Example 3.7 induce a reduced statistical model through the inverse map \mathbf{w}^{-1} ?

$$\begin{array}{cccccc} \cdot & & & & & \\ \cdot & 0.5 & & & & \\ 0.5 & \cdot & 2.5 & & & \\ \cdot & 2.5 & \cdot & 1 & & \\ 0.5 & \cdot & \cdot & 2.5 & \cdot & \\ \cdot & \cdot & 4 & \cdot & \cdot & 1 \\ -1 & \cdot & \cdot & \cdot & \cdot & 1 \end{array}.$$

The outcome would correspond to the reduced model

$$\mathcal{M} = ((0.5, 2, 0), (0.5, 4, 0), (2.5, 1, 3), (0.5, 1, 5), (4, 2, 1), (2.5, 2, 4), (2.5, 3, 2), (1, 3, 3), (1, 5, 0), (1, 5, 1))$$

in the probability simplex Δ_9 . As it turns out \mathcal{M} is indeed a reduced statistical model by the next theorem.

Theorem 3.25. *The map $\mathcal{M} \mapsto w(\mathcal{M})$ is a bijection between reduced statistical models and valid real outcomes $\mathbf{w} \in \mathbb{R}^{V_\infty}$ with $w_{0,0} = -1$.*

$$\text{reduced models} \longleftrightarrow \text{valid outcomes}$$

Figure 3.1: Bijection between reduced models and valid real outcomes \mathbf{w} with $w_{0,0} = -1$

To show this theorem, we need to do some preparations. Let $\mathbb{R}[\theta]_{\leq d}$ denote the vector space of polynomials in the variable θ of degree at most d with real coefficients. Similarly, we define $\mathbb{Z}[\theta]_{\leq d}$ and $\mathbb{Q}[\theta]_{\leq d}$. Next, we introduce the linear map $\alpha_d^{\mathbb{R}}$ that maps real chip configurations to real polynomials:

$$\begin{aligned} \alpha_d^{\mathbb{R}} : \mathbb{R}^{V_d} &\rightarrow \mathbb{R}[\theta]_{\leq d}, \\ \mathbf{w} &\mapsto \sum_{(i,j) \in V_d} w_{i,j} \theta^i (1 - \theta)^j. \end{aligned}$$

We define the map $\alpha_d^{\mathbb{Z}}$ and $\alpha_d^{\mathbb{Q}}$ for integer and rational chip configurations analogously.

Lemma 3.26. *The following statements hold true for all $d \in \mathbb{N} \cup \{\infty\}$:*

1. $\{\mathbf{w} \in \mathbb{R}^{V_d} \mid \mathbf{w} \text{ is an outcome}\} = \text{kernel}(\alpha_d^{\mathbb{R}});$
2. $\{\mathbf{w} \in \mathbb{Z}^{V_d} \mid \mathbf{w} \text{ is an outcome}\} = \text{kernel}(\alpha_d^{\mathbb{Z}});$
3. $\{\mathbf{w} \in \mathbb{Q}^{V_d} \mid \mathbf{w} \text{ is an outcome}\} = \text{kernel}(\alpha_d^{\mathbb{Q}}).$

Proof. We only prove the first statement. The other two statements are proven analogously. Note that it suffices to show the statement for $d < \infty$ since $\alpha_\infty^{\mathbb{R}}$ is the direct limit of $\alpha_0^{\mathbb{R}}, \alpha_1^{\mathbb{R}}, \alpha_2^{\mathbb{R}}, \dots$, and so on.

Let $d < \infty$. By Corollary 3.20, the codimension of the outcome space is $d + 1$, as it is defined by the roots of the Pascal forms $\text{col}(0), \dots, \text{col}(d)$.

Let $f(\theta) = \lambda_0 + \lambda_1 \theta + \dots + \lambda_d \theta^d$ be a polynomial in \mathbb{R} of degree at most d . Define a chipsplitting configuration \mathbf{w} by

$$w_{i,j} := \begin{cases} \lambda_i & \text{if } j = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Then, $\alpha_d^{\mathbb{R}}(\mathbf{w}) = f$, which shows that the map $\alpha_d^{\mathbb{R}}$ is surjective. Hence, the kernel of $\alpha_d^{\mathbb{R}}$ has codimension $d + 1$; it has equal codimension as the space of outcomes.

Finally, we just need to show that the space of outcomes is contained in the kernel of $\alpha_d^{\mathbb{R}}$. Since their codimensions are equal, the two spaces must be equal. Let $\mathbf{w} \in \mathbb{R}^{V_d}$ be an

outcome. The value of $\alpha_d^{\mathbb{R}}(\mathbf{w})$ remains the same if apply splitting or unsplitting moves at arbitrary vertices $(i, j) \in V_{d-1}$ because we have

$$-\theta^i(1-\theta)^j + \theta^{i+1}(1-\theta)^j + \theta^i(1-\theta)^{j+1} = \theta^i(1-\theta)^j(-1 + \theta + (1-\theta)) = 0.$$

The remaining claim follows from $\alpha_d^{\mathbb{R}}(\mathbf{0}) = 0$. \square

We now show Theorem 3.25.

Proof of Theorem 3.25. Let $\mathcal{M} = (w_k, i_k, j_k)_{k=0}^n$ be a reduced model.

- First, we need to show that $\mathbf{w} := w(\mathcal{M})$ is an outcome; a-priori we only know that it is some chip configuration. By definition of $w(\mathcal{M})$, we have that $w_{0,0} = -1$. Since \mathcal{M} is a statistical model, we know that $\sum_{k=0}^n w_k \theta^{i_k} (1-\theta)^{j_k} \equiv 1$. Thus, $\alpha_d^{\mathbb{R}}(\mathbf{w}) = \sum_{k=0}^n w_k \theta^{i_k} (1-\theta)^{j_k} - 1 \equiv 0$. Thus, $\mathbf{w} \in \text{kernel}(\alpha_d^{\mathbb{R}})$. By Lemma 3.26, the chip configuration \mathbf{w} is an outcome.
- **Injectivity:** Let $\mathcal{M} = (w_k, i_k, j_k)_{k=0}^n$ and $\mathcal{M}' = (w'_k, i'_k, j'_k)_{k=0}^n$ be two distinct models. Then, $w(\mathcal{M}) \neq w(\mathcal{M}')$ (see Remark 2.6).
- **Surjectivity:** Let $\mathbf{w} \in \mathbb{R}^{V_\infty}$ be a valid real outcome with $w_{0,0} = -1$. We define

$$w_k := w_{i_k, j_k} \quad \text{for all } k = 0, \dots, n.$$

Then, $\mathcal{M} = (w_k, i_k, j_k)_{k=0}^n$ is a reduced model by Lemma 3.26. We see that $w(\mathcal{M}) = \mathbf{w}$. Hence, $\mathcal{M} \mapsto w(\mathcal{M})$ is surjective. \square

Next, we collect simple results.

Proposition 3.27. *The following statements hold for all reduced models \mathcal{M} :*

1. $\text{supp}^+(w(\mathcal{M})) = \text{supp}^+(\mathcal{M})$.
2. The map $\mathcal{M} \mapsto w(\mathcal{M})$ is degree-preserving.
3. The outcome $w(\mathcal{M})$ is a rational outcome if and only if all the coefficients of \mathcal{M} are rational.

Proof. All three statements follow directly from definitions. \square

Proposition 3.28. *Let $\mathbf{w} \in \mathbb{Q}^{V_\infty}$ be a valid rational outcome. Then, there exist positive $\lambda \in \mathbb{Q}$ and integral valid outcome $\mathbf{z} \in \mathbb{Z}^{V_\infty}$ such that $\mathbf{w} = \lambda \mathbf{z}$.*

Proof. Let \mathbf{w} be a valid rational outcome. Its support is finite. Thus, there exist $\mu \in \mathbb{N}$ such that $\mu \mathbf{w} \in \mathbb{Z}^{V_\infty}$. Define $\lambda := \frac{1}{\mu}$ and $\mathbf{v} := \mu \mathbf{w}$. Clearly, $\alpha_{\deg(\mathbf{v})}^{\mathbb{Z}}(\mathbf{v}) = \mu \alpha_{\deg(\mathbf{w})}^{\mathbb{Q}}(\mathbf{w}) \equiv 0$. By Lemma 3.26, \mathbf{v} is an outcome. It is valid because $\mu \mathbf{w}$ is valid. \square

Proposition 3.29. *Let $\mathbf{w} \in \mathbb{R}^{V_\infty}$ be a valid real outcome. If $w_{0,0} = 0$, then $\mathbf{w} = \mathbf{0}$.*

Proof. By Lemma 3.26, we have $\sum w_{i,j} \theta^i (1 - \theta)^j \equiv 0$. By assumption, the negative support is empty. Hence, all the $w_{i,j}$ are non-negative. We evaluate at $\theta = \frac{1}{2}$ to conclude that the positive support of \mathbf{w} is empty. Hence, $\mathbf{w} = \mathbf{0}$. \square

Let us go back to Example 3.23.

Example 3.30. We have seen that the valid real outcome below induces a reduced statistical model by Theorem 3.25.

$$\begin{array}{cccc} 1 & & & \\ \cdot & 3 & & \\ \cdot & \cdot & 3 & \\ -1 & \cdot & \cdot & 1. \end{array}$$

The outcome has degree three and positive support size four. Can we find another outcome with the same degree but smaller positive support size? Indeed we can unsplitt at vertex $(1, 1)$ to get

$$\begin{array}{cccc} 1 & & & \\ \cdot & \cdot & & \\ \cdot & 3 & \cdot & \\ -1 & \cdot & \cdot & 1. \end{array}$$

Can we reduce the positive support size further? As it will turn out, we cannot. The positive support size is minimal for a degree three outcome.

Example 3.31. Let us fix the positive support size to be three. What is the largest degree of a valid real outcome with positive support size three? We have already found

$$\begin{array}{cccc} 1 & & & \\ \cdot & \cdot & & \\ \cdot & 3 & \cdot & \\ -1 & \cdot & \cdot & 1. \end{array}$$

However is there an even larger one? Once again, the answer is no: by the following theorem, the degree of a valid real outcome with positive support of size three is at most three.

Theorem 3.32. *The following upper bound*

$$\deg(\mathbf{w}) \leq 2 \cdot |\text{supp}^+(\mathbf{w})| - 3$$

holds for valid integral outcomes \mathbf{w} with $|\text{supp}^+(\mathbf{w})| \leq 5$.

Arthur Bik and Marigliano Orlando established this theorem in [2], and we will provide a proof in the following sections. The question of whether the upper bound extends to cases with larger positive support sizes, $|\text{supp}^+(\mathbf{w})| > 5$, remains open. A primary contribution of this thesis is the significant advancement in proving this bound for the case $|\text{supp}^+(\mathbf{w})| = 6$.

Pos. Support Size	Max Degree
2	1
3	3
4	5
5	7

Table 3.1: The following table shows the largest degree of a valid real outcome with positive support size 2, 3, 4 and 5.

Theorem 3.32 is of particular interest because it is equivalent to Theorem 2.23, i.e. proving that the degree of outcomes is bounded by their positive support size is equivalent to establishing that the degree of statistical models is bounded by their dimension.

$$\begin{aligned} \text{degree of outcomes } \mathbf{w} &\longleftrightarrow \text{degree of statistical models } \mathcal{M} \\ \text{supp}^+(\mathbf{w}) &\longleftrightarrow \text{dimension of } \mathcal{M} \end{aligned}$$

Proposition 3.33. *Theorem 2.23 and Theorem 3.32 are equivalent.*

Proof. Assume Theorem 2.23 holds. We want to show Theorem 3.32. Let \mathbf{w} be a valid integral outcome of positive support size $n \leq 5$. Normalize \mathbf{w} such that $w_{0,0} = -1$. The degree and positive support size do not change. By Theorem 3.25, the outcome \mathbf{w} induces a reduced statistical model \mathcal{M} in Δ_{n-1} . Then, $\deg(\mathcal{M}) \leq 2(n-1) - 1 = 2n - 3$. Thus, applying Theorem 3.25 let us go back to the outcome \mathbf{w} , and Proposition 3.27 establishes $\deg(\mathbf{w}) \leq 2n - 3$.

For the converse direction, assume Theorem 3.32 holds. Let $n \leq 5$. We want to show Theorem 2.23. Let $\mathcal{M} = (w_k, i_k, j_k)_{k=0}^{n-1} \subset \Delta_{n-1}$ be a one-dimensional discrete statistical model with rational MLE. By Theorem 2.25 we may assume that \mathcal{M} is fundamental. We use Theorem 3.25 to map \mathcal{M} to some valid real outcome $\mathbf{w} = (w_{i,j})$ with $w_{0,0} = -1$. Note that \mathbf{w} is even a *rational* outcome because by Definition 2.14 the weights $(w_k)_{k=0}^{n-1}$ of the fundamental model \mathcal{M} are uniquely determined by the equation $p_0(\theta) + p_1(\theta) + \cdots + p_{n-1}(\theta) - 1 \equiv 0$, and for some $\theta \in [0, 1]$ this equation becomes rational. Next, we use Proposition 3.28 to find some integral valid outcome \mathbf{z} such that $\mathbf{w} = \mu\mathbf{z}$ for some positive scalar $\mu \in \mathbb{Q}_{>0}$. Again, scaling does not affect the degree or size of the positive support. By Proposition the integral valid outcome 3.27 \mathbf{z} has positive support size n . By Theorem 3.32 we have $\deg(\mathbf{z}) \leq 2n - 3$. Thus, $\deg(\mathbf{w}) = \deg(\mu\mathbf{z}) = \deg(\mathbf{z}) \leq 2n - 3$. Hence, the degree of \mathcal{M} is smaller or equal to $2(n-1) - 1$ by Proposition 3.27. We proved Theorem 2.23. \square

Recall that our goal is to demonstrate that only finitely many fundamental statistical models exist in Δ_n for $n \leq 4$. To achieve this, we originally aimed to prove Theorem 2.24 from Chapter 2. However, as established in Chapter 3, we can alternatively prove Theorem 3.32, which serves as an equivalent approach to resolving the problem. We have chosen to focus on Theorem 3.32 because it allows us to address the problem from a combinatorial perspective.

Chapter 4

Supports of Valid Outcomes

Let us devote the remaining chapters to the study of Theorem 3.32, which for the sake of convenience we restate below.

Theorem. *The following upper bound*

$$\deg(\mathbf{w}) \leq 2 \cdot |\text{supp}^+(\mathbf{w})| - 3$$

holds for valid integral outcomes \mathbf{w} with $|\text{supp}^+(\mathbf{w})| \leq 5$.

From now on, valid outcomes \mathbf{w} refer to *integral* valid outcomes in \mathbb{Z}^{V_d} for *finite* $d \in \mathbb{N}$. To show the theorem, we will first study the supports of valid outcomes; knowing that some kinds of supports cannot be the supports of valid outcomes will help us to prove the theorem. For instance, integer configurations that have support in the entries below marked with an $*$ cannot be valid outcomes:

```

.
. .
* . .
. . . .
. . . . .
. . . . * .
* . * . . * *
```

This is the key result of this chapter, see Section 4.4.

4.1 Invertibility Criterion

Let $d \in \mathbb{N}$. One of the most important tools in the study of outcomes is the *invertibility criterion* first introduced in [2]. By Theorem 3.13 we can characterize outcomes as the roots of all Pascal forms on \mathbb{Z}^{V_d} . In the previous chapter we have already found two bases for the

space of Pascal forms, namely $(\text{row}(0), \dots, \text{row}(d))$ and $(\text{col}(0), \dots, \text{col}(d))$ (see Definition 3.16). Let us introduce a *new* basis for the space of Pascal forms.

Definition 4.1. Let $k = 0, \dots, d$ and $\mathbf{e}_k \in \mathbb{R}^{d+1}$ be the k -th unit vector. We define $\text{diag}(k)$ to be the unique Pascal form $\sum c_{i,j} x_{i,j}$ such that $c_{k,d-k} = \mathbf{e}_k$.

Example 4.2. Fix the degree $d = 7$. We visualized $\text{diag}(3)$ by

$$\begin{array}{cccccccc} & & & & & & & & \\ & & & & & & & & \\ & & & & & & & & \\ & & & & & & & & \\ 1 & & 1 & & 1 & & 1 & & \\ 4 & & 3 & & 2 & & 1 & & . \\ 10 & & 6 & & 3 & & 1 & & . & . \\ 20 & & 10 & & 4 & & 1 & & . & . & . \\ 35 & & 15 & & 5 & & 1 & & . & . & . & . \end{array}$$

Proposition 4.3. For all integers $k = 0, \dots, d$ we have:

$$\text{diag}(k) = \sum_{(i,j) \in V_d} \binom{d-i-j}{k-i} x_{i,j}.$$

Note that $\binom{a}{b} = 0$ for $b < 0$ or $b > a$.

Proof. Note that for all $(i,j) \in V_d$ with $i+j = d$ we have $\binom{d-i-j}{k-i} = 1$ if and only if $k = i$, and in all other cases $k \neq i$ the binomial coefficient is zero. Thus, it remains to show that $\sum_{(i,j) \in V_d} \binom{d-i-j}{k-i} x_{i,j}$ is a Pascal form. We have

$$\binom{d-i-j}{k-i} = \binom{d-i-1-j}{k-i-1} + \binom{d-i-j-1}{k-i}.$$

for all $(i,j) \in V_{d-1}$ because $\binom{a+1}{b+1} = \binom{a}{b+1} + \binom{a}{b}$. \square

Proposition 4.4. Let p be a Pascal form on \mathbb{Z}^{V_d} . There exist unique coefficients $\mu_0, \dots, \mu_d \in \mathbb{Z}$ such that $p = \mu_0 \text{diag}(0) + \dots + \mu_d \text{diag}(d)$.

Proof. Let $p = \sum c_{i,j} x_{i,j}$. Choose $\mu_k = c_{k,d-k}$ for $k = 0, \dots, d$. Since p is a Pascal form, the coefficients $c_{i,j}$ satisfy the Pascal recurrence relation. Thus, the coefficients μ_k are uniquely determined. \square

Given some set of vertices $S \subset V_d$ the invertibility criterion uses the diagonal basis $(\text{diag}(0), \dots, \text{diag}(d))$ to determine whether a nonzero outcome with support in S exists.

Definition 4.5. Let $E \subset \{0, \dots, d\}$ and $S \subset V_d$ with $|E| = |S| \neq 0$. The *pairing matrix* of (E, S) is defined as $A_{E,S}^{(d)} := \left[\binom{d-i-j}{k-i} \right]_{k \in E, (i,j) \in S}$.

Example 4.6. Let $d = 2$, $S = \{(1, 1), (0, 0)\}$ and $E = \{0, 1\}$. Then the pairing matrix is

$$A_{E,S}^{(d)} = \begin{bmatrix} \begin{pmatrix} 2 & -1 & -1 \\ 0 & -1 & -1 \\ 2 & -1 & -1 \\ 1 & -1 & \end{pmatrix} & \begin{pmatrix} 2 & -0 & -0 \\ 0 & -2 \\ 2 & -0 & -0 \\ 1 & -2 \end{pmatrix} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}.$$

Now, assume \mathbf{w} is an outcome with support in S . Since it is an outcome, we have $\text{diag}(k)(\mathbf{w}) = 0$ for all $k = 0, 1, 2, 3$. Thus,

$$A_{E,S}^{(d)} \mathbf{w} = \mathbf{0}.$$

We make the following observation: if the matrix $A_{E,S}^{(d)}$ were invertible (it is not for the given example), then we would have $\mathbf{w} = \mathbf{0}$; so in this case the initial configuration $\mathbf{0}$ is the *only* outcome with support in S . This is the invertibility criterion.

Proposition 4.7 (Invertibility Criterion). *Let \mathbf{w} be an outcome with $\text{supp}(\mathbf{w}) \subset S$. If $A_{E,S}^{(d)}$ is invertible, then $\mathbf{w} = \mathbf{0}$.*

Proof by Contraposition. Let $\mathbf{w} \neq \mathbf{0}$. Its support is non-empty. Then, $\mathbf{w}' := (w_{i,j})_{(i,j) \in S} \neq \mathbf{0}$. So, $A_{E,S}^{(d)} \cdot \mathbf{w}' = \mathbf{0}$. The kernel of the pairing matrix is non-trivial. Hence, the pairing matrix $A_{E,S}^{(d)}$ is not invertible. \square

Given a non-zero configuration \mathbf{w} we try to construct sets $S \supset \text{supp}(\mathbf{w})$ and E such that the pairing matrix $A_{E,S}^{(d)}$ is invertible. If we succeed, then \mathbf{w} is *not* an outcome since the initial configuration is the only valid outcome with support in S .

4.2 Divide and Conquer

The invertibility criterion is a powerful tool to determine whether a given configuration is an outcome. However, it is not always easy to find suitable sets S and E such that the pairing matrix is invertible. We will now introduce a method to construct such sets.

Divide

Instead of finding one large set S with $\text{supp}(\mathbf{w}) \subset S$, we divide S into smaller sets S_1, \dots, S_l . These smaller sets S_1, \dots, S_k will be implicitly defined by integers $\lambda_1, \dots, \lambda_l \in \mathbb{N}$ as we will shortly see. We choose $l \in \mathbb{N}$ and integers $\lambda_1, \dots, \lambda_l \in \mathbb{N}$ such that for all $i = 1, \dots, d$ we have

$$\begin{aligned} |S_i| &\in \{0, \lambda_i\} \\ S_i &:= \{(i, j) \in \text{supp}(\mathbf{w}) : i = c_{k-1}, \dots, c_k - 1\} \\ c_i &:= \lambda_1 + \dots + \lambda_i, \\ \lambda_1 + \dots + \lambda_l &= d + 1. \end{aligned}$$

Such decomposition will always work when $|\{(i, j) \in \text{supp}(\mathbf{w}) : i \geq d - k\}| \leq k + 1$ for all $k = 0, \dots, d$. This is because we can always choose λ_1 minimal such that $|S_1| \in \{0, \lambda_1\}$. We repeat this process until $c_l = d + 1$. This decomposition is illustrated in the following example.

Example 4.8. Fix the degree $d = 6$. Assume we have some configuration $\mathbf{w} \in \mathbb{Z}^{V_6}$ with support in the positions marked with an $*$ below.

$$\begin{array}{cccccc} & & & & & \\ & & & & & \\ & & & & & \\ * & . & . & & & \\ & . & . & . & . & \\ & . & . & . & . & . \\ & . & . & . & . & * \\ * & . & * & . & . & * \end{array}$$

The first column contains two non-zero entries. So we see $\lambda_1 = 2$. Then, we conclude that $\lambda_2 = \lambda_3 = \lambda_4 = \lambda_5 = \lambda_6 = 1$; otherwise $|S_i| \notin \{0, \lambda_i\}$ for $i > 1$.

Next with S_i defined, we define for all $i = 1, \dots, l$ the sets

$$E_i := \begin{cases} \{c_{i-1}, \dots, c_i - 1\} & \text{if } |S_i| = \lambda_i, \\ \emptyset & \text{if } |S_i| = 0. \end{cases}$$

We see that $|E_i| = |S_i|$ for all $i = 1, \dots, l$.

Conquer

Given some support S , we divide it into smaller sets S_1, \dots, S_l as described above. We also define sets E_1, \dots, E_l . Write $E := E_1 \cup \dots \cup E_l$.

Proposition 4.9. *We have*

$$A_{E,S}^{(d)} = \begin{bmatrix} A_{E_1,S_1}^{(d)} & 0 & \dots & 0 \\ A_{E_2,S_1}^{(d)} & A_{E_2,S_2}^{(d)} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ A_{E_l,S_1}^{(d)} & A_{E_l,S_2}^{(d)} & \dots & A_{E_l,S_l}^{(d)} \end{bmatrix}.$$

Proof. We have

$$A_{E,S}^{(d)} = \begin{bmatrix} A_{E_1,S_1}^{(d)} & A_{E_1,S_2}^{(d)} & \dots & A_{E_1,S_l}^{(d)} \\ A_{E_2,S_1}^{(d)} & A_{E_2,S_2}^{(d)} & \dots & A_{E_2,S_l}^{(d)} \\ \vdots & \vdots & \ddots & \vdots \\ A_{E_l,S_1}^{(d)} & A_{E_l,S_2}^{(d)} & \dots & A_{E_l,S_l}^{(d)} \end{bmatrix}.$$

Let $x, y = 1, \dots, l$ such that $x < y$. Let $k \in E_x$ and $(i, j) \in S_y$. Then, $k \leq c_x - 1 < c_x \leq c_{y-1} \leq i$; so $k - i < 0$. Thus, $\binom{d-i-j}{k-i} = 0$. This implies that the upper off-diagonal blocks are zero. \square

Corollary 4.10. *The pairing matrix $A_{E,S}^{(d)}$ is invertible if and only if $A_{E_1,S_1}^{(d)}, \dots, A_{E_l,S_l}^{(d)}$ are invertible.*

Corollary 4.11 (Invertibility Criterion, Divide and Conquer). *Let \mathbf{w} be an outcome with $\text{supp}(\mathbf{w}) \subset S$. If $A_{E_1,S_1}^{(d)}, \dots, A_{E_l,S_l}^{(d)}$ are invertible, then $\mathbf{w} = \mathbf{0}$.*

Example 4.12. We continue Example 4.8.

$$\begin{array}{ccccccc} & & & & & & \\ & & & & & & \\ & & & & & & \\ * & . & . & & & & \\ & . & . & . & . & & \\ & . & . & . & . & . & \\ & . & . & . & . & * & . \\ * & . & * & . & . & * & * \end{array}$$

With $\lambda = (2, 1, 1, 1, 1, 1)$ we obtain the following decomposition

$$S_1 = \{(0, 0), (0, 4)\}, S_2 = \{(2, 0)\}, S_3 = \emptyset, S_4 = \{(4, 1)\}, S_5 = \{(5, 0)\}, S_6 = \{(6, 0)\},$$

and

$$E_1 = \{0, 1\}, E_2 = \{2\}, E_3 = \emptyset, E_4 = \{4\}, E_5 = \{5\}, E_6 = \{6\}.$$

Then, the pairing matrix reads

$$A_{E,S}^{(d)} = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ 6 & 2 & 0 & 0 & 0 & 0 \\ * & * & 1 & 0 & 0 & 0 \\ * & * & * & 1 & 0 & 0 \\ * & * & * & * & 1 & 0 \\ * & * & * & * & * & 1 \end{bmatrix},$$

where $*$ stands for arbitrary entries. The matrix is invertible, so no nonzero outcome with support in $S = \{(0, 0), (0, 4), (2, 0), (4, 1), (5, 0), (6, 0)\}$ exists.

4.3 Symmetry

With the Invertibility Criterion we can exclude certain supports from being the supports of valid outcomes. We will now show that the supports of valid outcomes are symmetric with respect to the main diagonal.

Proposition 4.13. *Let $\mathbf{w} = (w_{i,j})_{(i,j) \in V_d}$ be a configuration in \mathbb{Z}^{V_d} . Then \mathbf{w} is an outcome if and only if $\tilde{\mathbf{w}} := (w_{j,i})_{(i,j) \in V_d}$ is an outcome.*

Proof. Observe that

$$\begin{aligned} \text{diag}(k) &= \sum_{(i,j) \in V_d} \binom{d-i-j}{k-i} x_{i,j} \\ &= \sum_{(i,j) \in V_d} \binom{d-i-j}{d-i-j-(k-i)} x_{i,j} \\ &= \sum_{(i,j) \in V_d} \binom{d-i-j}{d-k-j} x_{i,j}. \end{aligned}$$

Let \mathbf{w} be a valid outcome. Then, $\text{diag}(k)(\mathbf{w}) = 0$ for all $k = 0, \dots, d$. Thus,

$$\begin{aligned} \text{diag}(k)(\tilde{\mathbf{w}}) &= \sum_{(i,j) \in V_d} \binom{d-i-j}{k-j} w_{j,i} \\ &= \sum_{(i,j) \in V_d} \binom{d-i-j}{d-k-i} w_{j,i} \\ &= \sum_{(i,j) \in V_d} \binom{d-i-j}{d-k-j} w_{i,j} \\ &= \sum_{(i,j) \in V_d} \binom{d-i-j}{k-i} w_{i,j} \\ &= \text{diag}(k)(\mathbf{w}) = 0 \quad \forall k = 0, \dots, d. \end{aligned}$$

□

Example 4.14. We have already seen that the support below cannot be the support of a valid outcome.

```

.
. .
* . .
. . . .
. . . . .
. . . . * .
* . * . . * *
```

By symmetry, the support below cannot be the support of a valid outcome either.

```

*
* .
. * .
. . . .
* . . . .
. . . . .
* . . . * . .
    
```

Next, we introduce another kind of symmetry. Let $\mathbf{w} = (w_{i,j})$ be a configuration in \mathbb{Z}^{V_d} . We define

$$\mathbf{w} \mapsto \hat{\mathbf{w}} := \left((-1)^{d-j} w_{j,d-i-j} \right)_{(i,j) \in V_d}.$$

Proposition 4.15. *Let $\mathbf{w} = (w_{i,j})_{(i,j) \in V_d}$ be a configuration in \mathbb{Z}^{V_d} . Then \mathbf{w} is an outcome if and only if $\hat{\mathbf{w}} := \left((-1)^{d-j} w_{j,d-i-j} \right)_{(i,j) \in V_d}$ is an outcome.*

Proof. Let $k = 0, \dots, d$. We have

$$\begin{aligned} \text{col}(k)(\hat{\mathbf{w}}) &= (-1)^k \sum_{(i,j) \in V_d} (-1)^j \binom{i}{k-j} (-1)^{d-j} w_{j,d-i-j} \\ &= (-1)^{d-k} \sum_{(i,j) \in V_d} \binom{d-i-j}{k-i} w_{i,j} \\ &= (-1)^{d-k} \text{diag}(k)(\mathbf{w}) \\ &= 0. \end{aligned}$$

This proves the statement. □

The symmetry just introduced can be interpreted in the following way: we define a group action of the symmetry group S_3 on \mathbb{Z}^{V_d} by

$$(12) \cdot \mathbf{w} = \tilde{\mathbf{w}} \quad \text{and} \quad (123) \cdot \mathbf{w} = \hat{\mathbf{w}}.$$

Then, the group actions (12), (13) and (23) can be depicted in Figure 4.1.




Figure 4.1: This illustration is taken from [2]. The left most illustration shows mirroring the configuration with respect to the main diagonal. The middle illustration shows switching the order on the same row while also alternating the signs of the row. The right most illustration shows switching the order on the same column while also alternating the signs of the column.

4.4 Impossible Supports

Now, we show that specific supports cannot be the supports of valid integral outcomes. Hence, the title of this section *Impossible Supports*. For instance, can we have an outcome whose support is only contained in $S = \{(0, 0), (0, i)\}$ for some $i \in \mathbb{N}$? We will show that this is not possible.

Proposition 4.16. *Let $d \in \mathbb{N}$, and $i = 0, \dots, d$. If $S = \{(0, i)\}$ and $E = \{0\}$, then $A_{E,S}^{(d)}$ is invertible.*

Proof. We have $A_{E,S}^{(d)} = [1]$, which is invertible. \square

Proposition 4.17. *Let $d \in \mathbb{N}$. Assume $i, j = 0, \dots, d$ with $i < j$. If $S = \{(0, i), (0, j)\}$ and $E = \{0, 1\}$, then $A_{E,S}^{(d)}$ is invertible.*

Proof. We have $A_{E,S}^{(d)} = \begin{bmatrix} 1 & 1 \\ d-i & d-j \end{bmatrix}$, which is invertible. \square

Proposition 4.18. *Let $d \in \mathbb{N}$. Assume $i, j, k = 0, \dots, d$ with $i < j < k$. If $S = \{(0, i), (0, j), (0, k)\}$ and $E = \{0, 1, 2\}$, then $A_{E,S}^{(d)}$ is invertible.*

Proof. We have

$$A_{E,S}^{(d)} = \begin{bmatrix} \binom{d-i}{0} & \binom{d-j}{0} & \binom{d-k}{0} \\ \binom{d-i}{1} & \binom{d-j}{1} & \binom{d-k}{1} \\ \binom{d-i}{2} & \binom{d-j}{2} & \binom{d-k}{2} \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 \\ d-i & d-j & d-k \\ \frac{(d-i)(d-i-1)}{2} & \frac{(d-j)(d-j-1)}{2} & \frac{(d-k)(d-k-1)}{2} \end{bmatrix}.$$

We substitute

$$\begin{aligned} x &= d - i, \\ y &= d - j, \\ z &= d - k. \end{aligned}$$

Then, we see

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 2 \end{bmatrix} A_{E,S}^{(d)} = \begin{bmatrix} 1 & 1 & 1 \\ x & y & z \\ x^2 & y^2 & z^2 \end{bmatrix}.$$

The matrix on the right-hand side is invertible because it is a Vandermonde matrix. Thus, the pairing matrix $A_{E,S}^{(d)}$ is invertible. \square

Proposition 4.19. *Let $d \in \mathbb{N}$. Assume $i, j = 0, \dots, d$ with $i < j$. Moreover, let $k = 0, \dots, d - 1$. If $S = \{(0, i), (0, j), (1, k)\}$, $E = \{0, 1, 2\}$, and $i + j \neq 2k + 1$, then $A_{E,S}^{(d)}$ is invertible.*

Proof. We have

$$A_{E,S}^{(d)} = \begin{bmatrix} \binom{d-i}{0} & \binom{d-j}{0} & \binom{d-k-1}{-1} \\ \binom{d-i}{d-i} & \binom{d-j}{d-j} & \binom{d-k-1}{0} \\ \binom{d-i}{1} & \binom{d-j}{1} & \binom{d-k-1}{d-k-1} \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 \\ d-i & d-j & d-k-1 \\ \frac{(d-i)(d-i-1)}{2} & \frac{(d-j)(d-j-1)}{2} & \frac{(d-k-1)(d-k-2)}{2} \end{bmatrix}.$$

We substitute

$$\begin{aligned} x &= d - i, \\ y &= d - j, \\ z &= d - k - 1. \end{aligned}$$

Then, we see

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 2 \end{bmatrix} A_{E,S}^{(d)} \begin{bmatrix} 1 & 1 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{x-y} & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ x & 1 & 1 \\ x^2 & x+y & 2z+1 \end{bmatrix}.$$

We see that the determinant is nonzero because $x + y \neq 2z + 1$ by $i + j \neq 2k + 1$. \square

Remark 4.20. Without loss of generality, we may assume

$$\begin{aligned} S &\subset \{(i, j) \in V_d \mid i < |S|\} \\ E &= \{0, 1, \dots, |S| - 1\} \end{aligned}$$

because the matrices $A_{E,S}^{(d)} = A_{E-\rho, S-\rho}^{(d-\rho)}$ are equal, where $\rho := \min \{E \cup \{i \mid (i, j) \in S\}\}$ and $E-\rho := \{(i-\rho, j) \mid (i, j) \in E\}$. This assumption allows us to apply the previous propositions to more general S and E .

Example 4.21. Assume we have a configuration with support $S = \{(0, i), (0, j), (0, k)\}$ for $0 \leq i < j < k \leq d$. By Proposition 4.18, we know that no valid integral nonzero outcome can have this support.

Now, let us consider a configuration \mathbf{w} with support $\text{supp}(\mathbf{w}) \subset S$ such that S can be decomposed into S_1, \dots, S_l as described before. Let $\ell = 1, \dots, l$. If $S_\ell = \{(x, i), (x, j), (x, k)\}$ for $0 \leq i < j < k \leq d$ and $x \in \mathbb{N}$, then $\mathbf{w} = \mathbf{0}$ by Proposition 4.18 and the previous comment on the generality of S and E .

For instance, this configuration is not an outcome

$$\begin{array}{ccccccc}
 * & & & & & & \\
 . & . & & & & & \\
 . & . & * & & & & \\
 . & . & . & . & & & \\
 . & . & . & * & . & & \\
 . & . & . & . & . & . & \\
 . & . & . & * & . & . & . \\
 * & . & . & * & . & . & . & .
 \end{array}$$

where $*$ denotes a non-zero entry. This is because for $\lambda = (3, 3, 1, 1)$ we have $S_2 = \{(3, 0), (3, 1), (3, 3)\}$. Similarly, these configurations are not outcomes

$$\begin{array}{ccccccc}
 * & & & & & & \\
 . & . & & & & & \\
 . & . & * & & & & \\
 . & . & . & . & & & \\
 . & . & . & . & . & . & \\
 . & . & . & * & . & . & \\
 . & . & . & * & . & . & . \\
 * & . & . & * & . & . & . & .
 \end{array}$$

$$\begin{array}{ccccccc}
 * & & & & & & \\
 . & . & & & & & \\
 . & . & * & & & & \\
 . & . & . & . & & & \\
 . & . & . & . & . & & \\
 . & . & . & . & . & * & \\
 . & . & . & . & . & * & . \\
 * & . & . & . & . & * & . & .
 \end{array}$$

Of course, there are many more examples of impossible supports.

Chapter 5

Valid Outcomes of Positive Support Size One, Two, and Three

All the tools are ready to show the following three theorems, which were first proved in [2].

Theorem 5.1. *No valid integral outcomes of positive support size one exists.*

Theorem 5.2. *For valid integral outcomes \mathbf{w} with $|\text{supp}^+(\mathbf{w})| = 2$ we have $\deg(\mathbf{w}) = 1$.*

Theorem 5.3. *For valid integral outcomes \mathbf{w} with $|\text{supp}^+(\mathbf{w})| = 3$ we have $\deg(\mathbf{w}) \leq 3$.*

This proves our Main Theorem 3.32 for the case of positive support size three or less, i.e.

$$\deg(\mathbf{w}) \leq 2 \cdot |\text{supp}^+(\mathbf{w})| - 3$$

for all valid integral outcomes \mathbf{w} with $|\text{supp}^+(\mathbf{w})| \leq 3$.

We start with the proof of the first theorem.

Proof of Theorem 5.1. Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be a valid integral outcome. Since it is valid, we either have an empty negative support or a negative support that only contains $(0, 0)$. If the negative support is empty, then $\mathbf{w} = \mathbf{0}$ by Proposition 3.29. Hence, we assume $w_{0,0} < 0$.

Now, consider the Pascal form $\text{diag}(0) = \sum c_{i,j} x_{i,j}$. We have $c_{0,0} = c_{0,1} = \dots = c_{0,d} = 1$ and $c_{i,j} = 0$ for everything else. Similarly, we have for the Pascal form $\text{diag}(d) = \sum c'_{i,j} x_{i,j}$ that $c'_{i,0} = 1$ and $c'_{i,j} = 0$ for everything else. Since outcomes are roots of Pascal forms, we have

$$\text{diag}(0)(\mathbf{w}) = \text{diag}(d)(\mathbf{w}) = 0.$$

Since $w_{0,0} < 0$ we must have $w_{0,j} > 0$ and $w_{i,0} > 0$ for some $i, j > 0$. Hence, \mathbf{w} has positive support size at least two. \square

Next, we prove the second theorem.

Proof of Theorem 5.2. Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be an integral outcome with positive support size two and degree d . By the previous proof, we see that

$$\text{supp}^+(\mathbf{w}) = \{(0, j), (i, 0)\}.$$

Without loss of generality, we assume $i = d$. We want to show that $j = d$. Consider the Pascal form $\text{row}(d) = \sum c_{i,j} x_{i,j}$, which has only nonzero coefficients $c_{i,j}$ for $i + j = d$.

- If the degree d is odd, we have $c_{d,0} = 1$ and $c_{0,d} = -1$. Since $\text{row}(d)(\mathbf{w}) = 0$, we must have $j = d$.
- If the degree d is even, we have $c_{d,0} = c_{0,d} = 1$. Thus, $\text{row}(d)(\mathbf{w}) \neq 0$ for all $j = 0, \dots, d$. Hence, valid outcomes with positive support size two do not exist for even degrees.

From now on, we assume $\text{supp}^+(\mathbf{w}) = \{(0, d), (d, 0)\}$. We could further assume that the degree d is odd, but we do not need it.

For sake of contradiction, let $d \geq 2$. Then, we can divide the support

$$\text{supp}(\mathbf{w}) = \{(0, 0), (0, d), (d, 0)\},$$

via $\lambda = (2, 1, \dots, 1)$ as in Chapter 4.2 to obtain $S_1 = \{(0, 0), (0, d)\}$, $S_k = \emptyset$, and $S_l = \{(d, 0)\}$. By Proposition 4.17, the pairing matrix induced by S_1 and $E_1 = \{0, 1\}$ is invertible. For S_l we apply Proposition 4.16 and Remark 4.20 to get that the induced pairing matrix is invertible. By Corollary 4.11, the outcome \mathbf{w} is zero, which has an empty positive support. This is a contradiction to the assumption that the positive support size is two. Hence, the degree d equals one. \square

Example 5.4. The previous theorem shows that the only valid integral outcomes with positive support size two are multiples of

$$\begin{pmatrix} 1 \\ -1 & 1 \end{pmatrix}.$$

It remains to prove Theorem 5.3. For that consider the following lemma which characterizes the possible supports of valid integral outcomes with positive support size three.

Proposition 5.5. *Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be a valid integral outcome of degree d . If the positive support size of \mathbf{w} is three, then one of the following holds:*

1. We have $\text{supp}(\mathbf{w}) = \{(0, 0), (d, 0), (0, d), (i, j)\}$ for some $i, j > 0$ with $i + j < d$.
2. We have $\text{supp}(\mathbf{w}) = \{(0, 0), (d, 0), (0, d), (i, d - i)\}$ for some $i = 1, \dots, d - 1$.
3. We have $\text{supp}(\mathbf{w}) = \{(0, 0), (d, 0), (0, d), (i, 0)\}$ for some $i = 1, \dots, d - 1$.

4. We have $\text{supp}(\mathbf{w}) = \{(0, 0), (d, 0), (0, d), (0, i)\}$ for some $i = 1, \dots, d-1$.
5. We have $\text{supp}(\mathbf{w}) = \{(0, 0), (d, 0), (0, e), (d-f, f)\}$ for some $e, f = 1, \dots, d-1$.
6. We have $\text{supp}(\mathbf{w}) = \{(0, 0), (0, d), (e, 0), (d-f, f)\}$ for some $e, f = 1, \dots, d-1$.

Proof. Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be a valid integral outcome of degree d . Assume $\{(0, 0), (d, 0), (0, d)\} \subset \text{supp}(\mathbf{w})$. Clearly, statement 1, 2, 3, or 4 must hold.

So assume $(0, d) \notin \text{supp}(\mathbf{w})$ and $(d, 0) \notin \text{supp}(\mathbf{w})$. As in the proof of Theorem 5.1, consider the Pascal form $\text{diag}(0) = \sum c_{i,j} x_{i,j}$. We have $c_{0,\cdot} = \mathbf{1}$ and $c_{i,j} = 0$ for everything else. Similarly, we have for the Pascal form $\text{diag}(d) = \sum c'_{i,j} x_{i,j}$ that $c'_{\cdot,0} = \mathbf{1}$ and $c'_{i,j} = 0$ for everything else. Since outcomes are roots of Pascal forms, we have $\text{diag}(0)(\mathbf{w}) = \text{diag}(d)(\mathbf{w}) = 0$. Due to $w_{0,0} < 0$, we conclude $w_{0,j} > 0$ and $w_{i,0} > 0$ for some $i, j > 0$. Thus, we have

$$\{(i, 0), (0, j)\} \subset \text{supp}(\mathbf{w})$$

for some $i, j = 1, \dots, d-1$ using the assumption $(0, d) \notin \text{supp}(\mathbf{w})$ and $(d, 0) \notin \text{supp}(\mathbf{w})$. Since \mathbf{w} is of degree d , there exists $w_{k,d-k} > 0$ for some $k = 1, \dots, d-1$. However, $\text{row}(d)(\mathbf{w}) = 0$ implies that there must be some $w_{h,d-h} > 0$ for some $h \neq k$; this h cannot equal 0 or d . Thus, the positive support size of \mathbf{w} is at least four, which is a contradiction. Hence, we must have $(d, 0) \in \text{supp}(\mathbf{w})$ or $(0, d) \in \text{supp}(\mathbf{w})$.

Let $(d, 0) \in \text{supp}(\mathbf{w})$ and $(e, 0) \in \text{supp}(\mathbf{w})$ for some $e = 1, \dots, d-1$. Now using the same argument as before, there must exist some $w_{f,d-f} > 0$ for some $f = 1, \dots, d-1$; otherwise $\text{row}(d)(\mathbf{w}) > 0$ which is a contradiction since \mathbf{w} is a root of all Pascal forms. This proves statement 5.

The proof for statement 6 is analogous. □

Knowing the possible supports of valid integral outcomes with positive support size three, we apply the Invertibility Criterion 4.11 to each possible support to prove Theorem 5.3.

Proposition 5.6. *Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be a valid integral outcome of degree d . If $\text{supp}(\mathbf{w}) = \{(0, 0), (d, 0), (0, d), (i, j)\}$ for some $i, j > 0$ with $i + j < d$, then $d = 3$ and $(i, j) = (1, 1)$.*

Proof. Let $i > 1$. Choose $\lambda = (2, 1, \dots, 1)$. Then, $E_1 = \{0, 1\}$, $S_1 = \{(0, 0), (0, d)\}$, $E_{i-1} = \{i\}$, $S_{i-1} = \{(i, j)\}$, $E_{d-1} = \{d\}$, $S_{d-1} = \{(d, 0)\}$ and, $E_k = S_k = \emptyset$ for all $k \in \{1, \dots, d-1\} \setminus \{1, i-1, d-1\}$. The pairing matrices $A_{E_n, S_n}^{(d)}$ are all invertible for $n = 1, \dots, d-1$. Hence, the pairing matrix $A_{\{0,1,i,d\}, \text{supp}(\mathbf{w})}^{(d)}$ is also invertible. By the Invertibility Criterion, \mathbf{w} is the zero configuration, which is a contradiction. Thus, we have $i = 1$.

Now, we assume $j > 1$. The configuration $\tilde{\mathbf{w}} = (w_{ji})_{(i,j) \in V_d}$ is an outcome by Proposition 4.13 because \mathbf{w} is an outcome. Then $\tilde{\mathbf{w}}$ has support $\{(0, 0), (d, 0), (0, d), (1, \cdot)\}$ by the previous argument. However, then we have $j = 1$ which is a contradiction. So, we have $j = 1$.

Finally, we need to show that the degree d equals three. For the sake of contradiction, assume $d > 3$. Then, we can choose $\lambda = (3, 1, \dots, 1)$. We obtain $E_1 = \{0, 1, 2\}$ and

$S_1 = \{(0, 0), (0, d), (1, 1)\}$. By Proposition 4.19 this pairing matrix $A_{E_1, S_1}^{(d)}$ is invertible. The other relevant pairing matrix $A_{\{d\}, \{(d, 0)\}}^{(d)}$ is also invertible. Thus, the pairing matrix $A_{\{0, 1, 2, d\}, \text{supp}(\mathbf{w})}^{(d)}$ is invertible. By the Invertibility Criterion, the configuration \mathbf{w} is the zero configuration, which is a contradiction. Hence, we have $d = 3$. \square

Proposition 5.7. *Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be a valid integral outcome of degree d . Assume the outcome \mathbf{w} satisfies one of the following conditions:*

1. $\text{supp}(\mathbf{w}) = \{(0, 0), (d, 0), (0, d), (i, d - i)\}$ for some $i = 1, \dots, d - 1$,
2. $\text{supp}(\mathbf{w}) = \{(0, 0), (d, 0), (0, d), (i, 0)\}$ for some $i = 1, \dots, d - 1$,
3. $\text{supp}(\mathbf{w}) = \{(0, 0), (d, 0), (0, d), (0, i)\}$ for some $i = 1, \dots, d - 1$.

Then, $d = 2$ and $i = 1$ hold.

Proof. Assume $d > 2$. Let \mathbf{w} satisfy the third condition. Choose $\lambda = (3, 1, \dots, 1)$. Then, apply Proposition 4.18. So, we have that the pairing matrix is invertible. So, $\mathbf{w} = \mathbf{0}$ which is a contradiction. Thus, $d = 2$. By symmetry we have the same result for the second condition.

We want to show $d = 2$ for all outcomes \mathbf{w} satisfying the first condition. Let \mathbf{w}' satisfy the second condition. Then $\mathbf{w} = (123)\mathbf{w}'$ holds. Assume $d > 2$. By Proposition 4.15, we have found an outcome \mathbf{w}' of degree at least three. This contradicts Proposition 5.7 that we have just shown for the second condition. Thus, $d = 2$ holds.

Finally, we have $i = 1$ because $i = 1, \dots, d - 1$ and $d = 2$. \square

Proposition 5.8. *Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be a valid integral outcome of degree d . Assume the outcome \mathbf{w} satisfies one of the following conditions:*

1. $\text{supp}(\mathbf{w}) = \{(0, 0), (d, 0), (0, e), (d - f, f)\}$ for some $e, f = 1, \dots, d - 1$.
2. $\text{supp}(\mathbf{w}) = \{(0, 0), (0, d), (e, 0), (d - f, f)\}$ for some $e, f = 1, \dots, d - 1$.

Then, $d = 2$ and $e = f = 1$ hold.

Proof. By Proposition 4.13, it suffices to show the statement for outcomes \mathbf{w} satisfying the first condition.

Let $d > 2$. If $f = d - 1$, then choose $\lambda = (3, 1, \dots, 1)$. This allows us to apply Proposition 4.19 because $0 + e \neq 2d - 1$ for $d > 1$. So, $\mathbf{w} = \mathbf{0}$ which is a contradiction. Thus, we have $f < d - 1$. Then, we can choose $\lambda = (2, 1, \dots, 1)$. Use Proposition 4.17 to get that $\mathbf{w} = \mathbf{0}$. This is a contradiction. Hence, we have $d = 2$.

Let $d = 2$. Then, we have $e = f = 1$ by definition of e and f . \square

Finally, we can prove Theorem 5.3.

Proof of Theorem 5.3. Use Proposition 5.5. For each case, either apply Proposition 5.6, Proposition 5.7, or Proposition 5.8 to show that the degree d equals two or three. \square

Chapter 6

Constraints on Supports of Valid Outcomes

We want to show in the next chapter that for all valid integral outcomes \mathbf{w} with $|\text{supp}^+(\mathbf{w})| = 4$ we have $\deg(\mathbf{w}) \leq 2 \cdot |\text{supp}^+(\mathbf{w})| - 3 = 5$. For that goal we need new tools aside from the *Invertibility Criterion*. We will introduce the *Hyperfield Criterion* and *Contractions* to achieve it.

6.1 Hyperfield Criterion

In this section, we develop the *Hyperfield Criterion* which allows us to interpret Pascal forms as constraints on the support of valid outcomes.

Let us define the sign hyperfield. For some set A , the set 2^A denotes the power set of A .

Definition 6.1. Let $H := \{-1, 0, 1\}$. We define the addition $+: H \times H \rightarrow 2^H \setminus \{\emptyset\}$ on H as follows

$$0 + x = \{x\} \quad \forall x \in H, \quad 1 + 1 = \{1\}, \quad 1 + (-1) = H, \quad (-1) + (-1) = \{-1\}.$$

Multiplication $\times: H \times H \rightarrow H$ is defined as usual. We call H the *sign hyperfield*.

Often, for singleton sets $\{x\}$, we will write x instead of $\{x\}$. So,

$$1 + 1 = 1 \quad \text{or} \quad (-1) + 0 = -1.$$

Remark 6.2. The tuple $(H, +, \cdot, 0, 1)$ is called a *hyperfield*. A hyperfield satisfies the following properties:

1. The maps $+$ and \cdot are symmetric;
2. $(H \setminus \{0\}, \cdot, 1)$ is a group;

3. $0 \cdot x = 0$ and $0 + x = x$ hold for all $x \in H$;
4. $\bigcup_{q \in x+y} (q + z) = \bigcup_{q \in x+y} (x + q)$ hold for all $x, y, z \in H$;
5. $a \cdot (x + y) = (a \cdot x) + (a \cdot y)$ hold for all $a, x, y \in H$.
6. An inverse element $y \in H$ exists for every $x \in H$ such that the set $x + y$ contains 0. This inverse element y is unique for every x and is denoted by $-x$.

Refer to [2] Section 6.1 or [1] for more details.

Next, we define polynomials over the sign hyperfield.

Definition 6.3. A polynomial in n variables x_1, \dots, x_n over H is a formal sum

$$f = \sum_{\mathbf{k} \in \mathbb{Z}_{\geq 0}^n} \lambda_{\mathbf{k}} \mathbf{x}^{\mathbf{k}}, \quad \lambda_{\mathbf{k}} \in H,$$

where only a finite number of coefficients $\lambda_{\mathbf{k}}$ are non-zero, and $\mathbf{x}^{\mathbf{k}} = x_1^{k_1} \cdots x_n^{k_n}$. The set of all polynomials in n variables over H is denoted by $H[x_1, \dots, x_n]$.

Let $\mathbf{x} \in H$. Then, we define

$$f(\mathbf{x}) := \sum_{\mathbf{k} \in \mathbb{Z}_{\geq 0}^n} \lambda_{\mathbf{k}} \mathbf{x}^{\mathbf{k}} \in H.$$

We say that f *vanishes* at $\mathbf{x} \in H$ if $0 \in f(\mathbf{x})$. In this case, \mathbf{x} is a *hyperfield root* of f .

Any *real* polynomial can be turned into a polynomial over the sign hyperfield by replacing the coefficients with elements of H . We can then evaluate the polynomial at any point in H .

Definition 6.4. Let $f = \sum \lambda_{\mathbf{k}} \mathbf{x}^{\mathbf{k}} \in \mathbb{R}[\mathbf{x}]$ be a polynomial over \mathbb{R} . We call

$$\text{sign}(f) := \sum_{\mathbf{k} \in \mathbb{Z}_{\geq 0}^n} \text{sign}(\lambda_{\mathbf{k}}) \mathbf{x}^{\mathbf{k}} \in H[\mathbf{x}]$$

the polynomial over H induced by f .

For sake of simplicity, we also write for any real vector $\mathbf{w} \in \mathbb{R}^n$:

$$\text{sign}(\mathbf{w}) := (\text{sign}(w_1), \dots, \text{sign}(w_n)).$$

Example 6.5. Let $d = 5$. Consider the Pascal forms on \mathbb{Z}^{V_d} generated by $\text{diag}(0)$, $\text{diag}(1)$, $\text{diag}(2)$, $\text{diag}(3)$, $\text{diag}(4)$ and $\text{diag}(5)$. The polynomial over H induced by these forms can be depicted as follows:

Dots represent zero, and $+$ represents one. Similarly, consider $\text{col}(0), \dots, \text{col}(5)$:

A minus sign $-$ represents -1 . For $\text{row}(0), \dots, \text{row}(5)$ we have

Definition 6.6. A hyperfield Pascal form is just a polynomial over H induced by a Pascal form.

The reason we introduced the sign hyperfield is that it allows us to neglect the concrete values of the coefficients of a polynomial and focus on their signs. This makes reasoning about roots easier, which is helpful since we saw in earlier chapters that chipsplitting outcomes are roots of Pascal forms.

Proposition 6.7. *Let $f \in \mathbb{R}[\mathbf{x}]$ be a real polynomial. Let $\mathbf{w} \in \mathbb{R}^n$ be a root of f . Then, $\text{sign}(\mathbf{w})$ is a root of $\text{sign}(f)$.*

Proof. Define $\mathbf{s} := \text{sign}(\mathbf{w})$. Write $f = \sum \lambda_{\mathbf{k}} \mathbf{x}^{\mathbf{k}}$ with real coefficients $\lambda_{\mathbf{k}}$. If $\lambda_{\mathbf{k}} \mathbf{w}^{\mathbf{k}} = 0$ for all $\mathbf{k} \in \mathbb{Z}_{\geq 0}^n$, then clearly the sign of $\lambda_{\mathbf{k}} \mathbf{w}^{\mathbf{k}}$ is zero; hence the sign of f is the singleton set $\{0\}$ when evaluated at \mathbf{s} . So, \mathbf{s} is a root of $\text{sign}(f)$.

Now, suppose that there exists some $\mathbf{k} \in \mathbb{Z}_{\geq 0}^n$ such that $\lambda_{\mathbf{k}} \mathbf{w}^{\mathbf{k}} \neq 0$. Assume $\lambda_{\mathbf{k}} \mathbf{w}^{\mathbf{k}} > 0$. Then, there also exists some $\mathbf{j} \in \mathbb{Z}_{> 0}^n$ such that we have $\lambda_{\mathbf{j}} \mathbf{w}^{\mathbf{j}} < 0$; otherwise $f(\mathbf{w}) > 0$ which

is a contradiction to \mathbf{w} being a root of f . Thus, $\text{sign}(f)(\mathbf{s})$ has summands of both signs, and hence $\text{sign}(f)(\mathbf{s}) = H$. So $0 \in \text{sign}(f)(\mathbf{s})$ holds. Therefore, \mathbf{s} is a root of $\text{sign}(f)$. \square

Taking the contrapositive of the above proposition, we get the *Hyperfield Criterion* which was first presented by Bik and Marigliano in [2].

Proposition 6.8 (Hyperfield Criterion). *Let $\mathbf{s} = (s_{i,j})_{(i,j) \in V_d} \in H^{V_d}$. Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be a chipsplitting configuration with $\text{sign}(\mathbf{w}) = \mathbf{s}$. If \mathbf{s} is not a root of a hyperfield Pascal form, then \mathbf{w} is not a chipsplitting outcome.*

Proof. Follows from Proposition 6.7 and Theorem 3.13. \square

We call a vector $\mathbf{s} \in H^{V_d}$ a *sign configuration* or *hyperfield configuration*. For completeness, we state standard definitions for sign configurations $\mathbf{s} \in H^{V_d}$ similar to Definition 3.6.

Definition 6.9. Let $\mathbf{s} \in H^{V_d}$ be a sign configuration. We define the following:

1. The positive support is defined as $\text{supp}^+(\mathbf{s}) := \{(i, j) \in V_d \mid s_{i,j} = 1\}$.
2. The negative support is defined as $\text{supp}^-(\mathbf{s}) := \{(i, j) \in V_d \mid s_{i,j} = -1\}$.
3. The support is defined as $\text{supp}(\mathbf{s}) := \text{supp}^+(\mathbf{s}) \cup \text{supp}^-(\mathbf{s})$.
4. The degree of \mathbf{s} is defined as $\deg(\mathbf{s}) := \max \{i + j \mid (i, j) \in \text{supp}(\mathbf{s})\}$.
5. We call \mathbf{s} *valid* if its support is empty or $\text{supp}^-(\mathbf{s}) = \{(0, 0)\}$.

Lemma 6.10. *Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be a chipsplitting configuration. Then, the following statements hold:*

1. $\text{supp}^+(\text{sign}(\mathbf{w})) = \text{sign}^+(\mathbf{w})$,
2. $\text{supp}^-(\text{sign}(\mathbf{w})) = \text{sign}^-(\mathbf{w})$,
3. $\deg(\text{sign}(\mathbf{w})) = \deg(\mathbf{w})$.

Proof. Follows from the definitions. \square

To make use of the Hyperfield Criterion, we investigate hyperfield forms induced by the Pascal forms $\text{col}(k)$, $\text{row}(k)$, and $\text{diag}(k)$.

Proposition 6.11. *Let $k = 0, \dots, d$. Define*

$$\begin{aligned} A_k^+ &:= \{(i, j) \in V_d \mid j = 0, \dots, k \text{ and } i = k - j, \dots, d - j \text{ with } j \equiv k \pmod{2}\} \\ A_k^- &:= \{(i, j) \in V_d \mid j = 0, \dots, k \text{ and } i = k - j, \dots, d - j \text{ with } j \not\equiv k \pmod{2}\} \end{aligned}$$

Then, the following statements hold:

1. We have

$$\text{sign}(\text{diag}(k)) = \sum_{i=0}^k \sum_{j=0}^{d-k} x_{i,j}.$$

2. We have

$$\text{sign}(\text{col}(k)) = \sum_{(i,j) \in A_k^+} x_{i,j} - \sum_{(i,j) \in A_k^-} x_{i,j}.$$

3. We have

$$\text{sign}(\text{row}(k)) = \sum_{(i,j) \in A_k^+} x_{j,i} - \sum_{(i,j) \in A_k^-} x_{j,i}.$$

Proof. The first statement follows directly from Proposition 4.3 since $i \leq k$ and $d-i-j \geq k-i$ must hold for the binomial coefficient to be non-zero. The second and third statement follow similarly from Proposition 3.18. \square

Proposition 6.12. *Let $\mathbf{s} \in H^{V_d}$ be a valid nonzero sign configuration. The following statements hold:*

1. *Let $k = 0, \dots, d$. If $0 \in \text{sign}(\text{diag}(k))(\mathbf{s})$, then $\text{sign}(\text{diag}(k))(\mathbf{s}) = H$.*
2. *If $0 \in \text{sign}(\text{col}(k))(\mathbf{s})$ for all $k = 0, \dots, d$, then $\text{sign}(\text{col}(k))(\mathbf{s}) = H$.*
3. *If $0 \in \text{sign}(\text{row}(k))(\mathbf{s})$ for all $k = 0, \dots, d$, then $\text{sign}(\text{row}(k))(\mathbf{s}) = H$.*

Proof. We see that \mathbf{s} has at least degree $d \geq 1$ since it is nonzero and valid. All $s_{i,j}$ equal one for $i + j > 0$, and there exists $s_{k,d-k} = 1$ for some $k = 0, \dots, d$.

1. Assume $\text{sign}(\text{diag}(k))(\mathbf{s})$ contains zero. By Proposition 6.11, we have

$$0 \in \text{sign}(\text{diag}(k))(\mathbf{s}) = \sum_{i=0}^k \sum_{j=0}^{d-k} s_{i,j}.$$

We know that $s_{0,0}$ is minus one. So, we have $s_{i,j} = 1$ for some i, j with $i + j > 0$. Thus, $\text{sign}(\text{diag}(k))(\mathbf{s}) = H$.

2. First note that $\text{col}(0) = \text{diag}(d)$. So, the case $k = 0$ is proven. Let $k > 0$. We start with $k = d$. Then, the union of A_d^+ and A_d^- consists exactly of vertices of degree d . Since $\text{sign}(\text{col}(d))(\mathbf{s}) = \sum_{(i,j) \in A_d^+} s_{i,j} - \sum_{(i,j) \in A_d^-} s_{i,j}$ contains zero, we have $s_{i,j} = 1$ for some $(i, j) \in A_d^+$, and $s_{i',j'} = -1$ for some $(i', j') \in A_d^-$. Hence, $\text{sign}(\text{col}(d))(\mathbf{s}) = H$.

Let $k = d - 1$. Then, $s_{i,j} = 1$ for some $(i, j) \in A_{k+1}^+$, and $s_{i',j'} = -1$ for some $(i', j') \in A_{k+1}^-$. Note that $A_{k+1}^- \subset A_k^+$ by definition. Since $\text{sign}(\text{col}(k))(\mathbf{s}) = \sum_{(i,j) \in A_k^+} s_{i,j} - \sum_{(i,j) \in A_k^-} s_{i,j}$ contains zero, we have $s_{i'',j''} = -1$ for some $(i'', j'') \in A_k^-$. Hence, $\text{sign}(\text{col}(k))(\mathbf{s}) = H$.

Repeat this argument for $k = d - 2, \dots, 1$ to show that $\text{sign}(\text{col}(k))(\mathbf{s}) = H$.

3. The proof is analogous to the previous case.

□

Corollary 6.13. *Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be a valid outcome. Then,*

$$\text{sign}(p)(\text{sign}(\mathbf{w})) = H \quad \text{for all } p \in \{\text{diag}(k), \text{col}(k), \text{row}(k) \mid k = 0, \dots, d\}.$$

Proof. This follows from Theorem 3.13, Proposition 6.7, and Proposition 6.12. □

Example 6.14. Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be a valid outcome of degree $d = 5$. By the previous corollary and Example 6.5, we know that the outcome \mathbf{w} has at least one positive entry $w_{i,j}$ in each of the following marked areas + because $w_{0,0} < 0$:

+
+ .	+ +
+ . .	+ + .	+ + +
+ . . .	+ + . .	+ + + .	+ + + +
+	+ + . . .	+ + + . .	+ + + + .	+ + + + +
+	+ +	+ + + . . .	+ + + + . .	+ + + + + .	+ + + + + +

Moreover, for each triangle below the outcome \mathbf{w} must have some $w_{i,j} > 0$ for one vertex (i, j) in the plus area + and $w_{i',j'} > 0$ for another vertex (i', j') in the minus area − because $\text{sign}(\text{col})(\text{sign}(\mathbf{w})) = H$.

.	.	.	.	+
.	+ +	. −
.	+ + +	. − −	. . +
. . . .	+ + + +	. − − −	. . + +	. . . −
+ + + + +	. − − − −	. . + + +	. . . − − +
. − − − − −	. . + + + +	. . . − − − + + −

Similarly, the statement holds for $\text{sign}(\text{row})$:

−	+	−	+	−
− +	+ −	− +	+ −	. +
− + .	+ − +	− + −	. − +	. . −

- + . . . + - + . . . + - + . . . + - . . . +
 - + - + - + . . . - + -
 . + + + + +

The above example demonstrates that we can view Corollary 6.13 as constraints on the support of a valid outcome \mathbf{w} . Configurations that do not satisfy these constraints are not valid outcomes.

Corollary 6.15. *Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be a valid outcome of degree $d \geq 1$. Then, all of the following constraints hold:*

1. *For all $k = 0, \dots, d$, the positive support of \mathbf{w} contains the vertex (i, j) for at least one $i = 0, \dots, k$ and $j = 0, \dots, d - k$.*
2. *For all $k = 1, \dots, d$, the positive support of \mathbf{w} contains at least one $(i, j) \in A_k^+$ and $(i', j') \in A_k^-$.*
3. *For all $k = 1, \dots, d$, the positive support of \mathbf{w} contains at least one (i, j) and (i', j') from $(j, i) \in A_k^+$ and $(j', i') \in A_k^-$.*

We will use these constraints to efficiently compute all outcomes of degree some d .

6.2 Solving Homogeneous Hyperfield Linear Systems

We deviate from the approach by Bik and Marigliano in [2] by spending some considerable time in solving homogeneous hyperfield linear systems. It will help us path the way for some breakthrough for the efficient computation of valid outcomes.

Fix some degree d . Due to the Hyperfield Criterion, we are interested in valid configurations $\mathbf{w} \in \mathbb{Z}^{V_d}$ which satisfy

$$0 \in \text{sign}(p)(\text{sign}(\mathbf{w}))$$

for all $p \in \{\text{diag}(k), \text{row}(k), \text{col}(k)\}_{k=0}^d$. These configurations are potential valid outcomes. Let us consider how we can efficiently compute such valid configurations.

Problem: Given a set of linear forms $A = \{p_1, \dots, p_k\}$, compute the solution set $V(A) := \{\mathbf{x} \in H^{V_d} : 0 \in \text{sign}(p_i)(\mathbf{x}) \ \forall i = 1, \dots, k\}$.

We further simplify the problem by only considering configurations \mathbf{x} with $\text{supp}^-(\mathbf{x}) = \{(0, 0)\}$ and $|\text{supp}^+(\mathbf{x})| = n$ for some fixed $n \in \mathbb{N}$.

Problem: Given a set of linear forms $A = \{p_1, \dots, p_k\}$, compute the solution set $S_n(A) := V(A) \cap \{x \in H^{V_d} : \text{supp}^-(x) = \{(0, 0)\}, |\text{supp}^+(x)| = n\}$.

Note that $S_n(A)$ is a superset of valid outcomes of positive support size n , which will be useful in finding all valid outcomes.

A Naive Approach

To compute $S_n(A)$ a simple brute force algorithm can be used; just iterate over all positive support size n supports and check if they are hyperfield roots of some hyperfield Pascal basis.

Algorithm 2 Brute Force Algorithm

Require: Positive support size n , a set of linear forms $A = \{p_1, \dots, p_k\}$

Ensure: $S_n(A)$

```

1: function SOLVE( $A, n$ )
2:   initialize empty list solutions
3:   for  $n$ -combination  $S = \{(c_i, r_i) : i = 1, \dots, n\}$  of  $V_d$  do
4:     initialize  $x \in H^{V_d}$  with positive support  $S$  and  $x_{0,0} = -1$ 
5:     if  $x$  is a hyperfield root of every  $p \in A$  then
6:       add  $S$  to solutions
7:     end if
8:   end for
9:   return solutions
10: end function

```

The naive approach has an exponential time complexity since $\binom{(d+1)(d+2)/2}{n}$ many supports are checked.

Efficient Algorithm

For a specific type of system of linear forms A , we can greatly speed up the computation of $S_n(A)$.

Definition 6.16. Let p be a linear form in H^{V_d} , and let $\mathbf{x} \in H^{V_d}$ be some hyperfield root of p . If $\text{supp}(\mathbf{x}) \cap \text{supp}(p) = \emptyset$, then the root \mathbf{x} is called a *trivial root* of p . Otherwise, the root \mathbf{x} is called a *non-trivial root* of p .

Definition 6.17. Let A be a system of linear forms in H^{V_d} . We say $S_n(A)$ is *non-trivial* if $S_n(A) \neq \emptyset$ and every $\mathbf{x} \in S_n(A)$ is a non-trivial root for every form $p \in A$. We say A is *non-trivial* if $S_n(A)$ is non-trivial.

Proposition 6.18. Let A be a system of linear forms in H^{V_d} , $p \in A$ and $\mathbf{x} \in S_n(A)$. Then, the following statements hold:

1. If $(0, 0) \in \text{supp}^+(p)$, then $x_{i,j} = 1$ for some $(i, j) \in \text{supp}^+(p)$.
2. If $(0, 0) \in \text{supp}^-(p)$, then $x_{i,j} = 1$ for some $(i, j) \in \text{supp}^-(p)$.

Proof. Assume $(0, 0) \in \text{supp}^+(p)$. Since $x_{0,0} = -1$, we have $-1 \in \text{sign}(p)(x)$. By assumption, \mathbf{x} is a hyperfield root of p , so $0 \in \text{sign}(p)(\mathbf{x})$. This can only happen if $x_{i,j} = 1$ for some $(i, j) \in \text{supp}^+(p)$. The case $(0, 0) \in \text{supp}^-(p)$ is similar. \square

The next proposition assumes that A is non-trivial.

Proposition 6.19. *Let A be a non-trivial system of linear forms, $p \in A$ and $\mathbf{x} \in S_n(A)$. If $(0, 0) \notin \text{supp}(p)$, then $\text{supp}^+(p) \neq \emptyset$, $\text{supp}^-(p) \neq \emptyset$ as well as $x_{i,j} = x_{i',j'} = 1$ for some $(i, j) \in \text{supp}^+(p)$ and $(i', j') \in \text{supp}^-(p)$.*

Proof. Assume $(0, 0) \notin \text{supp}(p)$. First, $\text{supp}(p) \neq \emptyset$ because $S_n(A)$ is non-empty and consists only of non-trivial roots. If $\text{supp}^+(p) = \emptyset$, then $\text{supp}^+(x) \subset \text{supp}^-(p) = \text{supp}(p) \neq \emptyset$. Hence, $\text{sign}(p)(x) = \{-1\}$, which contradicts \mathbf{x} being a root. Thus, $\text{supp}^+(p)$ is non-empty. Similarly, $\text{supp}^-(p)$ is non-empty.

By non-triviality, $x_{i,j} = 1$ for some $(i, j) \in \text{supp}(p)$. Assume $(i, j) \in \text{supp}^+(p)$. Hence, $1 \in \text{sign}(p)(\mathbf{x})$. Since \mathbf{x} is a root, we also have $0 \in \text{sign}(p)(\mathbf{x})$. This can only occur if $x_{i',j'} = 1$ for some $(i', j') \in \text{supp}^-(p)$. The case $(i, j) \in \text{supp}^-(p)$ is similar. \square

Both propositions allow us to interpret hyperfield linear forms in a non-trivial system A as constraints on the positive supports of roots in $S_n(A)$.

Example 6.20. Fix the degree $d = 3$. Assume a system A and some linear form $p \in A$. Further assume $p = \text{diag}(0)$ is a diagonal Pascal equation of order 0. The support of p is represented by the following diagram:

$$\begin{array}{cccc} + & & & \\ + & . & & \\ + & . & . & \\ + & . & . & . \end{array}$$

We see that any nonzero hyperfield root \mathbf{x} of A satisfies $x_i = 1$ for some

$$(i, j) \in \{(0, 0), (0, 1), (0, 2), (0, 3)\}$$

because x_{00} is negative.

Now, assume A is non-trivial. Consider a row pascal equation $q = \text{row}(3)$ which is contained in A . Its support is depicted by the following diagram:

$$\begin{array}{cccc} - & & & \\ . & + & & \\ . & . & - & \\ . & . & . & + \end{array}$$

For some \mathbf{x} with $\text{supp}^-(\mathbf{x}) = \{(0, 0)\}$ to be a hyperfield root of q , we must have either

1. $x_{i,j} = x_{i',j'} = 1$ for some $(i, j) \in \{(0, 3), (2, 1)\}$ and $(i', j') \in \{(3, 0), (1, 2)\}$, or
2. $\text{supp}(\mathbf{x}) \subset V_3 \setminus \text{supp}(q)$.

Considering only non-trivial roots \mathbf{x} lets us exclude the latter case. Thus, if we have a non-trivial system A with $p, q \in A$, to compute $S_n(A)$, it suffices to check only those hyperfield roots whose support intersected with each of the three following regions is non-empty:

$$\begin{array}{cccc}
 + & & & \\
 + & . & & \\
 + & . & . & \\
 + & . & . & . \\
 \\
 + & & & \\
 . & . & & \\
 . & . & + & \\
 . & . & . & . \\
 \\
 . & & & \\
 . & + & & \\
 . & . & . & \\
 . & . & . & +
 \end{array}$$

Here are examples of such hyperfield roots illustrated in the following diagrams:

$$\begin{array}{cccc}
 + & & & \\
 . & + & & \\
 . & . & . & \\
 . & . & . & . \\
 \\
 . & & & \\
 . & . & & \\
 + & . & + & \\
 . & . & . & +
 \end{array}$$

Definition 6.21. To each linear form p in H^{V_d} we can associate a finite set of supports, which we call $\text{constraints}(p) := \{\text{supp}^+(p) \setminus \{(0, 0)\}, \text{supp}^-(p) \setminus \{(0, 0)\}\} \subset 2^{V_d}$.

The name is justified by the following proposition.

Proposition 6.22. Let p be a linear form in \mathbb{H}^{V_d} , and $\mathbf{x} \in H^{V_d}$ with $\text{supp}^-(\mathbf{x}) = \{(0, 0)\}$. Then, \mathbf{x} is a non-trivial hyperfield root of p if and only if $\text{supp}^+(\mathbf{x}) \cap S \neq \emptyset$ for all $S \in \text{constraints}(p)$.

Proof. Since \mathbf{x} is non-trivial, we clearly have non-empty intersection of $\text{supp}^+(\mathbf{x})$ and $S \in \text{constraints}(p)$. The converse direction is also clear since $p(\mathbf{x}) = 1 - 1 = H$. \square

We present an algorithm for computing $S_n(A)$ of non-trivial systems A .

Algorithm 3 Algorithm for Non-Trivial Systems**Require:** Positive support size n , non-trivial system A **Ensure:** $S_n(A)$

```

1: function SOLVE( $A, n$ )
2:    $C \leftarrow \bigcup_{p \in A} \text{constraints}(p)$ 
3:    $\text{solutions} \leftarrow \{\mathbf{x} \in H^{V_d} \mid \forall S \in C : \text{supp}^+(\mathbf{x}) \cap S \neq \emptyset, |\text{supp}^+(\mathbf{x})| = n, \text{supp}^-(\mathbf{x}) = \{(0, 0)\}\}$ 
4:   return solutions
5: end function

```

Proof of correctness. The correctness of $\text{solutions} = S_n(A)$ follows from Proposition 6.22 and the assumption that A is non-trivial. \square

6.3 Implementation of the Hyperfield Criterion

The Hyperfield Criterion states that only the common hyperfield roots of all Pascal forms can be supports of valid outcomes. The system of all Pascal forms is a-priori an *infinite* and *non-trivial* system. However, we found out that several bases of Pascal forms exist such as the row, col and diag Pascal basis. This allows us to consider *finite* systems. Define the finite system $A = \{\text{diag}(i)\}_{i=0}^d \cup \{\text{row}(i)\}_{i=0}^d \cup \{\text{col}(i)\}_{i=0}^d$.

Proposition 6.23. *The system A is non-trivial.*

Proof. First, $S_n(A)$ is non-empty because $\mathbf{x} = (x_i)_{i=0}^n$ defined as $x_{i,d-i} = \binom{n}{i}$ is a solution of the system A .

Let $\mathbf{x} \in S_n(A)$ and $i = 0, \dots, n$. Consider the following cases.

- Assume, $\mathbf{x} \notin \text{supp}(\text{diag}(i))$; then $\text{diag}(i)(\mathbf{x}) < 0$; we found a contradiction to \mathbf{x} being a root.
- Assume, $\mathbf{x} \notin \text{supp}(\text{row}(i))$. If $i = d$, then \mathbf{x} is not of degree n . Therefore, we assume $i < d$. Then, either \mathbf{x} is a trivial root for $\text{row}(i+1)$ or we have $\text{row}(i+1)(\mathbf{x}) \neq 0$. In the latter case, we found a contradiction to \mathbf{x} being a root. For the former case that \mathbf{x} is a trivial root, we conclude that there exists nonzero $x_{u,d-u}$ for some $u = i+2, \dots, d$ since \mathbf{x} is of degree n ; now we just repeat the argument for $\text{row}(i+1)$. More precisely, if \mathbf{x} is again a trivial root for $\text{row}(i+2)$, we repeat the argument for $\text{row}(i+2)$ until we will end up with a contradiction $\text{row}(u)(\mathbf{x}) \neq 0$.
- For the case $\text{col}(i)$, we can argue by symmetry.

\square

Corollary 6.24. *Configurations $\mathbf{x} \in \mathbb{Z}^{V_d}$ of the form $\text{supp}(\mathbf{x}) \subset \{(i, j) \in \mathbb{Z}^{V_d} : i + j \leq k \text{ or } i > k + 1\}$ are not valid outcomes for any $k = 0, \dots, d - 1$. Neither are configurations $x \in \mathbb{Z}^{V_d}$ of the form $\text{supp}(x) \subset \{(i, j) \in \mathbb{Z}^{V_d} : i + j \leq k \text{ or } j > k + 1\}$ for $k = 0, \dots, d - 1$ due to symmetry.*

Proof. Since the previously defined system A is non-trivial, we must have that supports of valid outcomes intersect the support of $\text{row}(k + 1)$ non-trivially. \square

Example 6.25. This is not a valid outcome:

```

.
. .
. . .
. . . *
* . . * *
* * . * * *
```

Now that we have shown that A is a trivial system, we have found an efficient way to apply the Hyperfield Criterion. Here is a detailed breakdown of an implementation of the algorithm.

6.4 Contractions

We remind that we want to show that for all valid integral outcomes \mathbf{w} with $|\text{supp}^+(\mathbf{w})| = 4$ its degree is at most five. The problem is that the number of valid outcomes to check is infinite since the degree could be arbitrarily large. To overcome this issue, we will introduce *contractions* that allows us to check finitely many cases while still covering all valid outcomes of arbitrarily large degree. We follow the approach of Bik and Marigliano in [2].

The idea of contraction is to *contract* or *consolidate* vertices in V_d by merging some rows and columns into a single vertex. We do this by defining new formal variables $b_i, c_i, d_i, e_i, y_{i,j}$ and $z_{i,j}$ called *contraction variables*.




Figure 6.1: This figure illustrates the contraction variables and is taken from [2]. The yellow areas X, Y, Z represent formal variables $x_{i,j}$ that are unaffected by the contraction. The mint areas B, C, D represent rows and columns of vertices that are merged into a single vertex. The area D is further divided into areas D_1, D_2 by alternating columns.

Definition 6.26. Let $x_{i,j}$ be formal variables indexed by V_d . We merge a subset of rows and columns of formal variables $x_{i,j}$ in V_d into a single vertex by defining the following *contraction*

$$\begin{aligned} y_{i,j} &:= x_{i,d-3-i+j} && \text{for } i, j = 0, \dots, 3, \\ z_{i,j} &:= x_{d-3-j+i,j} && \text{for } i, j = 0, \dots, 3, \\ b_j &:= x_{4,j} + \dots + x_{d-4-j,j} && \text{for } j = 0, \dots, 3, \\ c_i &:= x_{i,4} + \dots + x_{i,d-4-i} && \text{for } i = 0, \dots, 3, \\ d_k &:= \begin{cases} x_{4,d-4-k} + x_{6,d-6-k} + \dots + x_{d-4-k,4} & \text{if } d+k \text{ is even} \\ x_{4,d-4-k} + x_{6,d-6-k} + \dots + x_{d-5-k,5} & \text{if } d+k \text{ is odd} \end{cases} && \text{for } k = 0, \dots, 3, \\ e_k &:= \begin{cases} x_{5,d-5-k} + x_{7,d-7-k} + \dots + x_{d-5-k,5} & \text{if } d+k \text{ is even} \\ x_{5,d-5-k} + x_{7,d-7-k} + \dots + x_{d-4-k,4} & \text{if } d+k \text{ is odd} \end{cases} && \text{for } k = 0, \dots, 3. \end{aligned}$$

$y_{0,3}$																
$y_{0,2}$	$y_{1,3}$															
$y_{0,1}$	$y_{1,2}$	$y_{2,3}$														
$y_{0,0}$	$y_{1,1}$	$y_{2,2}$	$y_{3,3}$													
c_0	$y_{1,0}$	$y_{2,1}$	$y_{3,2}$	d_0												
c_0	c_1	$y_{2,0}$	$y_{3,1}$	d_1	e_0											
c_0	c_1	c_2	$y_{3,0}$	d_2	e_1	d_0										
c_0	c_1	c_2	c_3	d_3	e_2	d_1	e_0									
c_0	c_1	c_2	c_3	$*$	e_3	d_2	e_1	d_0								
c_0	c_1	c_2	c_3	$*$	$*$	d_3	e_2	d_1	e_0							
c_0	c_1	c_2	c_3	$*$	$*$	$*$	e_3	d_2	e_1	d_0						
c_0	c_1	c_2	c_3	$*$	$*$	$*$	$*$	d_3	e_2	d_1	e_0					
c_0	c_1	c_2	c_3	$*$	$*$	$*$	$*$	$*$	e_3	d_2	e_1	d_0				
$x_{0,3}$	$x_{1,3}$	$x_{2,3}$	$x_{3,3}$	b_3	b_3	b_3	b_3	b_3	b_3	$z_{0,3}$	$z_{1,3}$	$z_{2,3}$	$z_{3,3}$			
$x_{0,2}$	$x_{1,2}$	$x_{2,2}$	$x_{3,2}$	b_2	b_2	b_2	b_2	b_2	b_2	b_2	$z_{0,2}$	$z_{1,2}$	$z_{2,2}$	$z_{3,2}$		
$x_{0,1}$	$x_{1,1}$	$x_{2,1}$	$x_{3,1}$	b_1	b_1	b_1	b_1	b_1	b_1	b_1	b_1	$z_{0,1}$	$z_{1,1}$	$z_{2,1}$	$z_{3,1}$	
$x_{0,0}$	$x_{1,0}$	$x_{2,0}$	$x_{3,0}$	b_0	b_0	b_0	b_0	b_0	b_0	b_0	b_0	b_0	$z_{0,0}$	$z_{1,0}$	$z_{2,0}$	$z_{3,0}$

As we can see, the vertices $x_{0,4}, \dots, x_{0,d-4}$ are merged into the contraction variable c_0 by summing them up. Similarly, the formal variables $x_{i,4}, \dots, x_{i,d-4-i}$ are merged into c_i for $i = 1, 2, 3$.

We remind that hyperfield Pascal forms are expressed as sums $\sum_{(i,j) \in V_d} \lambda_{i,j} x_{i,j}$ with $\lambda_{i,j} \in H$. The key insight is that there are some Pascal forms that can be expressed in

terms of the contraction variables c_0, c_1, c_2 and c_3 instead of the original variables $x_{i,j}$ for all vertices (i, j) in the C -area of Figure 6.4.

Example 6.27. Consider the Pascal form $\text{diag}(1)$ in $\mathbb{Z}^{V_{16}}$. Its support is depicted in the following figure:



We see that $\text{diag}(1) = x_{0,0} + x_{0,1} + x_{0,2} + x_{0,3} + x_{1,0} + x_{1,1} + x_{1,2} + x_{1,3} + y_{0,0} + y_{0,1} + y_{0,2} + y_{1,0} + y_{1,1} + y_{1,2} + y_{1,3} + c_0 + c_1$.

The previous example also demonstrates that the expression $\text{diag}(1) = x_{0,0} + x_{0,1} + x_{0,2} + x_{0,3} + x_{1,0} + x_{1,1} + x_{1,2} + x_{1,3} + y_{0,0} + y_{0,1} + y_{0,2} + y_{1,0} + y_{1,1} + y_{1,2} + y_{1,3} + c_0 + c_1$ is *independent* of the degree d , i.e. if we were to consider the Pascal form $\text{diag}(1)$ in \mathbb{Z}^{V_d} for some arbitrary d , the expression would still hold. This is great news since it allows us to express Pascal forms in terms of contraction variables for all degrees d at once.

So far, we have only considered the contraction variables c_0, c_1, c_2 , and c_3 . As we might expect, we can also express some Pascal forms in terms of the contraction variables $b_0, b_1, b_2, b_3, d_0, d_1, d_2, d_3, e_0, e_1, e_2, e_3, y_{i,j}$, and $z_{i,j}$. We will now find these kinds of Pascal forms that can be represented in terms of the contraction variables independent of the degree d . A good set of Pascal forms to consider are the Pascal forms $\text{diag}(k), \text{col}(k)$ and $\text{row}(k)$ for $k = 0, 1, 2, 3, d-3, d-2, d-1, d$.

Proposition 6.28. *Let $d \geq 11$. Let p be a hyperfield form induced by one of the following Pascal forms on \mathbb{Z}^{V_d} :*

1. $\text{col}(1), \text{col}(2), \text{col}(3)$, or

2. $\text{row}(1), \text{row}(2), \text{row}(3)$, or
3. $\text{diag}(1), \text{diag}(2), \text{diag}(3)$, or
4. $\text{diag}(d-1), \text{diag}(d-2), \text{diag}(d-3)$.

Then, we have

$$p = \sum_{i,j=0}^3 \lambda_{i,j}^{(x)} x_{i,j} + \sum_{i,j=0}^3 \lambda_{i,j}^{(y)} y_{i,j} + \sum_{i,j=0}^3 \lambda_{i,j}^{(z)} z_{i,j} + \sum_{j=0}^3 \lambda_j^{(b)} b_j + \sum_{i=0}^3 \lambda_i^{(c)} c_i + \sum_{k=0}^3 \lambda_k^{(d)} d_k + \sum_{k=0}^3 \lambda_k^{(e)} e_k$$

with coefficients $\lambda_{i,j}^{(x)}, \lambda_{i,j}^{(y)}, \lambda_{i,j}^{(z)}, \lambda_j^{(b)}, \lambda_i^{(c)}, \lambda_k^{(d)}, \lambda_k^{(e)} \in H$.

Define $\lambda(p, d) := \left(\lambda_{i,j}^{(x)}, \lambda_{i,j}^{(y)}, \lambda_{i,j}^{(z)}, \lambda_j^{(b)}, \lambda_i^{(c)}, \lambda_k^{(d)}, \lambda_k^{(e)} \right)$ for the coefficients of p on H^{V_d} . Then, we have

$$\lambda(p, d) = \lambda(p, d+1) = \lambda(p, d+2) = \lambda(p, d+3) = \dots$$

In other words, all the coefficients $\lambda_{i,j}^{(x)}, \lambda_{i,j}^{(y)}, \lambda_{i,j}^{(z)}, \lambda_j^{(b)}, \lambda_i^{(c)}, \lambda_k^{(d)}, \lambda_k^{(e)}$ are independent of the degree d .

Proof. For case two and three we observe that the hyperfield form p has support contained in the areas X, C , and Y from Figure 6.4. This follows directly from Proposition 6.11. We also see that p depends only on the column sums on C .

For case one and four we see that p has support contained in the areas X, B , and Z from Figure 6.4 by Proposition 6.11. We conclude that p depends only on the row sums on B . \square

Here is another example.

Example 6.29. Consider $\text{diag}(3)$ and $d = 11$.

```

.
. .
. . .
+ + + +
+ + + + .
+ + + + . .
+ + + + . . .
+ + + + . . . .
+ + + + . . . . .
+ + + + . . . . .
+ + + + . . . . .
+ + + + . . . . .
+ + + + . . . . .

```


$$\begin{array}{cccccccc}
. & . & . & . & + & + & + & + \\
. & . & . & . & . & - & - & - - \\
. & . & . & . & . & . & + & + + + \\
. & . & . & . & . & . & . & - - - - \\
. & . & . & . & . & . & . & + + + + \\
. & . & . & . & . & . & . & - - - -
\end{array}$$

We write

$$\text{sign}(\text{col}(d-3)) = \sum_{i=0}^3 \sum_{j=0}^i (-1)^{i+j} y_{i,j} - \sum_{k=0}^3 (-1)^k d_k + \sum_{k=0}^3 (-1)^k e_k - \sum_{i,j=0}^3 (-1)^j z_{i,j}.$$

For $d = 13$, the support of $\text{sign}(\text{col}(d-3))$ is depicted in the following figure:

$$\begin{array}{cccccccc}
. & & & & & & & \\
. & . & & & & & & \\
. & . & . & & & & & \\
+ & + & + & + & & & & \\
. & - & - & - & - & & & \\
. & . & + & + & + & + & & \\
. & . & . & - & - & - & - & \\
. & . & . & . & + & + & + & + \\
. & . & . & . & . & - & - & - - \\
. & . & . & . & . & . & + & + + + \\
. & . & . & . & . & . & . & - - - - \\
. & . & . & . & . & . & . & + + + + \\
. & . & . & . & . & . & . & - - - - \\
. & . & . & . & . & . & . & + + + +
\end{array}$$

We write

$$\text{sign}(\text{col}(d-3)) = \sum_{i=0}^3 \sum_{j=0}^i (-1)^{i+j} y_{i,j} - \sum_{k=0}^3 (-1)^k d_k + \sum_{k=0}^3 (-1)^k e_k + \sum_{i,j=0}^3 (-1)^j z_{i,j}.$$

We have merged formal variables $x_{i,j}$ indexed by vertices (i, j) in the areas B, C, D into contraction variables. Now, we apply these contractions to concrete elements $\mathbf{s} \in H^{V_d}$.

Definition 6.32. Define the index set

$$\Xi := \{0, 1, 2, 3\}^2 \sqcup \{0, 1, 2, 3\}^2 \sqcup \{0, 1, 2, 3\}^2 \sqcup \{0, 1, 2, 3\} \sqcup \{0, 1, 2, 3\} \sqcup \{0, 1, 2, 3\} \sqcup \{0, 1, 2, 3\}.$$

Let $\mathbf{s} \in H^\Xi$. We call \mathbf{s} a *contracted hyperfield configuration*, and write \mathbf{s} as

$$\mathbf{s} = (\mathbf{x}, \mathbf{y}, \mathbf{z}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}) = (x_{i,j}, y_{i,j}, z_{i,j}, b_j, c_i, d_k, e_k).$$

Definition 6.33. Let $\mathbf{s} \in H^\Xi$ be a contracted hyperfield configuration. We say \mathbf{s} is *valid* if one of the following holds:

1. $\mathbf{s} = \mathbf{0}$ or
2. $s_{0,0} = -1$, $x_{i,j} \geq 0$ for all $i+j > 0$, and $y_{i,j}, z_{i,j}, b_j, c_i, d_k, e_k \geq 0$ for all $i, j, k = 0, 1, 2, 3$.

Going from the world of hyperfield configurations to the world of *contracted* hyperfield configurations is done via the following map.

Definition 6.34. Let $d \geq 11$. Let $\mathbf{s} \in H^{V_d}$ be a hyperfield configuration. We define

$$\text{contr}_d(\mathbf{s}) : \mathbf{s} \mapsto (\mathbf{x}, \mathbf{y}, \mathbf{z}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}) = (x_{i,j}, y_{i,j}, z_{i,j}, b_j, c_i, d_k, e_k)$$

where we set

$$\begin{aligned} x_{i,j} &:= s_{i,j} && \text{for } i, j = 0, \dots, 3, \\ y_{i,j} &:= s_{i,d-3-i+j} && \text{for } i, j = 0, \dots, 3, \\ z_{i,j} &:= s_{d-3-j+i,j} && \text{for } i, j = 0, \dots, 3, \\ b_j &:= s_{4,j} + \dots + s_{d-4-j,j} && \text{for } j = 0, \dots, 3, \\ c_i &:= s_{i,4} + \dots + s_{i,d-4-i} && \text{for } i = 0, \dots, 3, \\ d_k &:= \begin{cases} s_{4,d-4-k} + s_{6,d-6-k} + \dots + s_{d-4-k,4} & \text{if } d+k \text{ is even} \\ s_{4,d-4-k} + s_{6,d-6-k} + \dots + s_{d-5-k,5} & \text{if } d+k \text{ is odd} \end{cases} && \text{for } k = 0, \dots, 3, \\ e_k &:= \begin{cases} s_{5,d-5-k} + s_{7,d-7-k} + \dots + s_{d-5-k,5} & \text{if } d+k \text{ is even} \\ s_{5,d-5-k} + s_{7,d-7-k} + \dots + s_{d-4-k,4} & \text{if } d+k \text{ is odd} \end{cases} && \text{for } k = 0, \dots, 3. \end{aligned}$$

exactly as in Definition 6.26.

The contraction map contr_d maps hyperfield configurations $\mathbf{s} = (\mathbf{x}, \mathbf{y}, \mathbf{z}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}) \in H^{V_d}$ to elements in H^Ξ if $\mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e} \geq 0$. If one of the entries is negative, the map may output to some element $(2^H)^\Xi$. To make life easier, we introduce the following definition.

Definition 6.35. Let $\mathbf{s} \in H^{V_d}$ be a hyperfield configuration. We say \mathbf{s} is *weakly valid* if for all $(i, j) \in \text{supp}^-(\mathbf{s})$ one of the following holds:

1. $i, j = 0, \dots, 3$, or
2. $i = 0, \dots, 3$ and $i + j \geq d - 3$, or
3. $j = 0, \dots, 3$ and $i + j \geq d - 3$.




Figure 6.3: A hyperfield configuration is weakly valid if its negative support is contained in the yellow area. The figure is taken from [2].

From now on, we only consider *weakly valid* hyperfield configurations because in this case the contraction map contr_d always outputs elements in H^Ξ .

Definition 6.36. Let $\mathbf{s} \in H^\Xi$ be a contracted hyperfield configuration. The *positive support* of $\mathbf{s} = (\mathbf{x}, \mathbf{y}, \mathbf{z}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e})$ is defined as the set of all symbols $x_{i,j}, y_{i,j}, z_{i,j}, b_j, c_i, d_k, e_k$ such that the corresponding coefficients of \mathbf{s} equal to one.

Example 6.37. Let $\mathbf{s} = (\mathbf{x}, \mathbf{y}, \mathbf{z}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}) \in H^\Xi$ be a contracted hyperfield configuration defined by

$$x_{0,0} = -1, \quad x_{0,3} = 1, \quad x_{1,1} = 1, \quad x_{3,0} = 1, \quad d_0 = 1, \quad e_0 = 1$$

where all other entries are zero. Then, the positive support of \mathbf{s} is given by

$$\text{supp}^+(\mathbf{s}) = \{x_{0,3}, x_{1,1}, x_{3,0}, d_0, e_0\}.$$

Chapter 7

Valid Outcomes of Positive Support Size Four

We have all the tools to prove the main result of this chapter, namely for all valid integral outcomes \mathbf{w} with $|\text{supp}^+(\mathbf{w})| = 4$ we have

$$\deg(\mathbf{w}) \leq 5.$$

As in previous chapters, we characterize outcomes as roots of Pascal forms. So, we define the following two systems of Pascal forms that valid outcomes must be roots of:

$$\begin{aligned} \Phi_1 := \{ & \text{col}(1), \text{col}(2), \text{col}(3), \text{row}(1), \text{row}(2), \text{row}(3), \\ & \text{diag}(1), \text{diag}(2), \text{diag}(3), \text{diag}(d-1), \text{diag}(d-2), \text{diag}(d-3) \}, \end{aligned}$$

and

$$\begin{aligned} \Phi_2 := \{ & \text{col}(d), \text{col}(d-1), \text{col}(d-2), \text{col}(d-3), \\ & \text{row}(d), \text{row}(d-1), \text{row}(d-2), \text{row}(d-3) \}. \end{aligned}$$

We also define $\Phi := \Phi_1 \cup \Phi_2$.

By Proposition 6.28, we can write all hyperfield forms induced by Pascal forms p in Φ_1 as

$$\text{sign}(p) = \hat{p}$$

for some linear form $\hat{p} \in H[\mathbf{x}, \mathbf{y}, \mathbf{z}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}]$ if $d \geq 11$. This linear form is independent of the degree d . To make notations consistent later, we set $\hat{p}^{\text{even}} := \hat{p}^{\text{odd}} := \hat{p}$.

Similarly, by Proposition 6.30, we can write all hyperfield forms induced by Pascal forms p in Φ_2 as

$$\text{sign}(p) = \begin{cases} \hat{p}^{\text{even}} & \text{if } d \text{ is even} \\ \hat{p}^{\text{odd}} & \text{if } d \text{ is odd} \end{cases}$$

for some linear forms $\hat{p}^{\text{even}}, \hat{p}^{\text{odd}} \in H[\mathbf{x}, \mathbf{y}, \mathbf{z}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}]$ if $d \geq 12$. These linear forms $\hat{p}^{\text{even}}, \hat{p}^{\text{odd}}$ are independent of the degree d .

Definition 7.1. We define the following three solution sets:

1. Define Γ_d to be the set of all valid hyperfield configurations $\mathbf{s} \in H^{V_d}$ of degree d such that $\text{sign}(p)(\mathbf{s}) = H$ for all $p \in \Phi$.
2. Define Γ^{even} to be the set of all valid contracted hyperfield configurations $\mathbf{s} \in H^{\Xi}$ such that $\hat{p}^{\text{even}}(\mathbf{s}) = H$ for all $p \in \Phi$.
3. Define Γ^{odd} to be the set of all valid contracted hyperfield configurations $\mathbf{s} \in H^{\Xi}$ such that $\hat{p}^{\text{odd}}(\mathbf{s}) = H$ for all $p \in \Phi$.

By Proposition 6.8, valid chipsplitting outcomes of degree d have supports in Γ_d . This is the reason why we have defined Γ_d in the first place.

Proposition 7.2. *Let $d \geq 12$. Then, the following holds:*

1. *If d is even, then $\Gamma_d = \text{contr}_d^{-1}(\Gamma^{\text{even}})$.*
2. *If d is odd, then $\Gamma_d = \text{contr}_d^{-1}(\Gamma^{\text{odd}})$.*

Proof. Let $d \geq 12$ be even. Let $\mathbf{s} \in H^{V_d}$ be a hyperfield configuration and $p \in \Phi$. Then, we have

$$\text{sign}(p)(\mathbf{s}) = \hat{p}^{\text{even}}(\text{contr}_d(\mathbf{s})).$$

by definition of \hat{p}^{even} . If $\mathbf{s} \in \Gamma_d$, then $H = \text{sign}(p)(\mathbf{s}) = \hat{p}^{\text{even}}(\text{contr}_d(\mathbf{s}))$. Hence, $\text{contr}_d(\mathbf{s})$ is contained in Γ^{even} . If $\text{contr}_d(\mathbf{s}) \in \Gamma^{\text{even}}$ holds, using the equation above we also see that $\mathbf{s} \in \Gamma_d$. This shows that $\Gamma_d = \text{contr}_d^{-1}(\Gamma^{\text{even}})$.

The second statement for odd degrees d follows analogously. \square

Corollary 7.3. *Let $d \geq 12$. Let $\mathbf{w} \in \mathbb{Z}^{V_d}$ be a valid outcome. Then, we have*

$$\text{contr}_d(\text{sign}(\mathbf{w})) \in \Gamma^{\text{even}} \cup \Gamma^{\text{odd}}.$$

Proof. Define $\mathbf{s} := \text{sign}(\mathbf{w})$. By Proposition 6.12 we have $\mathbf{s} \in \Gamma_d$. If d is even, then $\text{contr}_d(\mathbf{s}) \in \Gamma^{\text{even}}$ by the previous proposition. If d is odd, then $\text{contr}_d(\mathbf{s}) \in \Gamma^{\text{odd}}$ by the previous proposition. This shows the claim. \square

This corollary allows us to exclude certain outcomes as valid outcomes. Assume we have some contracted hyperfield configuration $\xi \in H^{\Xi}$ that is not a root of some of the linear forms $\hat{p}^{\text{even}}, \hat{p}^{\text{odd}}$ for $p \in \Phi$. Then, any chipsplitting configuration $\mathbf{w} \in \mathbb{Z}^{V_d}$ with $\text{contr}_d(\text{sign}(\mathbf{w})) = \xi$ is not a valid outcome.

Proposition 7.4. *Let $\mathbf{s} \in H^{V_d}$ be a valid hyperfield configuration of degree d and positive support size four. Define the set Ω_d to be*

$$\left\{ \mathbf{s} \in H^{V_d} \mid \deg(\mathbf{s}) = d, |\text{supp}^+(\mathbf{s})| = 4, \text{sign}(p)(\mathbf{s}) = H \forall p \in \{\text{row}(k), \text{col}(k), \text{diag}(k)\}_{k=0}^d \right\}.$$

Then, all of the following hold:

1. Let $d = 6$. Then, \mathbf{s} is contained in Ω_d if and only if its positive support is exactly one of the following:

$$\{(0, 3), (1, 5), (4, 1), (6, 0)\}, \{(0, 5), (1, 1), (3, 3), (6, 0)\}, \{(0, 6), (1, 1), (3, 3), (5, 0)\}, \\ \{(0, 6), (1, 1), (3, 3), (6, 0)\}, \{(0, 6), (1, 4), (3, 0), (5, 1)\}.$$

2. Let $d = 7$. Then, \mathbf{s} is contained in Ω_d if and only if its positive support is exactly one of the following:

$$\{(0, 7), (1, 1), (3, 3), (7, 0)\}, \{(0, 7), (1, 3), (5, 1), (7, 0)\}, \{(0, 7), (1, 5), (3, 1), (7, 0)\}.$$

3. Let $d \geq 8$. Then $\mathbf{s} \notin \Omega_d$.

Proof. \mathbf{s}

□

Bibliography

- [1] Matt Baker and Nathan Bowler. “Matroids over hyperfields”. In: *AIP Conference Proceedings*. Vol. 1978. 1. AIP Publishing, 2018.
- [2] Arthur Bik and Orlando Marigliano. “Classifying one-dimensional discrete models with maximum likelihood degree one”. In: *arXiv preprint arXiv:2205.09547* (2022).
- [3] Eliana Duarte, Orlando Marigliano, and Bernd Sturmfels. “Discrete statistical models with rational maximum likelihood estimator”. In: (2021).
- [4] June Huh. “Varieties with maximum likelihood degree one”. In: *arXiv preprint arXiv:1301.2732* (2013).
- [5] Caroline J Klivans. *The mathematics of chip-firing*. Chapman and Hall/CRC, 2018.