

Investigating the Crystal Cargo of Fiamme from the Ora Ignimbrite (Permian, Italy): Elucidating the Spatio-Temporal Evolution of a Crystal-Rich, Supereruption-Sized Magma System

This manuscript ([permalink](#)) was automatically generated from [gennachiaro/phd_proposal@f670c82](#) on February 18, 2020.

Authors

- **Genna Chiaro**

 [0000-0002-8981-7679](#) ·  [gennachiaro](#) ·  [gennachiaro](#)

Department of Earth and Environmental Sciences, Vanderbilt University

Motivation

Supereruptions, events that expel >450 km³ of magma from the subsurface, are the most extreme expressions of eruptive volcanism on Earth (Sparks et al., 2005). These catastrophic events tend to occur every ~100,000 years (Mason et al., 2004). Due to their scarcity, we are limited to studying the resulting deposits of these colossal eruptions. By investigating the evolutionary history preserved in mineral rims, unaltered volcanic glass compositions, spatial distribution of the deposits, and timescales of magmatic processes, we can inform our current hypotheses regarding the formation of large silicic magma systems that are capable of supereruption.

Our inability to obtain high-resolution subsurface images has spurred innovative methods for resolving the organization of magma in the crust. If we know the major element concentrations of the original melt and the minerals present, we can obtain temperature and pressure estimates using the thermodynamics software, Rhyolite-MELTS (Gualda and Ghiorso, 2014). Importantly, this can help determine where the magma bodies were situated in the crust prior to eruption. Major and trace element compositions of volcanic glasses can provide information on the number of magma bodies that contributed to the supereruption (Begué et al., 2014). Furthermore, both the lateral distribution of volcanic glasses and ⁴⁰Ar/³⁹Ar sanidine geochronology can provide insight on the evolution of the eruption and whether the deposit was emplaced instantaneously or had distinct volcanic pulses (Begué et al., 2014; Kay et al., 2011).

I plan to investigate the pre-eruptive conditions of a crystal-rich supereruption, the Ora Ignimbrite, in order to fill current gaps in knowledge regarding the structure and evolution of highly-evolved, crystal-rich, trans-crustal magma systems.

The following research questions are the main focus of my dissertation:

- 1. Can we use mineral major and trace element compositions to identify fiamme type and population?**
- 2. Do we observe similar histories recorded in the mineral populations? What can this tell us about magmatic processes during the assembly and eruption of large silicic systems?**
- 3. Can we observe clustering of fiamme populations in space and time? What does this suggest about the subsurface architecture of the Ora magmatic system? Are we able to identify different phases of the eruption and the tapping of separate vents?**

4. Can high-precision $^{40}\text{Ar}/^{39}\text{Ar}$ sanidine ages subdivide the Ora eruption into distinguishable volcanic pulses, thereby illuminating the episodic tempo and dynamic behavior of this very large magmatic system?

The Permian Ora Ignimbrites ($>1,200 \text{ km}^3$) are advantageous deposits for this study because glacial incision has revealed exceptional intracaldera exposures with $>1,000 \text{ m}$ of vertical relief. I plan to expand upon my M.S. work on the Ora vitrophyre horizons and develop major and trace element-based hierarchical clustering algorithms in python to identify minerals that correspond to the pre-defined fiamme types and populations identified in Chiaro (2019). A hot-spot clustering analysis (Ripley's k-function) in ArcGIS will be used to determine the distribution of fiamme types and populations throughout the entire deposit and high-precision $^{40}\text{Ar}/^{39}\text{Ar}$ sanidine ages will provide eruption ages. By integrating this project with the Rhyolite-MELTS pressures calculated in Chiaro et al., (in prep), we hope to elucidate the spatio-temporal evolution of a supereruption-sized, crystal-rich magma system.

Background

The supereruption-scale Ora Ignimbrites are products of an ignimbrite flareup resulting from subduction of the Paleotethys Ocean and the consequent formation of Pangaea (Cassinis et al., 2012). Glacial incision of the Permian Ora Ignimbrites ($>1,200 \text{ km}^3$) reveals heterogeneous intracaldera deposits with $>1,000 \text{ m}$ of relief. Less voluminous outflow was emplaced to the north and south of the caldera and is stratigraphically similar to late-erupted intracaldera units (Marocchi et al., 2008). The Ora deposits are interpreted to be low-intensity eruptions that tapped multiple magma bodies and were sourced from discrete vents (Willcock et al., 2015). Bulk rock chemistry and mineralogy suggest that caldera collapse and eruption initiated in the south and migrated northward through time (Willcock et al., 2013).



Figure 1: A map of the field location. The four fiamme types identified in Chiaro et al. (in prep) are listed with their corresponding vitrophyre locations.

Table 1: Samples collected in Fall 2017 and Spring 2019. Sample locations are shown in Fig. 1. The number of bulk tuff samples and fiamme are listed.

| Samples: | Location: | Type: | Bulk Tuff: | Fiamme: |
|--------------|-------------------------|---------------|------------|---------|
| ORA 2 | Outflow Vitrophyre | Vitrophyre | 9 | 31 |
| ORA 3 | Northern Intracaldera | Bulk Tuff | 1 | 1 |
| ORA 4 | Odorizzi Quarry | Bulk Tuff | 2 | 50 |
| ORA 5 | Intracaldera Vitrophyre | Vitrophyre | 1 | 30 |
| ORA 6 | Northern Intracaldera | Rhyolite Dike | 4 | |

| Samples: | Location: | Type: | Bulk Tuff: | Fiamme: |
|---------------|-------------------------------|---------------------------------|------------|---------|
| ORA 7 | Base of Northern Intracaldera | Bulk Tuff | 1 | |
| ORA 8 | Northern Intracaldera | Bulk Tuff | 1 | |
| ORA 9 | NW Outflow 1 | Biotite-Rich Fiamme | | 1 |
| ORA 10 | NW Outflow 2 | Medium-Grained Fiamme | | 1 |
| ORA 11 | Castel Firmiano | Ash/Ignimbrite Contact | 3 | |
| ORA 12 | Rhyolite Dike | Granite Dike | 4 | |
| ORA 13 | Rhyolite Dike | Potential Fiamma | | 1 |
| ORA 14 | Caldera Rim Cutting Dike | Porphyritic Coarse-Grained Dike | 1 | |
| ORA 15 | Caldera Rim Cutting Dike | Medium-Grained Dike | 1 | |
| ORA 16 | Cava Flor Quarry | Fine and Coarse-Grained Fiamme | | 4 |
| ORA 17 | NW Outflow 3 | Fine to Medium-Grained Fiamme | | 3 |
| ORA 18 | Magdalena Hike | Pseudotachylite | 1 | |
| ORA 19 | Magdalena Hike @ Sculpture | Fine and Coarse-Grained Dikes | 4 | |

Recent work investigated the pre-eruptive architecture of the Ora system with textural, mineralogical, and geochemical features of fiamme (Chiaro, 2019). Fiamme were collected from two vitrophyre horizons, an early-erupted intracaldera deposit and a late-erupted outflow deposit, in order to find well-preserved glass. These fiamme were initially grouped into four types based on crystal content and biotite content (Table 2). Glass major elements further verified the categorization scheme (Fig. 2). Tight clustering of MG and VCCR glass major element compositions suggests that the glasses are unaltered and thus represent pristine melt compositions (Fig. 2).

Table 2: The four fiamme types and their location, crystal content, maximum phenocryst size, and mafic content (Chiaro et al., in prep).

| FIAMMA TYPES: | DESCRIPTION: | LOCATION | CRYSTAL CONTENT: | MAX PHENOCRYST SIZE: | MAFICS: |
|----------------|--|--------------|------------------|----------------------|---------|
| TYPE 1: | Very Coarse-Grained Crystal-Rich (VCCR) | Intracaldera | ~40-50% | >5 mm | ~2-3% |
| TYPE 2: | Medium-Grained (MG) | Outflow | ~20-40% | 1-3 mm | <0.5% |
| TYPE 3: | Fine-Grained (FG) | Intracaldera | ~20% | ≤1 mm | <0.1% |
| TYPE 4: | Fine-Grained Crystal-Poor (FGCP) | Outflow | ~10% | ≤1 mm | ≤0.15% |




Figure 2: All fiamme major element glass compositions. The colors correspond to the fiamme types shown above in Table 2. (a) The tight clustering of VCCR and MG fiamme suggest that these glasses did not undergo alteration. (b) Clustering in the Na2O vs. K2O plot provides further evidence that the VCCR and MG glasses represent pristine melt compositions.

Glass trace elements illuminate at least three distinct populations for both the VCCR and MG fiamme, suggesting the presence of multiple magma batches within the Ora system (Fig. 3). Pre-eruptive storage pressures from the Rhyolite-MELTS (Q2F) geobarometer suggest that there were two potential scenarios for the subsurface organization of magma bodies in the Ora system (Fig. 4). Either there were two distinct crystal mush zones located at slightly different depths within the crust, or there was a continuous crystal mush present with multiple heterogeneous zones located throughout (Chiaro, 2019).




Figure 3: (a) A rare earth element diagram showing differences between the early-erupted intracaldera (VCCR) and late-erupted outflow (MG) fiamme. (b) A plot of strontium vs. barium illustrates six discrete glass populations from the VCCR and MG fiamme. Each population is inferred to represent a separate magma batch.




Figure 4: Different fiamme types were stored at different pressures in the crust, with a progression of erupting shallower to deeper magmas through time. Silica content vs. pressure shows variable storage depths for the FGCP melt, suggesting that there were multiple, small melt bodies located at different depths throughout the magma chamber (from Chiaro et al., in prep).

Approach

Sampling horizons for this project were chosen to (1) span the full duration of ignimbrite emplacement, (2) represent key lithologies based on observed textural variations and bulk rock geochemistry, and (3) reflect features within the section that might relate to possible time gaps (Fig. 5). These locations were identified from stratigraphic logs in Willcock et al. (2013) which characterize observed changes in volcanic lithofacies. I also utilized work detailing the stratigraphic variation of bulk rock compositions and crystal contents from Willcock et al. (2015) to inform my sampling strategy.



Figure 5: (a) A map of the Ora Ignimbrite deposits with acquired samples and future sampling sites. Modified from Willcock et al., 2013. (b) A cross section of the Athesian Volcanic Group from NNW to SSE. Future sampling sites are shown with their approximate stratigraphic horizons in the intracaldera and outflows. Modified from Bradner et al., 2016.

Mineral-Based Fiamme Classification:

I plan to build on the work completed for my M.S. (Chiaro, 2019) and determine a mineral-based characterization scheme in order to categorize fiamme throughout the entire Ora deposit. Since fiamme with unaltered glass are only present at the vitrophyre horizons, we will use mineral compositions to fingerprint each fiamma. This involves collecting major and trace elements of plagioclase, alkali feldspar, and biotite for each fiamme type and population. Preliminary work shows that biotite and plagioclase major elements can be used to differentiate intracaldera vs. outflow fiamme (Fig. 6). Furthermore, biotite major element compositions can be used to infer both fiamme type and population (Fig. 6).

Sliwinski et al. (2017) has shown that biotite trace element compositions can be used to fingerprint magma batches at Las Cañadas volcano in Tenerife. Our glass trace element compositions differentiate fiamme populations in Ora; each fiamme population is interpreted to represent an individual magma batch (Fig. 3, 6). Biotite grown from these discrete magma batches will likely retain different rim trace element compositions, allowing for mineral-based classification for different magma batches.




Figure 6: Biotite and plagioclase major element compositions from fiamme. Fiamme are divided into population based on hue. Both plagioclase and biotite can be used to determine fiamme type and biotite can also determine fiamme population (from Chiaro, 2019).

Hierarchical Clustering and Machine Learning in Python:

The foundation for this project requires major and trace element compositions for entire minerals. The major element compositions for feldspars and biotite will be collected from EDS maps. Afterwards, we will obtain trace element profiles for each of the grains.

I will develop a hierarchical agglomerative clustering algorithm in python for major elements in order to statistically determine how many clusters of minerals are present in the sampled fiamme. Another clustering algorithm will be completed using trace elements and we will compare the results. Once the clusters are identified, we can observe whether they correlate with fiamme type and population (Chiaro, 2019). If we see a correlation, we can confidently use mineral compositions to classify fiamme throughout the Ora Ignimbrite.

We can also use the mineral maps to visualize core to rim cluster evolution (Fig. 7). By noting where the clusters occur in each mineral and by looking at a large enough population of core to rim clusters, we can infer whether crystals experienced similar histories. This approach may reveal information regarding the magmatic processes that occurred prior to the Ora eruption (injection, decompression-driven growth, etc).




Figure 7: (A-C) BSE and EDS scans of a clinopyroxene phenocryst from the Holuhraun-Bardarbunga eruption in 2014-2015. The circles show the location of the electron microprobe analysis and the color indicates the corresponding cluster from major element analysis. (D) Trace element profiles from A-B. The colors and numbers at the bottom indicate the corresponding cluster identified with major elements. (E) The same as D, but for another crystal that is not shown (from Luca Caricchi).

We are also interested in identifying the mineral populations present in the ignimbrite matrix and observing whether this varies throughout the deposit. Machine learning algorithms are becoming widely incorporated in scientific studies and they provide an opportunity to make tedious point-counting obsolete. I plan to develop a machine learning algorithm in python that can classify each mineral in a phase composition map into its associated population.

This involves creating phase composition maps (PCMs) from full thin section BSE images. Utilizing BSE scans instead of EDS scans is much faster and we can show mineral phases and compositions rather than elemental concentrations (Willis et al., 2017). This process involves calibrating BSE greyscale values with EDS mineral analyses.

We then have to train the code to classify each mineral with its correct population (based on major element clustering). The image recognition will be completed with the scikitlearn package in python. Initial formulation will involve training a supervised machine learning algorithm to identify feldspars. If successful, we will work to develop an unsupervised neural network algorithm. The unsupervised neural network algorithm will effectively create clusters automatically via self-organizing maps and then it will tag each mineral with its identified population.

Hot Spot Cluster Analysis in ArcGIS:

We plan to utilize a hot-spot clustering analysis (Ripley's k-function) in ArcGIS in order to see whether fiamme populations are constrained to specific regions. This multi-distance spatial cluster analyses investigates the clustering of features over a defined study area. We believe that this may provide insight on the pre-eruptive magmatic architecture. For example, if the Ora system had two laterally juxtaposed magma chambers that were tapped by different eruptive conduits, we would expect to find two separate clusters with discrete fiamme populations. Integrating spatial statistics and petrology can help us determine how magma was stored in the crust prior to eruption (single chamber, multiple chambers?) and the approximate locations of storage in the x-y plane. If we

combine our previously obtained Rhyolite-MELTS data to this model, we can elucidate pre-eruptive storage conditions in three-dimensions.

Sequence Stratigraphy and Ar-Ar Age Dating:

This aspect of my project will provide a critical temporal framework for the Ora deposits and a context for understanding the spatio-temporal evolution of a giant magmatic system. Single crystal $^{40}\text{Ar}/^{39}\text{Ar}$ sanidine geochronology has the potential to distinguish episodes within supereruption-scale sequences; e.g. Kay et al. (2011) identified three volcanic pulses within the 2 Ma Cerro Galan Ignimbrite. Ages of the Permian Ora deposits cannot achieve the same absolute precision, but if they span $>\sim 50$ kyr we should be able to evaluate the eruptive tempo.

The eruption duration will be determined by obtaining $^{40}\text{Ar}/^{39}\text{Ar}$ sanidine ages for early, middle, and late deposits in both the southern and northern calderas (Fig. 5). Comparison of early-erupted sanidine ages from both calderas can reveal whether there was a break in time for the two-stage caldera collapse model suggested by Willcock et al. (2015). To determine whether the Ora supereruption can be subdivided into distinct volcanic pulses, we will sample from strata with observable changes in lithofacies and bulk rock geochemistry (Willcock et al., 2013). These variations may represent the tapping of distinct magma bodies.

We are also interested in seeing whether we can identify different phases of the eruption and the activation of new volcanic vents. Can we observe the transition from the early-erupted material (VCCR + FG) to the late-erupted material (MG + FGCP) in the stratigraphic record? Is it an instantaneous switch? Or do we find a horizon which includes all four fiamme types? The possibility that some supereruptions occurred episodically, rather than \sim instantaneously, is significant for understanding the storage and mobilization of eruptible magma in the shallow crust (e.g. Cashman et al., 2017). Evidence for multiple distinct eruptions can alter how we view the evolution and behavior of supereruption-sized magma systems through time.

A creative approach to these questions may provide insight on the eruption dynamics of the Ora Ignimbrite. Similar to extinction horizons in biostratigraphy, we may be able to observe the extinction of fiamme populations in the volcanic strata. Coupling $^{40}\text{Ar}/^{39}\text{Ar}$ ages with observed fiamme extinction trends may demonstrate whether the eruption migrated northward through time as suggested by Willcock et al (2013). If we observe a northward trend, this may suggest the sequential tapping of laterally juxtaposed magma bodies (Begu  et al., 2014). If not, this may represent the eruption of multiple magma bodies emplaced at different depths within a transcrustal magmatic system (Cashman et al., 2017).



Figure 6: Bulk rock compositions, biotite crystal content (Bt), and total crystal content (CF) variation through time. The yellow stars indicate horizons of interest that could show potential changes in eruptive material. Locations 3 and 8 are intracaldera deposits and correlate with Ora 3 and Ora 8. We have yet to sample from location 20, which is just south of the Ora 2 vitrophyre unit. Figure is modified from Willcock et al., 2015.

Methods

Sample Collection:

The samples collected during the 2017 and 2019 field seasons are shown in figure 1 and their descriptions are listed in table 1. Currently, we have a disproportionate amount of samples from the northern intracaldera. In order to reduce the sampling bias for our spatial statistics, we will obtain more samples from the southern intracaldera and southern outflow. Clustering analyses in ArcGIS requires at least 30 features to be reliable. In order to sample a significant number of fiamme, we will make fiamme mounts to maximize sampling potential and analysis speed. The ignimbrite matrix samples will be made into 75 x 50 mm billets.

Analytical Work:

We plan to obtain EDS scans of individual mineral grains and corresponding BSE images using the Tescan VEGA3 SEM at Vanderbilt University. We will also collect trace element profiles using line scans with consistently spaced laser pits from the ThermoFisher iCAP Qc quadrupole ICPMS at Vanderbilt University.

Hierarchical Clustering and Machine Learning in Python:

The skills to develop code for this project will be gained from taking DataCamp online courses and also by utilizing the abundant wealth of information on the internet. The hierarchical clustering algorithm will effectively calculate the Euclidean distance between all the points and store them in a proximity matrix. Points with the smallest distances separating them will be merged into clusters until only one single cluster is left. A dendrogram is then created and the optimum number of clusters is determined visually by intersecting the tallest vertical line in the dendrogram.

The machine learning algorithm will be trained to classify minerals into these pre-defined clusters based on major element data from an EDS map. In order to test the efficacy of this method, we will first test it on a large thin section of bulk tuff from the vitrophyre. If we can estimate the bulk eruptive material here, we can apply the methodology to a single stratigraphic section. If successful, we can apply this to a stratigraphic horizon in another part of the caldera and see if we observe similar results.

Spatial Statistics in ArcGIS:

Proficiency in ArcGIS was gained from an Advanced Spatial Analysis course taken in 2018 at Vanderbilt University. ArcGIS skills will be refined by participating in an ESRI MOOC titled Spatial Data Science: The New Frontier in Analytics which runs from February 12-April 9, 2020. The Spatial Analysis Research Laboratory (SARL) at Vanderbilt University will also provide expertise in GIS if any roadblocks may arise.

Ar-Ar Dating:

Ar-Ar sanidine ages will be obtained from the New Mexico Geochronology Research Laboratory (NMGRl) at New Mexico Tech. Sanidines will be separated, irradiated, and prepared for analysis using a Thermo Scientific Helix Multicollector. Funding for Ar-Ar dating will hopefully be provided by the Awards for Geochronology Student Research 2 (AGeS2) grant which will provide support for one year in 2020.

Timeline

Year 1 (2019-2020):

Year 2 (2020-2021):

Year 3 (2021-2022):

Broader Impacts

Our community has developed novel methods to assess the records preserved within individual crystals in order to understand the evolution of magmatic systems. With the expansion of open source science, we have capitalized on collaboration and creating workflows that are accessible to all. This proposed project will develop python scripts that will automate statistical analyses and tedious point counting to further the open science revolution!

Developing open-source science with python with hopes of uploading to the ENKI server for distribution.

Mentoring undergraduates! Female Power!!

References
