

Genomic architecture of ancestry sorting: diagnostic index, not recombination, governs its direction

Draft results summary (Module B). Methods are deferred to Supplementary Materials. All findings are descriptive: any recombination effect awaits the recombination-matched neutral null, because a non-recombining autosomal block does not have lower N_e than its surroundings under neutrality.

Results

Diagnostic index is separable from recombination and reflects reduced diversity. Across the genome the diagnostic index (DI) is essentially orthogonal to the map-interpolated recombination rate (Spearman $\rho = -0.03$) and to pooled parental minor-allele frequency ($\rho = -0.02$), making it a genuinely independent predictor of sorting. Its dominant correlate is *low* within-species diversity ($\rho = -0.46$ with π), not elevated absolute divergence ($\rho = +0.13$ with d_{xy}), so DI is largely a reduced-diversity differentiation signal rather than a divergence-island one. A classical differentiation-island signature—jointly elevated F_{ST} and d_{xy} , reduced π , and very large linkage blocks—is confined to the lowest-recombination decile (Fig. 1a), yet DI is flat across recombination and does not track it. DI and recombination are therefore separable axes.

Sorting is depleted, not enriched, in low-recombination regions. At the independent-unit level, the fraction of units that sort is *lowest* in the least-recombining decile (0.06 versus ~ 0.42 elsewhere; Fig. 1b), because those regions consist of a few very large linkage blocks whose consensus washes out. The higher SNP-level value in that decile (0.17) relative to the unit level (0.06) is spatial pseudo-replication: adjacent, redundant SNPs in a single low-recombination block. Consistently, sorting magnitude *increases* with recombination (prop.fixed $\sim$ recombination, standardised slope $+0.05$, $z = 117$). Sorting is thus not concentrated where recombination is low—contrary to a naive linked-selection expectation, and consistent with neutral book-keeping (fewer independent units per low-recombination block) pending the null.

Direction is governed by diagnostic index; recombination adds only a weak independent pull. In a logistic model of sorting direction among unidirectional units, $P(\text{aquilonia}) \sim \text{DI} + \text{recombination} + \text{parental MAF} + \log(\text{block size})$, diagnostic index dominates (standardised coefficient $+1.46$, $z = 125$): high-DI units fix toward *F. aquilonia*, low-DI toward *F. polycytena*, reproducing the Module A polarity. Recombination contributes a small but significant independent effect in the opposite-to-expected-noise direction—low recombination leans *F. aquilonia* after DI, parental MAF and size are controlled (coefficient -0.09 , $z = -7.6$)—a modest linked-selection lead. It is, however, roughly sixteen-fold weaker than DI, and at the raw unit level direction is nearly flat across recombination (Fig. 1c); the stronger low-recombination *F. aquilonia* excess seen at the SNP level is largely pseudo-replication that flattens once independent units are counted.

Outlook. Module B establishes that the predictable polarity of ancestry sorting is organised by locus diagnosticity, a reduced-diversity axis orthogonal to recombination, and that sorting is depleted rather than enriched in low-recombination regions once pseudo-replication is removed. A weak residual low-recombination pull toward *F. aquilonia* survives control for DI—a lead for the intrinsic (linked-selection / incompatibility) hypothesis. Adjudicating it requires the recombination-matched, haplodiploid neutral null; the intrinsic and extrinsic drivers are then discriminated, respectively, by

among-population linkage disequilibrium between unlinked diagnostic units and by overlap with climate genotype–environment outliers.




Figure 1: **Genomic architecture of sorting.** (a) Standardised relative differentiation (F_{ST}), absolute divergence (d_{xy}) and within-species diversity (π) across recombination deciles: the differentiation-island signature is confined to the lowest-recombination decile. (b) Fraction of loci sorted versus recombination at the SNP level and at the independent-unit level; the unit-level collapse in low recombination is the removal of spatial pseudo-replication. (c) Direction of unidirectional sorting (fraction fixing toward *F. aquilonia*) versus recombination at the unit level: the diagnostic-index polarity is not mirrored by a recombination gradient.