

# Developments of Corey-Fuchs Reaction in Organic and Total Synthesis of Natural Products

Majid M. Heravi\*, Shima Asadi, Niousha Nazari and Boshra Malekzadeh Lashkariani

Department of Chemistry, School of Science, Alzahra University, PO Box 1993891176, Vanak, Tehran, Iran

**Abstract:** Corey-Fuchs olefination is a two-step reaction, involving the reaction of an aldehyde and tetrabromomethane ( $\text{CBr}_4$ ) in the presence of the triphenylphosphine ( $\text{Ph}_3\text{P}$ ) to synthesize 1,1-dibromoolefins. In the second step, the conversion of the dibromoolefins to alkynes occurred in the presence of  $n\text{-BuLi}$  as the base. This transformation was first discovered by E. J. Corey and P. L. Fuchs in 1972. Due to the importance of terminal alkynes in the synthetic organic chemistry, the Corey-Fuchs reaction has found many applications in organic transformations; particularly in the total synthesis of several biologically active natural products, often as starting materials.

**Keywords:** Corey-Fuchs, Olefination, 1,1-Dibromoolefins, Total synthesis, Organic synthesis, Terminal alkynes, Triphenylphosphine.



Majid M. Heravi

## 1. INTRODUCTION

The Corey-Fuchs reaction, also known as the Ramirez-Corey-Fuchs reaction is a two-step methodology, allowing the synthesis of terminal alkynes from aldehyde [1]. The first step, which was discovered by Desai, McKelvie, and Ramirez, leads to 1,1-dibromoolefins, which involves the reaction of an aldehyde with phosphine-dibromomethylenes, obtained from the reaction of triphenylphosphine ( $\text{Ph}_3\text{P}$ ) and tetrabromomethane ( $\text{CBr}_4$ ) [2]. The second step is the Fritsch-Buttenberg-Wiechell rearrangement, which involves the conversion of dibromoolefins to alkynes in the presence of  $n\text{-BuLi}$  or LDA as a base [3]. Corey and Fuchs combined these two reactions to develop a useful strategy for the transformation of aldehydes to alkynes [1].

The terminal alkynes have several uses in the synthetic organic chemistry, including many name reactions such as Sonogashira [4], Glaser [5], and Eglinton coupling reactions [6]. As an example, Sonogashira coupling is an effective C-C bond formation strategy, involving the Pd-catalyzed reaction of aryl halides with terminal acetylenes. This carbon-carbon bond formation has been widely applied in organic reactions and natural product synthesis [4]. Above all, terminal alkynes are one of the essential components of an important and versatile reaction, for the synthesis of 1,2,3-triazoles, namely Huisgen reaction [7], which was developed by K.B Sharpless and his co-workers [8] as click reaction, which attracted attentions of the synthetic organic research groups, for the regioselective synthesis of 1,2,3-triazole derivatives [9-11].

In the continuation of our interest in the recent advancement of the named reactions in the organic and total synthesis [12-19] using asymmetric synthesis [14, 20-22] and application of terminal alkynes [23-27]. In this review, we tried to underscore the applications of another important name reaction, called Corey-Fuchs reaction.

## 2. COREY-FUCHS REACTION IN ORGANIC SYNTHESIS

The Corey-Fuchs reaction occurs *via* initial reaction of triphenylphosphine ( $\text{Ph}_3\text{P}$ ) (**1**) and tetrabromomethane ( $\text{CBr}_4$ ). The


provided active reagent is triphenyl(trihalomethyl)phosphonium halide **3**. The one-carbon homologation of an aldehyde (**4**) to dibromoolefin (**5**) occurred by the brominating agent (**3**). Finally, dibromoolefin was treated with  $n\text{-BuLi}$  to produce a terminal alkyne (**8**) (Scheme 1) [28].

The Corey-Fuchs reaction was employed in the preparation of ( $\pm$ )-4-ethynyl[2.2]paracyclophane ( $\text{PCP-C}\equiv\text{CH}$ ) (**12**) and 4,16-diethynyl[2.2]paracyclophane (**14**). The racemic terminal alkynes (**13** and **14**) were prepared *via* a two-step synthesis [1]. In the first step, interaction of zinc,  $\text{CBr}_4$  and  $\text{PPh}_3$  followed by addition of the formyl derivatives **9-10** gave the dibromoolefins **11** and **12** with 90-95% yields. The desired Corey-Fuchs products (**13** and **14**) were obtained by the treatment of **11** and **12** with  $n\text{-BuLi}$  in THF, followed by hydrolysis, in 80-90% yields. Furthermore, the ethynyl derivative **13** was coordinated to various mono- and dinuclear complexes and played a key role in a carbon-carbon coupling reaction (Scheme 2) [29].


Corey-Fuchs reaction is a prominent transformation using strong bases such as  $\text{BuLi}$  for the conversion of an aldehyde **4** by the chain extension into a 1,1-dibromovinyl derivative **15**, which leads to an alkyne. 1,1-dibromoalkenes **15** which was subjected to Suzuki-Miyaura conditions using different kinds of boronic acids [30]. Graham *et al.* represented cyclobutene-1,2-bis(imidazolium) salt **17** is an effective catalyst for the one-pot tandem Suzuki-Miyaura/dehydrobromination reactions (Scheme 3). This procedure involved the treatment of 1,1-dibromoalkenes **15** and palladium(II) acetate, aryl boronic acids, and potassium *tert*-butoxide in toluene under Corey-Fuchs conditions to give the corresponding alkynes **16** in the temperature range of 65-90 °C in good to excellent yields (71-83%) [31]. The Suzuki-Miyaura treatment of a variety of boronic acids and boronic esters with tri- and tetra-substituted alkenes, (*Z*)-1-aryl- or (*Z*)-alkenyl-1-bromoalk-1-enes and internal alkynes were shown in Fig. (1) [32-35].

Charette *et al.* [36] reported the synthesis of 1-iodoalkynes **19** as an important building block applicable in the several synthetic methodologies, total syntheses, pharmaceutical, and material/polymer applications [37-41]. There are several methods for the synthesis of 1-iodoalkynes **19** such as iodination of metal acetylides [42-43], direct iodination of the corresponding terminal alkyne [44-45],


\*Address correspondence to this author at the Department of Chemistry, School of Science, Alzahra University, PO Box 1993891176, Vanak, Tehran, Iran; Tel: +98 2188044051; Fax: +98 21 88041344; E-mail: mmh1331@yahoo.com




**Scheme 1.** Mechanism of Corey–Fuchs reaction *via* preparation of dibromoolefin (**5**) and terminal alkyne (**8**).




**Scheme 2.**



**Fig. (1).**


**Scheme 3.**


Et<sub>3</sub>N-catalyzed Hunsdiecker reaction with NIS [46-47], and two step homologation/iodination sequence, starting from an aldehyde (Corey-Fuchs reaction) (Scheme 4, A-C) [1, 48-49] or AgNO<sub>3</sub>-mediated direct iodination of trimethylsilylacetylenes (D and E) [50].

**Scheme 4.** Various methodologies employed for the synthesis of 1-iodoalkynes **19**.


Charette and his group employed Corey-Fuchs iodoalkynylation conditions through one-pot homologation/double elimination from commercially available benzylic and allylic bromides **18** (Scheme 5). This reaction was extended to various benzylic bromides derivatives **18** and the results showed that the electron-donating groups provided higher yields than those performed with electron-withdrawing substituents as shown in Fig. (2).

Scott *et al.* described the synthesis of the polycyclic aromatic hydrocarbons (PAHs) containing a newly formed naphthalene ring,

**Fig. (2).**



**Scheme 5.** Synthesis of 1-iodoalkynes **19** from their corresponding benzylic or allylic bromides **18**.



**Scheme 7.**


starting from diaryl ketones. The procedure involve four steps, which begins with a Corey-Fuchs olefination of the starting ketone **20**. Then a double Sonogashira coupling [51-53] with trimethylsilylacetylene would occur and is followed by desilylation with tetra-*n*-butylammonium fluoride, which provide the parent diyne **23**. In the last step, a ruthenium-catalyzed or Pt-catalyzed cycloisomerization is performed in refluxing 1,2-dichloroethane to give 1,8,9-perinaphthothioxanthene **24** in 95% yield (Scheme 6) [54].

As shown in Scheme 7, this step works well when the distance between the atoms being joined are less than about 3.35-3.40 Å.


The Corey-Fuchs olefination was used for several xanthenone **25**, leading to dibromoalkene **26** with subsequent Sonogashira coupling and transition metal-catalyzed double ring closures which gave cyclized product **27** (Scheme 8) [54].

The Corey-Fuchs reaction of diaryl ketones **25a** led to dibromo olefination **26a**, followed by Suzuki or Suzuki/Stille coupling leading to novel TAE fluorophores with tunable fluorescence (**28** and **29**). The products of this reaction, showed the aggregation-induced emission characteristic (Scheme 9) [55].


In 2014, Kantevari *et al.* reported a novel method for the generation of dibenzo[*b,d*] furan-1,2,3-triazole conjugates **33** using click chemistry and Corey-Fuchs reaction (Scheme 10) [56]. Dibenzo[*b,d*]furan-2-carboxaldehyde **30** as starting material was prepared by the formylation of dibenzofuran. In the next step, 2-




**Scheme 6.**




**Scheme 8.**




Scheme 9. Synthetic Route of the TAE Compounds.



Scheme 10. Synthesis of dibenzo[b,d]furan-1,2,3-triazole 33.



Scheme 11.



Scheme 12.


(2,2-dibromovinyl)dibenzo[*b,d*]furan **31** was synthesized in the presence of  $\text{CBr}_4$  and  $\text{PPh}_3$  in high yield. Noticeably, the conversion of **31** to 2-ethynyl-dibenzo[*b,d*]furan **32** under standard conditions of Corey-Fuchs reaction [57-59] failed. The further experiments revealed the optimized conditions for this reaction is using *n*-butyl lithium (1.6 M) in THF at  $0^\circ\text{C}$  followed by 3h stirring of the reaction mixture at ambient temperature. Finally the desired products were obtained *via* Huisgen's (3+2) cycloaddition using  $\text{CuSO}_4$  catalyst, sodium ascorbate in *t*-butanol and water to give dibenzo[*b,d*]furan-1,2,3-triazole **33**.

Another application of Corey-Fuchs reaction was reported by König *et al.* in the synthesis of alkyne-substituted porphyrins **36** as a building block for the synthesis of extended photoactive porphyrin arrays. In this procedure, 2-formyl-5,10,15,20-tetraphenylporphyrin **34** was used as a starting material and the desired product **36** was obtained in two steps. First, a Corey-Fuchs reaction con-

verts **34** to the *gem*-dibromoalkene **35** under standard conditions using a large excess of  $\text{CBr}_4$ , triphenylphosphine and zinc [60]. Using trifluoroacetic acid leads to demetallation of the zinc porphyrin **35a** to **35b** [61] then the introduction of  $\text{BuLi}$  caused the rearrangement of **2a** to the corresponding product **36** with overall yield 53% over two steps (Scheme 11) [62].

The product **36** can be applied in palladium-catalyzed coupling reactions with aryl iodides to give the extended photoactive porphyrin arrays under standard conditions [63-64].

Sankararaman and his group disclosed a new multistep system for the preparation of dehydrobenzoannulenes **41** (DBAs), which known as benzocyclynes. DBAs are in great interest because of their applications as starting materials for new forms of carbon allotropes [65], building blocks for 2D carbon networks [66-67], optoelectronic materials [68-69], and as sensitive probes for the investigation of tropicity of the annulene ring [70-71].



Scheme 13.

In this multistep system, the Corey-Fuchs methodology was used in the conversion of polyaldehyde precursors **38** to poly terminal alkynes **40**. In addition, for the synthesis of final pyrene-based dehydrobenzoannulenes **41**, Sonogashira and oxidative coupling (Eglinton coupling) were applied (Scheme 12) [72].

Polin and co-workers disclosed a new multicatalytic system involving, both Corey-Fuchs olefination and hydrolysis for the synthesis of 5-ethynyl-2,2'-bipyridine **46** from 2,2'-bipyridine-5-carbaldehyde **44** as outlined in Scheme 15. 5-Methyl-2,2'-bipyridine **42** was used as a common starting material, which was converted to **44** conversion of the methyl derivative (Scheme 13) [73].

Neidlein *et al.* [74] employed a Corey-Fuchs procedure in the reaction of cyclic diketones **47** and **51** to have access to dibromoalkenes **48** and **52** respectively. These products in addition to **55** underwent an alkynylation reaction to give geminal enediynes **49a-e**, **53a-e**, and **56a-c**. The results showed that when the backbone of the precursor was cyclohexane, which could be oxidized with DDQ (2,3-dichloro-5,6-dicyano-1,4-benzoquinone) leading to the *p*-quinoid systems **50a,b** (Scheme 14).

In the past decade, triptycene derivatives have received considerable attention for their applications in the supramolecular chemistry [75-76], materials science [77-79], and synthetic molecular machines [80-83] with unique three dimensional rigid structures.

As illustrated in Scheme 15, 2-formyltriptycene **57** could be conveniently converted into 2-ethynyltriptycene **59** by Corey-Fuchs reaction in two steps with 59% total yield [84].

3-Deoxy-2-ulonic acids is an important class of acidic monosaccharides and have specific roles in biological systems [85-86]. After oxidation to the corresponding aldehyd while having alcohol **60b** on hand, it underwent Corey-Fuchs reaction to create the de-

sired alkyne **61b**. The latter can be used for the synthesis of 3-deoxy-D-*manno*-octulosonic acid (KDO) as a key core structure of lipopolysaccharides (LPSs) in several steps (Scheme 16) [87].


The same sequence, which is used for KDO, the total synthesis of 3-deoxy-D-*arabino*-2-heptulosonic acid (DAH) **65** is an intermediate in the biosynthesis of amino acids. In a key step, the Corey-Fuchs alkynylation of alditol **63** turns into alkyne **64** and was employed (Scheme 17) [88].

Ferrocene alkynes are the building blocks of some natural products [89] and valuable ligands for material science and organometallic synthesis [90].


A Corey-Fuchs methodology was employed for the synthesis of novel functionalized ferrocene alkyne Ph<sub>2</sub>P(S)fcC≡CH (**68**; fc = ferrocene-1,1'-diyl) from Ph<sub>2</sub>P(S)fcCHO **66**. The latter can be converted to the mercury(II) acetylide Hg(C≡CfP(S)Ph<sub>2</sub>)<sub>2</sub> **69** and trimetallic Co<sub>2</sub>Fe complex [(μ-η<sup>2</sup>:η<sup>2</sup>-1)(Co(CO)<sub>3</sub>)<sub>2</sub>](Co-Co) **70** (Scheme 18) [91].

Hafner *et al.* developed a synthetic route of mono- and poly-ethynylated azulenes **73a-h**, which employs a versatile procedure for the new carbon-carbon bond formation. The classical Corey-Fuchs reaction was used as a key step. Azulenes ethynylated in the five-member ring can be obtained on a multigram scale via Pd-catalyzed cross-coupling of iodo- or bromoazulenes by trimethylsilyl-acetylene followed by desilylation.


The reaction of the corresponding 5,7-diformylazulene **71** with triphenylphosphine and tetrabromomethane in dichloromethane furnished, after chromatography on alumina (BII-III) using *n*-hexane as an eluent, greenish blue crystals of the tetrabromodiolfin **72** was obtained in 70% yield. The latter could be converted into the 5,7-diethynylazulene **73h** (86%) upon treatment with 6 equiv. of LDA followed by hydrolysis (Scheme 19) [92].



Scheme 14.




Scheme 15.



i) 1)  $(\text{COCl})_2$ , DMSO,  $\text{Et}_3\text{N}$ ,  $\text{CH}_2\text{Cl}_2$ ,  $-78^\circ\text{C}$ , 2 h;  
 2)  $\text{CBr}_4$ , TPP,  $\text{Et}_3\text{N}$ ,  $\text{CH}_2\text{Cl}_2$ ,  $0^\circ\text{C}$ , 2 h;  
 3)  $\text{EtMgBr}$ , THF,  $-78$  to  $0^\circ\text{C}$ , 1 h;  
 for 61a 65% over three steps and 61b 70% over three steps


Scheme 16.






(ii) 1) (COCl)<sub>2</sub>, DMSO, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>, -78 °C, 2 h; 2) CBr<sub>4</sub>, TPP, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>, 0 °C, 1 h;  
3) EtMgBr, THF, -78 to 0 °C, 1 h, 60% over three steps

Scheme 17.




Scheme 18.




(i) PPh<sub>3</sub>, CBr<sub>4</sub>, CH<sub>2</sub>Cl<sub>2</sub>, rt; (ii) 6 equiv. LDA, THF, -90 °C, rt

Scheme 19.




(i) (Boc)<sub>2</sub>O, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>, rt, 99%; (ii) IBX, EtOAc, 75 °C, 95%; (iii) **78a**: CBr<sub>4</sub>, PPh<sub>3</sub>, Et<sub>3</sub>N, THF, -78 °C to rt, 87%. **78b**: CHI<sub>3</sub>, PPh<sub>3</sub>, *t*-BuOK, THF, rt, 68%; (iv) *n*-BuLi, THF, -78 to 0 °C then I<sub>2</sub>, 95-99%; (v) *n*-Bu<sub>3</sub>SnH, Pd(PPh<sub>3</sub>)<sub>4</sub>, toluene, rt, **79a**: 82%, **79b**: 39%; (vi) CHI<sub>3</sub>, PPh<sub>3</sub>, *t*-BuOK, THF, rt, 40%; (vii) *o*-NBSH, Et<sub>3</sub>N, *i*-PrOH-THF, rt, 73%; (viii) NaI, CuI, (CH<sub>3</sub>NHCH<sub>2</sub>)<sub>2</sub>, *n*-BuOH, 120 °C, 47%; (ix) TFA, CH<sub>2</sub>Cl<sub>2</sub>, rt, **80a**: 92%, **80b**: 94%.

Scheme 20.




Scheme 21.

Fig. (3). (-)-cytisine **74** and model lactam **75**.


In 2004, a novel lactam annulation on a model compound (lactam **75**) has been developed by Rouden *et al.* (Fig. 3). This compound was established from (2-hydroxyethyl)-piperidine **76** in a linear synthesis involving a Corey-Fuchs olefination **81** of the aldehyde by a selective reduction. *o*-Nitrobenzenesulfonylhydrazide (*o*-NBSH) was applied as diimide equivalent for an iodoalkyne into a (*Z*)-iodopropene piperidine (Scheme 20) [93].

Brown *et al.* reported the preparation of a versatility of dibromoolefins (DBOs). These reactions proceeded in two steps; the first one was oxidative cleavage of alkene **83** by ozonolysis of esters which can be combined with a Corey-Fuchs type dibromoolefination. The second step involves using phosphorous ylide to produce alkene **85** (Scheme 21) [94].

Another application of Corey-Fuchs-type sequence was described by Pajkert and his co-worker. This methodology led to novel alkynes with a difluoromethylenephosphonate function using (diethoxyphosphoryl)difluoroacetic aldehyde **86** as the precursor. The first step the Corey-Fuchs-type reaction of corresponding aldehyde **86** led to (3,3-dibromodifluoroallyl)phosphonate **87**. Then the dehydrobromination of intermediate **87** occurred in the presence of potassium *tert*-butoxide to create bromoalkyne **88**. For the first time, the reaction **87** with lithium base led to the preparation of ((diethoxyphosphoryl)difluoropropynyl)lithium **89**. The compound **89** can be used as a valuable reagent for the synthesis of different propargyl phosphonates by the treatment with various electrophiles



Scheme 22.


Scheme 23. Degradation product **93** observed in the preparation of **91** (variable amounts).

such as aldehydes, ketones, triflates, chlorophosphines, and chlorosilanes (Scheme 22) [95].

The synthesis of ethynylglycine synthon **91** as a valuable material is in great demand. There are two common pathways for the generation **91**, as illustrated in Scheme 25. The first one is known as the Seyferth-Gilbert strategy and the second method is the Corey-Fuchs reaction. The synthesis of ethynylglycine synthon **91** via Corey-Fuchs reaction occurred by the addition of BuLi **90** fol-



lowed by preparation of dibromovinyl **92** as an intermediate to give the corresponding alkyne **91**. The ratio of main product **91** is higher than by-product **93** and the amount of the by-product increased by introducing a large excess of base for longer time (Scheme 23) [96-100].

Nierengarten and his co-workers synthesized branched conjugated systems with a terminal alkyne function via Corey-Fuchs dibromoolefination. This reaction was carried out by treatment of



- (i) triisopropylsilylacetylene,  $\text{PdCl}_2(\text{PPh}_3)_2$ ,  $\text{CuI}$ ,  $\text{PPh}_3$ ,  $\text{Et}_3\text{N}$ , THF (98%);  
(ii)  $\text{CBr}_4$ ,  $\text{PPh}_3$ , Zn dust,  $\text{CH}_2\text{Cl}_2$  (99%);  
(iii) LDA, THF then  $\text{NH}_4\text{Cl}$ ,  $\text{H}_2\text{O}$  (90%);  
(iv) 1,4-diiodobenzene,  $\text{PdCl}_2(\text{PPh}_3)_2$ ,  $\text{CuI}$ ,  $\text{PPh}_3$ ,  $\text{Et}_3\text{N}$ , THF (84%).

Scheme 24.




- (ii) BSC (1.2 equiv per OH), TBAF (0.01 equiv), NMP, rt.  
(iii) (a)  $(\text{COCl})_2$  (3 equiv), DMSO (6 equiv),  $\text{CH}_2\text{Cl}_2$ ,  $-70^\circ\text{C}$  then  $\text{Et}_3\text{N}$ ,  
(b)  $\text{Ph}_3\text{PCHBr}_3$  (2.2 equiv),  $t\text{-BuOK}$  (2.1 equiv), THF, rt.

Scheme 25.


**94** with an excess of LDA. Compound **HX0 98** was prepared using  $\text{CBr}_4/\text{PPh}_3/\text{Zn}$  under the conditions introduced by Corey-Fuchs. Under this reaction conditions, dibromoolefine **96** in 99% yield was produced (Scheme 24) [101].

### 3. COREY-FUCHS REACTION IN TOTAL SYNTHESIS

Corey-Fuchs olefination as a powerful tool, has been used as a key step in total synthesis of several, naturally occurring com-





Scheme 26.



Scheme 27.

pounds. Lievre and his co-workers reported the efficient and quick synthesis of carbohydrate-derived 2,7- and 2,8-dienes [102]. This reaction started by the quick conversion of *D*-glucose **99** to the corresponding **100**, bearing carbon-carbon double bond, *via* Wittig reaction. The latter was transformed **101** in several steps. The compound **101** can be converted to the desired natural product, **102** under Corey-Fuchs reaction conditions. It's worthy to mention that in **101** the hydroxyl groups were protected as -OTMS. (Scheme 25) [103].

The synthesis of naturally occurring compound, jaspine B (pachastrissamine) **107**, has been accomplished *via* the Corey-Fuchs reaction as an important step. Jaspine B **107** was first isolated from the Okinawa marine sponge *Pachastrissa sp.* by Higa and his co-workers in 2002 [104]. In this route, compound **104** as a suitable precursor for Corey-Fuchs reaction was provided *via* a multi-step reaction, starting from *L*-serine **103**. The obtained, aldehyde **105**, then underwent Corey-Fuchs reaction to afford the respect **105**. Treatment of **105** with 2 equiv. of *n*-BuLi at -78 °C generated al-



kyne **106** which was converted to the desired natural product **107** in several steps (Scheme **26**) [105].

Terminal halogenated antazirine analogs were initially isolated from *D. fragilis* sponge and *D. fragilis*. It has been found that they show modest cytotoxicity toward HCT-116 cells [106-107]. Correspondingly, motualevic acids such as **114** as  $\omega$ -dibrominated unsaturated fatty acids were isolated and extracted from *Siliquariasporgia* sp [108].

In 2015, Sudhakar *et al.* described the total synthesis of geometrical isomer of (*E*) and (*Z*)-antazirine **111** and motualevic acids **114** as key intermediates for the total synthesis of several naturally occurring compounds. These compounds could be prepared *via* using several reactions involving Wittig olefination, Corey-Fuchs reaction, Neber reaction, amide coupling.

1,10-Decanediol **108** was employed as a readily available starting material. Initially, Corey-Fuchs reaction resulted in the transformation of **109** to **110** and ultimately the desired target **111** was synthesized. In another Corey-Fuchs strategy, **113** was created from **112** and motualevic acids **114** was provided (Scheme **27**) [109].


The main step in the first total synthesis of motualevic acids A-E **117-121**, is a new class of brominated long chain acids relied on Corey-Fuchs reaction.

Under the Corey-Fuchs reaction conditions, aldehyde **115** gave *gem*-dibromoalkene **116** in 86% yield which can be directly converted to motualevic acid E **117**. The latter is a key intermediate in the synthetic pathway of motualevic acids A **118**, C **119** and D **120** (Scheme **28**). In another synthetic strategy, aldehyde **122** yielded terminal dibromide **123** by Corey-Fuchs reaction [110] in 70% yield to be converted into the motualevic acid B **121**, which is the *Z* isomer of motualevic acid A **118** (Scheme **29**) [111].

The enantiopure C<sub>1</sub>-C<sub>11</sub> fragment of bafilomycin A<sub>1</sub> **128** which related to macrolides (16-membered lactones) was synthesized with a 4% overall yield and high stereoselectivity by Lett and co-workers [112].


For the first time, Corey and Ponde assigned the absolute configuration and stereochemistry of bafilomycin and some related macrolides based on the <sup>1</sup>H NMR coupled with the computer modeling [113] and also X-ray crystallography [114].

Formation of alkyne **127** from **124** was achieved through the methodology of Corey and Fuchs [115]. The optimal reaction conditions for obtaining the intermediate dibromoolefine **125** from **124** without deprotection of the silyl ether and further change the corresponding alcohol into the bromide were determined in the mixture of Ph<sub>3</sub>P, CBr<sub>4</sub> and NEt<sub>3</sub> [116-118]. After reduction of **125** with



- (i)  $\text{PPh}_3$  (4 equiv),  $\text{CBr}_4$  (2 equiv),  $\text{CH}_2\text{Cl}_2$ , then  $\text{NEt}_3$  (8 equiv), rt, 15 min, then  $-78^\circ\text{C}$ , 10 and  $-78^\circ\text{C}$  to rt, 7 h;  
 (ii) DIBAH (3 equiv), toluene,  $-78^\circ\text{C}$ , 3 h;  
 (iii) TBAF 1 M (1.1 equiv), THF, rt, 2 h;  
 (iv) 12, THF,  $-78^\circ\text{C}$ , then *n*-BuLi 1.6 M in hexanes (5 equiv), 1 h and  $-78^\circ\text{C}$  to rt, 2 h.

Scheme 30.



Scheme 31.


DIBAH, the hydroxyl group of product was deprotected as TMS ether group in order to avoid partial deprotection and intermolecular *trans*-silylation in the next step. Then, dibromoolefin **126** generated compound **127**, derived from **125** in 71% overall yield (four steps) (Scheme 30) [112].

The first total synthesis of (+)-violapyrone C **133**, isolated from *Streptomyces* sp., has been reported by Lee and his research group in 2014. The total synthesis, started from (*S*)-(-)-2-methylbutanol **129** which was converted to the aldehyde **130** in several steps. The latter was then subjected to the Corey-Fuchs reagents in  $\text{CH}_2\text{Cl}_2$  to


give **131** followed by lithium-bromine exchange with *n*-butyllithium and then elimination to give corresponding terminal alkyne **132** which is a key intermediate in this total synthesis (Scheme 31) [119].

Traditional organic chromophores are present in a series of organic molecules that include electron-accepting and electron-donating groups joined by a conjugated  $\pi$  system called D- $\pi$ -A structure.

In a series of H-shaped chromophores, which have two parallel and non-conjugated D- $\pi$ -A units, 9,10-dihydroanthracene **134** was



Scheme 32.



Scheme 33.

applied as molecular backbone. Lu and his co-workers synthesized compound **136** of these H-shaped chromophores *via* Corey-Fuchs reaction in a total yield of about 12.5 % [120]. The Corey-Fuchs reaction of dialdehyde **135** with triphenylphosphine and carbon tetrabromide carried out under nitrogen atmosphere at  $0^\circ\text{C}$  to yield geminated dibromo alkene chromophore **136** (Scheme 32) [121-123].

Mulzer and his co-workers suggested an efficient synthesis of the protected seco acid **139** of the antibiotic macrodiolide tartrolon **140** as an inhibitor of Gram-positive bacteria with a broad antibiotic spectrum [124], based on the Corey-Bakshi-Shibata (CBS) reduction. In this synthesis pathway, (*S*)-ethyl lactate was utilized as an appropriate precursor and alkyne **138** can be produced from aldehyde **137** *via* a Corey-Fuchs sequence in 64% yield (Scheme 33) [125].



Amphidinolide C **145** with potent bioactivities is one of the most complex members of amphidinolides, which is present in various families of biologically active macrolides and linear polyketides. Notably, they are initially isolated by extraction and fractional separation from the symbiotic marine dinoflagellate *Am-*

*phidinium* sp. [126-127]. Initially, the ring opening of epoxide **141**, provided by Sharpless asymmetric epoxidation [128], protected by PMB ether afforded **142** after several steps. The key step in the synthesis of the latter is the Corey-Fuchs reaction of **143** to produce alkyne **144** in 85% yield (Scheme 34) [129].

The alkynyl side chain in vitamin D analogues was prepared *via* Corey-Fuchs homologation, followed by alkylation. Enzymatic  $\text{C}_{17}$ -hydroxylation of 25-hydroxyvitamin  $\text{D}_3$  [25-OH- $\text{D}_3$ ], which is the main circulating form of vitamin D produced hormonally active form of vitamin  $\text{D}_3$ , calcitriol [ $1\alpha,25$ -dihydroxyvitamin  $\text{D}_3$ ,  $1\alpha,25$ -(OH) $_2$ - $\text{D}_3$ ] [130]. Calcitriol also has a wide spectrum of biological functions which regulate the calcium and phosphorus metabolism in the inhibition of the proliferation of various types of malignant cells, and in the immune response [131].

Vitamin D analogues **149a**, **150a** ( $\text{C}_{20}$  stereochemistry similar to  $20R$ -natural configuration) and their  $20$ -epimers **149b**, **150b** ( $\text{C}_{20}$  stereochemistry similar to  $20S$ -epiconfiguration) were synthesized in 20-32% and 25% overall yield from **146**, respectively (Fig. 4). In this synthesis pathway, the installation of both the alkynyl and the (*Z*)-alkenyl side chains was achieved *via* Corey-Fuchs homologation



Scheme 34. Completion of the C(1)-C(9) fragment of amphidinolide C **145**.

Scheme 35.





Fig. (4).



Scheme 36.


of **147a** and **147b** to afford the alkyne **148a** and **148b** respectively (Scheme 35) [132].

A long, but successful route for the synthesis of complex is the natural product azadirachtin **154** [133-137], isolated from the Indian neem tree *Azadirachta indica* in 1968, was reported by Ley *et al* [138].


Under the presented conditions in Scheme 36, the two epimers of **152**, which originated from compound **151**, provided the desired propargylic mesylates  $\alpha$ -**153** and  $\beta$ -**153** via Corey-Fuchs alkynyla-

tion following by homologation with *para*-formaldehyde and treatment of resulting propargylic alcohols with methanesulfonic anhydride [139].

The efficient and selective synthesis of 6,7-dehydrostipiamide **160** as a non-natural multi drug resistance reversal agent of high potency and low toxicity was achievable via Corey-Peterson olefination and Corey-Fuchs reaction over 15 steps in 23% overall yield [140-141].



Scheme 37.



*i*: (a) SO<sub>3</sub>, Py, Et<sub>3</sub>N, DMSO, CH<sub>2</sub>Cl<sub>2</sub>, 0 °C to 20 °C, 1 h (81%);  
 (b) Ph<sub>3</sub>P, CBr<sub>4</sub>, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>, 0 °C to 20 °C, 2 h (70%);  
 (c) *n*-BuLi, ClCO<sub>2</sub>Et, THF, -78 °C to 20 °C, 4 h (78%)

Scheme 38.

Aldehyde **156** of >99% *E* in 81% yield was produced after Swern oxidation of **155** and the Corey-Peterson olefination of the crudely formed aldehyde in the presence of *N*-cyclohexyl(2-triethylsilylpropylidene)imine and *s*-BuLi at -78 °C, followed by treatment with CF<sub>3</sub>COOH. Negishi *et al.* initially transformed **155** into **156** in a three steps sequential reaction, involving the Swern oxidation of **155**, Corey-Fuchs reaction/ conversion of the obtained 1,1-dibromo-1-alkene into the corresponding methylalkyne and hydrozirconation and carbonylation with *n*-BuNC. However, the conditions shown in Scheme 37 found to be more efficient for this conversion. After several steps the **156** was transformed into the intermediate **157** which under another effective Corey-Peterson olefination gave compound **158** in 80% yield (>99% *E*). The Corey-Fuchs reaction of **158** followed by acidification afforded intermediate **159** in 92% yield [142]. After several steps, the latter was converted to the desired target **160**.


The Corey-Fuchs reaction is a key step in the introduction of the alkynyl moiety in the synthesis of racemic and enantiomerically pure acetylenic ω-keto esters **164**. In the first stage, Corey-Fuchs reaction of the hydroxy derivative **162**, which derived from 2-

methyl-1,3-cyclopentanone **161**. The oxidation of the hydroxy group in **162**, afforded *bis*-dioxolane **163** as an intermediated for the total synthesis of desired acetylenic ω-keto ester **164** in 13% overall yield from **161** (Scheme 38).

To increase the length of the tether, the same reaction condition as mentioned before was used to produce the di-protected diones **167a,b**. Monodeprotection of the latter gave the acetylenic ω-keto esters **168a,b** in 43% overall yield from **165a,b** and 39% overall yield from **166a,b** (Scheme 39) [143].


In recent years, the vitamin D metabolic system shows a broad range of biological activities in the treatment of the human diseases as skin diseases, cancer, and diseases related to calcium metabolism and aberrant immunological responses [144-145]. Isotope-labeled drug molecules are useful for searching by NMR spectroscopy the formation of ligand with biological hosts as proteins and membranes.

Triple-labeled [7,9,19-<sup>13</sup>C<sub>3</sub>]-vitamin D<sub>3</sub> **175** and its 25-hydroxylated **176** and 1α,25-dihydroxylated **177** synthesized employing Corey-Fuchs reaction following a normal Wittig process on keto aldehyde **169** to achieve the C-7,19 double labeling of the A-



ii: (a)  $\text{SO}_3$ , Py,  $\text{Et}_3\text{N}$ , DMSO,  $\text{CH}_2\text{Cl}_2$ , 0 °C to 20 °C, 1 h (n = 1: 91%; n = 2: 70%);  
 (b)  $\text{Ph}_3\text{P}$ ,  $\text{CBr}_4$ ,  $\text{Et}_3\text{N}$ ,  $\text{CH}_2\text{Cl}_2$ , 0 °C to 20 °C, 12 h (n = 1: 77%; n = 2: 95%);  
 (c) *n*-BuLi,  $\text{ClCO}_2\text{Et}$ , THF, -78 °C to 20 °C, 20 h (n = 1: 86%; n = 2: 89%)

Scheme 39.



Scheme 40.

ring enyne **174** which is an intermediate in this route. The latter was transformed into desired target **175** in several other steps including functional group transformations (Scheme 40) [145].

Piper species are wide spread plant groups. They were being used as traditional medicine which have been used in China and India [146]. These materials initially were isolated from different sources *i.e.*, the roots of *Piper chaba* and exhibited hepatoprotective activity [147]. Piperamides (alkenamides) frameworks contain conjugated linear chain attached in the sides with as well as different amides and aromatic moiety.

Suresh Babu and his co-workers described the synthesis of guineensine **185**. The latter was found being secondary metabolite of the Piperaceae family. The strategy started with **178**, which upon deprotection and oxidation is converted to **180** via **179**. Subsequent Corey-Fuchs homologation of aldehyde **180** followed by dibromomethylenation with  $\text{CBr}_4$  and  $\text{PPh}_3$  furnished the dibromo olefin **181** in essentially quantitative yield. In continuation of the strategy, followed by esterification the latter was transformed to **182**. The latter was converted *via* a two-step synthesis, to the desired target **185**. (Scheme 41) [148].

PMB ether of 5,6-epoxyisoprostane E2 **191** was prepared in 11.6% overall yield by bromohydrination, aldol and Corey-Fuchs

reactions *via* several steps. The Corey-Fuchs reaction of **188** followed by debromination with *n*-BuLi gave the Li anion **190**, which was quenched to produce **189** (Scheme 42) [149].

## CONCLUSION


In the progress of the modern organic synthesis, especially in the field of total synthesis of natural product, the Corey-Fuchs reaction plays a key role in one or more steps, to provide the appropriate alkyne useful for many divers conversion, usually as the precursor. The aim of this review is to exhibit the advantages and merits of the Corey-Fuchs reaction presenting it as an invaluable strategy to the organic synthetic chemists to be considered in their design toward the total synthesis of natural products.

## CONFLICT OF INTEREST

The authors confirm that this article content has no conflict of interest.


## ACKNOWLEDGEMENTS

The authors thank the Alzahra Research Council for partial financial support.



- (i) PTSA, MeOH, 0°C, 45 min, 95%;  
(ii) IBX, DMSO, THF, rt, 98%;  
(iii) CBr<sub>4</sub>, TPP, dry DCM, 0°C, 2 h, 95%;  
(iv) *n*-BuLi, dry THF, -78°C;  
(v) chloroethyl formate, dry THF, -78°C, 2 h, 90%;  
(vi) TPP, phenol, benzene, reflux, 95%;  
(vii) LiOH, THF/H<sub>2</sub>O, rt, 12 h, 96%;  
(viii) EDC, HOBT, isobutyl amine, DCM, 0°C to rt, 8 h, 90%.

Scheme 41.



Scheme 42.

## REFERENCES

- [1] Corey, E.J.; Fuchs, P.L. A synthetic method for formyl ethynyl conversion (RCHO→RC-CH or RC-CR'). *Tetrahedron Lett.*, **1972**, 13(36), 3769-3772.
- [2] Desai, N.B.; McKelvie, N.; Ramirez, F. A New Synthesis of 1,1-dibromoolefins via phosphine-dibromomethylenes. The reaction of triphenylphosphine with carbon tetrabromide. *J. Am. Chem. Soc.*, **1962**, 84(9), 1745-1747.
- [3] Fritsch, P. Fritsch-butenberg-wiechell rearrangement. *Liebigs Ann. Chem.*, **1894**, 279, 319-324.
- [4] Chinchilla, R.; Najera, C. Recent advances in Sonogashira reactions *Chem. Soc. Rev.*, **2011**, 40(10), 5084-5121.
- [5] Glaser, C. Beiträge zur Kenntniss des Acetenylbenzols. *Ber.*, **1869**, 2(1), 422-424.
- [6] Eglinton, G.; McCrae, W. The coupling of acetylenic compounds. *Adv. Org. Synth.*, **1963**, 4, 225-328.
- [7] Huisgen, R. *1,3-Dipolare Cycloadditionen*. Wiley Interscience: New York, **1984**; Vol. 75.
- [8] Kolb, H.C.; Finn, M.G.; Sharpless, K.B. Click chemistry: Diverse chemical function from a few good reactions. *Angew. Chem.*, **2001**, 113(11), 2056-2075.
- [9] Heravi, M.M.; Hamidi, H.; Zadsirjan, V. Recent applications of click reaction in the syntheses of 1,2,3-triazoles. *Curr. Org. Synth.*, **2014**, 11(5), 647-675.
- [10] Fazeli, A.; Oskooie, H.A.; Beheshtiha, Y.S.; Heravi, M.M.; Valizadeh, H. Green and facile synthesis of 1,4-disubstituted 1,2,3-triazoles via a Click reaction of alpha-bromo ketones, [bmim]N-3 and terminal acetylenes. *Lett. Org. Chem.*, **2013**, 10(10), 738-743.
- [11] Heravi, M.M.; Fazeli, A.; Oskooie, H.A.; Beheshtiha, Y.S.; Valizadeh, H. Click synthesis of 5-substituted 1H-tetrazoles from aldehydes, hydroxylamine, and [bmim]N-3 via one-pot, three-component reaction. *Synlett*, **2012**, (20), 2927-2930.
- [12] Heravi, M.M.; Hashemi, E.; Azimian, F. Recent developments of the Stille reaction as a revolutionized method in total synthesis. *Tetrahedron*, **2014**, 70(1), 7-21.
- [13] Heravi, M.M.; Zadsirjan, V. Oxazolidinones as chiral auxiliaries in asymmetric aldol reactions applied to total synthesis. *Tetrahedron: Asymmetry*, **2013**, 24(19), 1149-1188.
- [14] Heravi, M.M.; Asadi, S.; Lashkariani, B. Recent progress in asymmetric Biginelli reaction. *Mol Divers.*, **2013**, 17(2), 389-407.
- [15] Heravi, M.M.; Faghihi, Z. McMurry coupling of aldehydes and ketones for the formation of heterocycles via olefination. *Curr. Org. Chem.*, **2012**, 16(18), 2097-2123.
- [16] Heravi, M.M.; Asadi, S.; Azarakhshi, F. Recent applications of Doebner, Doebner-von Miller and Knoevenagel-Doebner reactions in organic syntheses. *Curr. Org. Synth.*, **2014**, 11, 701-731.
- [17] Heravi, M.M.; Asadi, S. Recent applications of organocatalysts in asymmetric aldol reactions. *Tetrahedron: Asymmetry*, **2012**, 23(20-21), 1431-1465.
- [18] Heravi, M.M.; Zadsirjan, V.; Bozorgpour, Z. Applications of Mannich reaction in total syntheses of natural products. *Curr. Org. Chem.*, **2014**, 18(22), 2857-2891.
- [19] Heravi, M.M.; Hashemi, E.; Ghobadi, N. Development of recent total syntheses based on the Heck reaction. *Curr. Org. Chem.*, **2013**, 17(19), 2192-2224.
- [20] Heravi, M.M.; Zadsirjan, V. Recent advances in the application of the Oppolzer camphorsultam as a chiral auxiliary. *Tetrahedron Asymmetry*, **2014**, 25(15), 1061-1090.
- [21] Heravi, M.M.; Hajiabbasi, P. Recent advances in C-heteroatom bond forming by asymmetric Michael addition. *Mol Divers.*, **2014**, 18(2), 411-439.
- [22] Heravi, M.M.; Hajiabbasi, P.; Hamidi, H. Recent development in the asymmetric michael addition for carbon-carbon bond formation. *Curr. Org. Chem.*, **2014**, 18(4), 489-511.
- [23] Heravi, M.M.; Hashemi, E.; Shirazi Beheshtiha, Y.; Ahmadi, S.; Hosseinejad, T. PdCl<sub>2</sub> on modified poly(styrene-co-maleic anhydride): A highly active and recyclable catalyst for the Suzuki-Miyaura and Sonogashira reactions. *J. Mol. Catal. A: Chem.*, **2014**, 394, 74-82.
- [24] Heravi, M.M.; Sadjadi, S. Recent advances in the application of the Sonogashira method in the synthesis of heterocyclic compounds. *Tetrahedron*, **2009**, 65, 7761-7775.
- [25] Oskooie, H.A.; Heravi, M.M.; Behbahani, F.K. A facile, mild and efficient one-pot synthesis of 2-substituted indole derivatives catalyzed by Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub>. *Molecules*, **2007**, 12, 1438-1446.
- [26] Heravi, M.M.; Keivanloo, A.; Rahimizadeh, M.; Bakavoli, M.; Ghassemzadeh, M. Pd-Cu catalyzed heterocyclization during Sonogashira coupling: Synthesis of 3-benzylthiazolo[3,2-a]benzimidazole. *Tetrahedron Lett.*, **2004**, 45, 5747-5749.
- [27] Heravi, M.M.; Kivanloo, A.; Rahimizadeh, M.; Bakavoli, M.; Ghassemzadeh, M.; Neumu, B. Regioselective synthesis of 6-benzylthiazolo[3,2-b]1,2,4-triazoles during Sonogashira coupling. *Tetrahedron Lett.*, **2005**, 46, 1607-1610.
- [28] Cao, Z.Y. Carbon Tetrabromide. *Synlett*, **2013**, 24, 889-890.
- [29] Clément, S.; Guyard, L.; Knorr, M.; Dilsky, S.; Strohmman, C.; Arroyo, M. Ethynyl[2.2]paracyclophanes and 4-isocyano[2.2]paracyclophane as ligands in organometallic chemistry. *J. Organomet. Chem.*, **2007**, 692(4), 839-850.
- [30] Yang, W.; Gao, X.; Wang, B. Boronic acid compounds as potential pharmaceutical agents. *Med. Res. Rev.*, **2003**, 23(3), 346-368.
- [31] Rahimi, A.; Schmidt, A. Tandem suzuki-miyaura cross-coupling/dehydrobromination of 1,1-dibromoalkenes to alkynes with a cyclobutene-1,2-diylbis(imidazolium) salt as catalyst precursor. *Synthesis*, **2010**, (15), 2621-2625.
- [32] Chelucci, G.; Capitta, F.; Baldino, S.; Pinna, G.A. One-pot conversion of 1,1-dibromoalkenes into internal alkynes by sequential Suzuki-Miyaura and dehydrobromination reactions. *Tetrahedron Lett.*, **2007**, 48(37), 6514-6517.
- [33] Chelucci, G.; Capitta, F.; Baldino, S. Synthesis of internal alkynes via one-pot palladium-catalyzed and dehydrobromination reactions of 1,1-dibromo-1-alkenes. *Tetrahedron*, **2008**, 64(44), 10250-10257.
- [34] Jacobsen, M.F.; Moses, J.E.; Adlington, R.M.; Baldwin, J.E. The biomimetic synthesis of SNF4435C and SNF4435D, and the total synthesis of the polyene metabolites aureothin, N-acetyl-aureothamine and spectinabilin. *Tetrahedron*, **2006**, 62(8), 1675-1689.
- [35] Jacobsen, M.F.; Moses, J.E.; Adlington, R.M.; Baldwin, J.E. A short total synthesis of aureothin and N-acetylaureothamine. *Org. Lett.*, **2005**, 7(4), 641-644.
- [36] Pelletier, G.; Lie, S.; Mousseau, J.J.; Charette, A.B. One-pot synthesis of 1-iodoalkynes and trisubstituted alkenes from benzylic and allylic bromides. *Org. Lett.*, **2012**, 14(21), 5464-5467.
- [37] Usanov, D.L.; Yamamoto, H. Enantioselective alkylation of aldehydes with 1-haloalkynes catalyzed by tethered bis(8-quinolinate) chromium complex. *J. Am. Chem. Soc.*, **2011**, 133(5), 1286-1289.
- [38] Hofmeister, H.; Annen, K.; Laurent, H.; Wiechert, R. A novel entry to 17 $\alpha$ -bromo- and 17 $\alpha$ -iodoethynyl steroids. *Angew. Chem., Int. Ed.*, **1984**, 23(9), 727-729.
- [39] Inuki, S.; Iwata, A.; Oishi, S.; Fujii, N.; Ohno, H. Enantioselective total synthesis of (+)-lysergic acid, (+)-lysergol, and (+)-isolysergol by palladium-catalyzed domino cyclization of allenes bearing amino and bromindolyl groups. *J. Org. Chem.*, **2011**, 76(7), 2072-2083.
- [40] Schwartz, E.; Breitenkamp, K.; Fokin, V.V. Synthesis and postpolymerization functionalization of poly(5-iodo-1,2,3-triazole)s. *Macromolecules*, **2011**, 44(12), 4735-4741.
- [41] Sun, A.; Lauher, J.W.; Goroff, N.S. Preparation of poly(diiododiacetylene), an ordered conjugated Polymer of carbon and iodine. *Science*, **2006**, 312(5776), 1030-1034.
- [42] Kabalka, G.W.; Reddy Mereddy, A. Iodination of organotrifluoroborates: Synthesis of vinyl and alkynyl iodides. *Tetrahedron Lett.*, **2004**, 45(7), 1417-1419.
- [43] Luthle, J.E.; Pietruszka, J. Synthesis of enantiomerically pure cis-cyclopropylboronic esters. *Eur. J. Org. Chem.*, **2000**, 2000(14), 2557-2562.
- [44] Rajender Reddy, K.; Venkateswar, M.; Uma Maheswari, C.; Santhosh Kumar, P. Mild and efficient oxy-iodination of alkynes and phenols with potassium iodide and tert-butyl hydroperoxide. *Tetrahedron Lett.*, **2010**, 51(16), 2170-2173.
- [45] Meng, L.G.; Cai, P.J.; Guo, Q.X.; Xue, S. Direct iodination of monosubstituted aryl acetylenes and acetylenic ketones. *Synth. Commun.*, **2008**, 38(2), 225-231.
- [46] Das, J.P.; Roy, S. Catalytic Hunsdiecker reaction of  $\alpha,\beta$ -unsaturated carboxylic acids: How efficient is the catalyst? *J. Org. Chem.*, **2002**, 67(22), 7861-7864.
- [47] Naskar, D.; Roy, S. 1-haloalkynes from propiolic acids: A novel catalytic halodecarboxylation protocol. *J. Org. Chem.*, **1999**, 64(18), 6896-6897.
- [48] Hollingworth, G.J.; Sweeney, J.B. Synthesis of clevidolide and the scobinolides. *Synlett*, **1993**, 1993(07), 463-465.
- [49] Michel, P.; Gennet, D.; Rassat, A. A one-pot procedure for the synthesis of alkynes and bromoalkynes from aldehydes. *Tetrahedron Lett.*, **1999**, 40(49), 8575-8578.
- [50] Nishikawa, T.; Shibuya, S.; Hosokawa, S.; Isobe, M. One pot synthesis of haloacetylenes from trimethylsilylacetylenes. *Synlett*, **1994**, 1994(07), 485-486.
- [51] Sonogashira, K. Palladium-catalyzed alkylation: Sonogashira alkyne synthesis. In: *Handbook of Organopalladium Chemistry for Organic Synthesis*, John Wiley & Sons, Inc.: **2003**; pp. 493-529.
- [52] Sonogashira, K. Development of Pd-Cu catalyzed cross-coupling of terminal acetylenes with sp<sup>2</sup>-carbon halides. *J. Organomet. Chem.*, **2002**, 653(1-2), 46-49.
- [53] Sonogashira, K. Cross-Coupling Reactions to sp carbon atoms. In: *Metal-Catalyzed Cross-Coupling Reactions*, Wiley-VCH Verlag GmbH: **2007**; pp. 203-229.
- [54] Donovan, P.M.; Scott, L.T. Elaboration of diaryl ketones into naphthalenes fused on two or four sides: A naphthoannulation procedure. *J. Am. Chem. Soc.*, **2004**, 126(10), 3108-3112.
- [55] Zhang, G.F.; Wang, H.; Aldred, M.P.; Chen, T.; Chen, Z.Q.; Meng, X.; Zhu, M.Q. General synthetic approach toward geminal-substituted tetraarylethene fluorophores with tunable emission properties: X-ray crystallography, aggregation-induced emission and piezofluorochromism. *Chem. Mater.*, **2014**, 26(15), 4433-4446.

- [56] Yempala, T.; Sridevi, J.P.; Yogeewari, P.; Sriram, D.; Kantevari, S. Rational design and synthesis of novel dibenzo[b,d]furan-1,2,3-triazole conjugates as potent inhibitors of Mycobacterium tuberculosis. *Eur. J. Med. Chem.*, **2014**, *71*, 160-167.
- [57] Mori, M.; Tonogaki, K.; Kinoshita, A. Synthesis of 1,3-dienes from alkynes and ethylene: Acetic acid 2-methylene-3-phenethylbut-3-enyl ester. In: *Org. Synth.*, John Wiley & Sons, Inc.: **2003**.
- [58] Gibtner, T.; Hampel, J.P. Gisselbrecht, A. Hirsch end-cap stabilized oligoynes: Model compounds for the linear sp carbon allotrope carbyne. *Chem. Eur. J.*, **2002**, *68*, 408-432.
- [59] Wang, Z.; Campagna, S.; Yang, K.; Xu, G.; Pierce, M.E.; Fortunak, J.M.; Confalone, P.N. A practical preparation of terminal alkynes from aldehydes. *J. Org. Chem.*, **2000**, *65*, 1889-1891.
- [60] Callot, H.J. *Bull. Soc. Chim. Fr.*, **1973**, 3413-3416.
- [61] Lindsey, J.S.; Prathapan, S.; Johnson, T.E.; Wagner, R.W. Porphyrin building blocks for modular construction of bioorganic model systems. *Tetrahedron*, **1994**, *50*(30), 8941-8968.
- [62] König, B.; Zieg, H. Efficient synthesis and coupling reactions of (2-ethynyl-5,10,15,20-tetraphenylporphyrinato)zinc(II). *Synthesis*, **1998**, (2), 171-174.
- [63] Hegedus, L.; Schlosser, M. *Organometallics in synthesis*. Wiley: New York, **1994**; 383.
- [64] Heck, R.F. *Palladium reagents in organic syntheses*. Academic Press: London, **1985**.
- [65] Diederich, F.; Rubin, Y. Synthetic approaches toward molecular and polymeric carbon allotropes. *Angew. Chem. Int. Ed.*, **1992**, *31*(9), 1101-1123.
- [66] Haley, M.M. It takes alkynes to make a world - new methods for dehydrobenzoannulene synthesis. *Synlett*, **1998**, (6), 557-565.
- [67] Marsden, J.A.; Palmer, G.J.; Haley, M.M. Synthetic strategies for dehydrobenzo[n]annulenes. *Eur. J. Org. Chem.*, **2003**, *2003*(13), 2355-2369.
- [68] Baughman, R.H.; Eckhardt, H.; Kertesz, M. Structure-property predictions for new planar forms of carbon: Layered phases containing sp<sup>2</sup> and sp atoms. *J. Chem. Phys.*, **1987**, *87*(11), 6687-6699.
- [69] Sarkar, A.; Pak, J.J.; Rayfield, G.W.; Haley, M.M. Nonlinear optical properties of dehydrobenzo [18] annulenes: Expanded two-dimensional dipolar and octupolar NLO chromophores. *J. Mater. Chem.*, **2001**, *11*(12), 2943-2945.
- [70] Mitchell, R.H. Measuring aromaticity by NMR. *Chem. Rev.*, **2001**, *101*(5), 1301-1316.
- [71] Boydston, A.J.; Haley, M.M.; Williams, R.V.; Armantrout, J.R. Diatropicity of 3,4,7,8,9,10,13,14-octadehydro[14]annulenes: A combined experimental and theoretical investigation. *J. Org. Chem.*, **2002**, *67*(25), 8812-8819.
- [72] Joseph, A.; Venkataramana, G.; Sankaraman, S. Synthesis of dehydrobenzoannulenes with pyrene core. *J. Chem. Sci.*, **2012**, *124*(3), 597-607.
- [73] Polin, J.; Schmohel, E.; Balzani, V. Functionalization of 2,2'-bipyridines in their 4 and 5 positions. Synthesis of 5-ethynyl-2,2'-bipyridine. *Synthesis*, **1998**, (3), 321-324.
- [74] Neidlein, R.; Winter, M. Synthesis of geminal enediynes with saturated and unsaturated carbocyclic backbones by palladium-catalyzed alkylation of dibromoolefins. *Synthesis*, **1998**, (9), 1362-1366.
- [75] Chen, C.F. Novel triptycene-derived hosts: Synthesis and their applications in supramolecular chemistry. *Chem. Commun.*, **2011**, *47*(6), 1674-1688.
- [76] Jiang, Y.; Chen, C.F. Recent developments in synthesis and applications of triptycene and pentiptycene derivatives. *Eur. J. Org. Chem.*, **2011**, *32*, 6377-6403.
- [77] Hoogboom, J.; Swager, T.M. Increased alignment of electronic polymers in liquid crystals via hydrogen bonding extension. *J. Am. Chem. Soc.*, **2006**, *128*(47), 15058-15059.
- [78] Goldsmith, R.H.; Vura-Weis, J.; Scott, A.M.; Borkar, S.; Sen, A.; Ratner, M.A.; Wasielewski, M.R. Unexpectedly similar charge transfer rates through benzo-annulated bicyclo[2.2.2]octanes. *J. Am. Chem. Soc.*, **2008**, *130*(24), 7659-7669.
- [79] Gu, X.; Lai, Y.H. Triptycene: No homoconjugation effect for extending optical properties of  $\pi$ -conjugated oligomers. *Org. Lett.*, **2010**, *12*(22), 5200-5203.
- [80] Kelly, T.R.; Cai, X.; Damkaci, F.; Panicker, S.B.; Tu, B.; Bushell, S.M.; Cornelia, I.; Piggott, M.J.; Salives, R.; Caverio, M.; Zhao, Y.; Jasmin, S. Progress toward a rationally designed, chemically powered rotary molecular motor. *J. Am. Chem. Soc.*, **2006**, *129*(2), 376-386.
- [81] Yang, J.S.; Huang, Y.T.; Ho, J.H.; Sun, W.T.; Huang, H.H.; Lin, Y.C.; Huang, S.J.; Huang, S.L.; Lu, H.F.; Chao, I. A pentiptycene-derived light-driven molecular brake. *Org. Lett.*, **2008**, *10*(11), 2279-2282.
- [82] Jiang, Y.; Guo, J.B.; Chen, C.F. A new [3]rotaxane molecular machine based on a dibenzylammonium ion and a triazolium station. *Org. Lett.*, **2010**, *12*(19), 4248-4251.
- [83] Nikitin, K.; Müller-Bunz, H.; Ortin, Y.; Muldoon, J.; McGlinchey, M.J. Molecular dials: Hindered rotations in mono- and diferrocenyl anthracenes and triptycenes. *J. Am. Chem. Soc.*, **2010**, *132*(49), 17617-17622.
- [84] Li, P.F.; Chen, C.F. Synthesis, structures, and solid state self-assemblies of formyl and acetyl substituted triptycenes and their derivatives. *J. Org. Chem.*, **2012**, *77*(20), 9250-9259.
- [85] Unger, F.M. The Chemistry and Biological Significance of 3-Deoxy-d-manno-2-Octulosonic Acid (KDO). In: *Advances in Carbohydrate Chemistry and Biochemistry*; Tipson, R.S.; Derek, H., Eds.; Academic Press, **1981**; Vol. 38, pp. 323-388.
- [86] Cipolla, L.; Gabrielli, L.; Bini, D.; Russo, L.; Shaikh, N. Kdo: A critical monosaccharide for bacteria viability. *Natural Prod. Rep.*, **2010**, *27*(11), 1618-1629.
- [87] York, W.S.; Darvill, A.G.; McNeil, M.; Albersheim, P. 3-deoxy-d-manno-2-octulosonic acid (KDO) is a component of rhamnagalacturonan II, a pectic polysaccharide in the primary cell walls of plants. *Carbohydr. Res.*, **1985**, *138*(1), 109-126.
- [88] Pradhan, T.K.; Lin, C.C.; Mong, K.K.T. Formal synthesis of 3-deoxy-D-manno-octulosonic acid (KDO) and 3-deoxy-D-arabino-2-heptulosonic acid (DAH). *Synlett*, **2013**, *24*(2), 219-222.
- [89] Hüsken, N.; Gasser, G.; Köster, S.D.; Metzler-Nolte, N. "Four-potential" ferrocene labeling of PNA oligomers via click chemistry. *Bioconjugate Chem.*, **2009**, *20*(8), 1578-1586.
- [90] Long, N.J.; Williams, C.K. Metal alkynyl  $\sigma$  complexes: Synthesis and materials. *Angew. Chem. Int. Ed.*, **2003**, *42*(23), 2586-2617.
- [91] Štěpnička, P.; Císařová, I. Synthesis of [1'-(diphenylthiophosphoryl)ferrocenyl]ethyne and alkyne-metal complexes thereof. *J. Organomet. Chem.*, **2006**, *691*(13), 2863-2871.
- [92] Fabian, K.H.H.; Elwahy, A.H.M.; Hafner, K. Syntheses of mono-, di- and triethynylazulenes. *Tetrahedron Lett.*, **2000**, *41*(16), 2855-2858.
- [93] Rouden, J.; Seitz, T.; Lemoucheux, L.; Lasne, M.C. Studies toward labeling cytosine with [11C]phosgene: Rapid synthesis of a  $\delta$ -lactam involving a new chemoselective lithiation-annulation method. *J. Org. Chem.*, **2004**, *69*(11), 3787-3793.
- [94] Brown, B.A.; Veinot, J.G.C. Synthesis of dibromoolefins via a tandem ozonolysis-dibromoolefination reaction. *Tetrahedron Lett.*, **2013**, *54*(8), 792-795.
- [95] Pajkert, R.; Röschenthaler, G.V. Synthesis of ( $\alpha,\alpha$ -difluoropropargyl)phosphonates via aldehyde-to-alkyne homologation. *J. Org. Chem.*, **2013**, *78*, 3697-3708.
- [96] Branquet, E.; Meffre, P.; Durand, P.; Le Goffic, F. Synthesis of new chiral vinyl halides from L-serinal. *Synth. Commun.*, **1998**, *28*(4), 613-622.
- [97] Meffre, P.; Gauzy, L.; Perdigues, C.; Desanges-Levecque, F.; Branquet, E.; Durand, P.; Le Goffic, F. En route to optically active ethynylglycine derivatives. *Tetrahedron Lett.*, **1995**, *36*(6), 877-880.
- [98] Meffre, P.; Gauzy, L.; Branquet, E.; Durand, P.; Le Goffic, F. Synthesis of optically active  $\beta$ ,  $\gamma$ -alkynylglycine derivatives. *Tetrahedron*, **1996**, *52*(34), 11215-11238.
- [99] Reginato, G.; Mordini, A.; Degl'Innocenti, A.; Caracciolo, M. Stereoselective synthesis of (R)-(-)-2, 2-dimethyl-3-*t*-butoxycarbonyl-4-ethynyl-oxazolidine: A chiral building block for the synthesis of a new class of substituted alkynes. *Tetrahedron Lett.*, **1995**, *36*(45), 8275-8278.
- [100] Reginato, G.; Mordini, A.; Caracciolo, M. Synthetic elaboration of the side chain of (R)-2, 2-dimethyl-3-(*tert*-butoxycarbonyl)-4-ethynylloxazolidine: A new regio- and stereoselective strategy to  $\delta$ -functionalized  $\beta$ -amino alcohols. *J. Org. Chem.*, **1997**, *62*(18), 6187-6192.
- [101] Nierengarten, J.F.; Zhang, S.; Ge'gout, A.; Urbani, M. Synthesis and optical properties of isomeric branched  $\pi$ -conjugated systems. *J. Org. Chem.*, **2005**, *70*, 7550-7557.
- [102] Dolhem, F.; Smiljanic, N.; Lièvre, C.; Demailly, G. A straightforward synthesis of glyco-2,7- and 2,8-dienes. *Tetrahedron*, **2006**, *62*(33), 7756-7761.
- [103] Muzart, J. Silyl ethers as protective groups for alcohols: Oxidative deprotection and stability under alcohol oxidation conditions. *Synthesis*, **1993**, (1), 11-27.
- [104] Ledroit, V.; Debitus, C.; Lavaud, C.; Massiot, G. Jaspines A and B: Two new cytotoxic sphingosine derivatives from the marine sponge Jaspis sp. *Tetrahedron Lett.*, **2003**, *44*, 225-228.
- [105] Tang, L.J.; Lv, C.S.; Liu, D.W.; Qin, S.L.; Wang, L.P.; Li, Y.P.; Wang, X.J.; Huang, S.P. Study on synthesis of (R)-*tert*-butyl (1-hydroxybut-3-yn-2-yl)carbamate. *Adv. Mater. Res.*, **2014**, *881*-883, 438-441.
- [106] Salomon, C.E.; Williams, D.H.; Faulkner, D.J. New azacyclopropene derivatives from *Dysidea fragilis* collected in Pohpei. *J. Nat. Prod.*, **1995**, *58*, 1463-1466.
- [107] Skepper, C.K.; Molinski, T.F. Long-chain 2 H-azirines with heterogeneous terminal halogenation from the marine sponge *Dysidea fragilis*. *J. Org. Chem.*, **2008**, *73*, 2592-2597.
- [108] Keffer, J.L.; Plaza, A.; Bewley, C.A. Motualevic acids A-F, antimicrobial acids from the sponge *siliquariaspongia* sp. *Org. Lett.*, **2009**, *11*, 1087-1090.
- [109] Kadam, V.D.; Sudhakar, G. Total synthesis of motualevic acids A-F, (E) and (Z)-antiazirines. *Tetrahedron*, **2015**, *71*(7), 1058-1067.
- [110] Ishigami, K.; Katsuta, R.; Watanabe, H. Stereoselective synthesis of Sch 642305, an inhibitor of bacterial DNA primase. *Tetrahedron*, **2006**, *62*(10), 2224-2230.
- [111] Sudhakar, G.; Kadam, V.D.; Reddy, V.V.N. Total synthesis of motualevic acids A-E. *Tetrahedron Lett.*, **2010**, *51*(7), 1124-1125.
- [112] Quéron, E.; Lett, R. Synthetic studies on bafilomycin A1: Stereoselective synthesis of the enantiopure C1-C11 fragment. *Tetrahedron Lett.*, **2004**, *45*(23), 4527-4531.
- [113] Corey, E.J.; Ponder, J.W. Stereochemistry of the hygrolidins. *Tetrahedron Lett.*, **1984**, *25*(39), 4325-4328.
- [114] Baker, G.H.; Brown, P.J.; Dorgan, R.J.J.; Everett, J.R.; Ley, S.V.; Slawin, A.M.Z.; Williams, D.J. A conformational study of bafilomycin A1 by x-ray crystallography and nmr techniques. *Tetrahedron Lett.*, **1987**, *28*(45), 5565-5568.

- [115] Corey, E.J.; Fuchs, P.L. A synthetic method for formyl→ethynyl conversion (RCHO→RC-CH or RC-CR'). *Tetrahedron Lett.*, **1972**, 13(36), 3769-3772.
- [116] Marshall, J.A.; Andersen, M.W. Synthesis of 12-, 14-, and 16-membered propargylic alcohols through Lewis acid-promoted ene cyclization. *J. Org. Chem.*, **1993**, 58(15), 3912-3918.
- [117] McIntosh, M.C.; Weinreb, S.M. An approach to total synthesis of (+)-lycoricidine. *J. Org. Chem.*, **1993**, 58(18), 4823-4832.
- [118] Grandjean, D.; Pale, P.; Chuche, J. An improved procedure for aldehyde-to-alkyne homologation via 1,1-dibromoalkenes; synthesis of 1-bromoalkynes. *Tetrahedron Lett.*, **1994**, 35(21), 3529-3530.
- [119] Lee, J.S.; Shin, J.; Shin, H.J.; Lee, H.S.; Lee, Y.J.; Won, H. Total synthesis and configurational validation of (+)-violapyrone C. *Eur. J. Org. Chem.*, **2014**, (21), 4472-4476.
- [120] Shi, J.P.; Wu, D.L.; Ding, Y.; Wu, D.H.; Hu, H.W.; Lu, G.Y. Synthesis of novel H-shaped chromophores. *Tetrahedron*, **2012**, 68(13), 2770-2777.
- [121] Hwang, G.T.; Son, H.S.; Ku, J.K.; Kim, B.H. Synthesis and photophysical studies of bis-enediynes as tunable fluorophores. *J. Am. Chem. Soc.*, **2003**, 125(37), 11241-11248.
- [122] Hajbi, Y.; Neagoie, C.; Biannic, B.; Chilloux, A.; Vedrenne, E.; Baldeyrou, B.; Bailly, C.; Mérou, J.Y.; Rosca, S.; Routier, S.; Lansiaux, A. Synthesis and biological activities of new furo[3,4-b]carbazoles: Potential topoisomerase II inhibitors. *Eur. J. Med. Chem.*, **2010**, 45(11), 5428-5437.
- [123] Raghavan, S.; Sudheer Babu, V.; Sridhar, B. Mercury(II)-mediated cleavage of cyclopropylcarbinols by an intramolecular sulfinyl group as a stereo- and regioselective route to stereotriads and stereotetrads. *J. Org. Chem.*, **2010**, 76(2), 557-565.
- [124] Irschik, H.; Schummer, D.; Gerth, K.; Hofle, G.R.H. The tartrolons, new boron-containing antibiotics from a myxobacterium, sorangium cellulosum. *J. Antibiot.*, **1995**, 48, 26-30.
- [125] Mulzer, J.; Berger, M. Studies directed towards the total synthesis of the antibiotic macrolide tartrolon: EPC synthesis of the protected seco acid. *Tetrahedron Lett.*, **1998**, 39(8), 803-806.
- [126] Kubota, T.; Tsuda, M.; Kobayashi, J.I. Absolute stereochemistry of amphidinolide C. *Organic Lett.*, **2001**, 3(9), 1363-1366.
- [127] Kobayashi, J.I.; Kubota, T. Bioactive macrolides and polyketides from marine dinoflagellates of the genus amphidinium. *J. Nat. Prod.*, **2007**, 70(3), 451-460.
- [128] Heravi, M. M.; Lashaki, T.B.; Poorahmad N. Applications of sharpless asymmetric epoxidation in total synthesis. *Tetrahedron Asymmetry*, **2015** In press.
- [129] Morra, N.A.; Pagenkopf, B.L. Gram scale synthesis of the C(1)-C(9) fragment of amphidinolide C. *Tetrahedron*, **2013**, 69(40), 8632-8644.
- [130] White, P.; Cooke, N. The multifunctional properties and characteristics of vitamin D-binding protein. *Trends Endocrinol. Metab.*, **2000**, 11(8), 320-327.
- [131] Dusso, A.; Slatopolsky, E.; Brown, A.J. The role of phosphorus in the development of secondary hyperparathyroidism and parathyroid cell proliferation in chronic renal failure. *Am. J. Med. Sci.*, **1999**, 317(6), 370-376.
- [132] Riveiros, R.; Rumbo, A.; Sarandeses, L.A.; Mourino, A. Synthesis and conformational analysis of 17 $\alpha$ ,21-cyclo-22-unsaturated analogues of calcitriol†. *J. Org. Chem.*, **2007**, 72(15), 5477-5485.
- [133] Kraus, W.; Bokel, M.; Klenk, A.; Pöhl, H. The structure of azadirachtin and 22,23-dihydro-23 $\beta$ -methoxyazadirachtin. *Tetrahedron Lett.*, **1985**, 26(52), 6435-6438.
- [134] Broughton, H.B.; Ley, S.V.; Slawin, A.M.Z.; Williams, D.J.; Morgan, E.D. X-ray crystallographic structure determination of detigloyldihydroazadirachtin and reassignment of the structure of the limonoid insect antifeedant azadirachtin. *Chem. Commun.*, **1986**, (1), 46-47.
- [135] Kraus, W.; Bokel, M.; Bruhn, A.; Cramer, R.; Klaiber, I.; Klenk, A.; Nagl, G.; Pöhl, H.; Sadlo, H.; Vogler, B. Structure determination by NMR of azadirachtin and related compounds from azadirachta indica a. Juss (Meliaceae). *Tetrahedron*, **1987**, 43(12), 2817-2830.
- [136] Ley, S.V.; Lovell, H.; Williams, D.J. Chemistry of insect antifeedants from Azadirachta indica, Part 14: absolute configuration of azadirachtin. *J. Chem. Soc., Chem. Commun.*, **1992**, (18), 1304-1306.
- [137] Ley, S.V.; Doberty, K.; Massiot, G.; Nuzillard, J.M. "Connectivist" approach to organic structure determination Lsd-program assisted Nmr analysis of the insect antifeedant Azadirachtin. *Tetrahedron*, **1994**, 50(42), 12267-12280.
- [138] Butterworth, J.H.; Morgan, E.D. Isolation of a substance that suppresses feeding in locusts. *Chem. Commun.*, **1968**, (1), 23-24.
- [139] Veitch, G.E.; Beckmann, E.; Burke, B.J.; Boyer, A.; Maslen, S.L.; Ley, S.V. Synthesis of Azadirachtin: A long but successful journey. *Angew. Chem. Int. Ed.*, **2007**, 46(40), 7629-7632.
- [140] Andrus, M.B.; Lepore, S.D. Synthesis of stipiamide and a new multidrug resistance reversal agent, 6,7-dehydrostipiamide. *J. Am. Chem. Soc.*, **1997**, 119(9), 2327-2328.
- [141] Andrus, M.B.; Lepore, S.D.; Turner, T.M. Total synthesis of stipiamide and designed polyenes as new agents for the reversal of multidrug resistance. *J. Am. Chem. Soc.*, **1997**, 119(50), 12159-12169.
- [142] Zeng, X.; Zeng, F.; Negishi, E.I. Efficient and selective synthesis of 6,7-dehydrostipiamide via Zr-catalyzed asymmetric carboalumination and Pd-catalyzed cross-coupling of organozinc†. *Org. Lett.*, **2004**, 6(19), 3245-3248.
- [143] Geoffroy, P.; Ballet, M.P.; Finck, S.; Marchioni, E.B.; Marcic, C.; Miesch, M. Synthesis of racemic and enantiomerically pure acetylenic  $\alpha$ -keto esters derived from 2-methyl-1,3-cycloalkanediones and 2-methylcycloalkanones respectively. *Synthesis*, **2010**, (1), 171-179.
- [144] Zhu, G.D.; Okamura, W.H. Synthesis of vitamin D (calciferol). *Chem. Rev.*, **1995**, 95(6), 1877-1952.
- [145] Okamura, W.H.; Zhu, G.D.; Hill, D.K.; Thomas, R.J.; Ringe, K.; Borchardt, D.B.; Norman, A.W.; Mueller, L.J. Synthesis and NMR studies of 13C-labeled vitamin D metabolites. *J. Org. Chem.*, **2002**, 67(5), 1637-1650.
- [146] Parmar, V.S.; Jain, S.C.; Bisht, K.S.; Jain, R.; Taneja, P.; Jha, A.; Tyagi, O.D.; Prasad, A.K.; Wengel, J.; Olsen, C.E. Phytochemistry of the genus piper. *Phytochemistry*, **1997**, 46(4), 597-673.
- [147] Matsuda, H.; Ninomiya, K.; Morikawa, T.; Yasuda, D.; Yamaguchi, I.; Yoshikawa, M. Hepatoprotective amide constituents from the fruit of piper chaba: Structural requirements, mode of action, and new amides. *Bioorg. Med. Chem.*, **2009**, 17(20), 7313-7323.
- [148] Shingala, S.D.; Reddy, G.V.; Kumar, R.S.C.; Yadav, P.A.; Babu, K.S. Practical and efficient approach to the synthesis of guineensine. *J. Asian Nat. Prod. Res.*, **2011**, 13(2), 128-135.
- [149] Kawashima, H.; Kobayashi, Y. Synthesis of the PMB ether of 5,6-epoxyisoprostane E2 through aldol reaction of the  $\alpha$ -bromocyclopentanone. *Org. Lett.*, **2014**, 16, 2598-2601.