Index

Abyssomicin C, synthesis, 291	oxidative functionalization, 173–181
Acetoxybenziodazole, X-ray structure, 62	preparation of 1,2-diperchlorates or 1,2-ditosylates,
Acetoxybenziodoxaboroles, 65	174
1-Acetoxy-1,2-benziodoxole-3(1 <i>H</i>)-one, preparation, 54	reaction with activated iodosylbenzene, 174
α-Acetoxylation of ketones	reaction with [bis(trifluoroacetoxy)iodo]benzene, 178
Bu ₄ NI-catalyzed, 369	reaction with HTIB, 175
ESI-MS/MS study, 339	reactions with μ-oxo-bridged triflate and perchlorate,
iodobenzene-catalyzed, 338	176
mechanism, 338, 339	reactions with phenyliodine(III) sulfates, 176-178
Acetylation, of alkenes, enantioselective, 178	Alkenyl(aryl)iodonium salts, preparation
10-Acetylphenaziodonium salt, 74, 75	by the addition of hypervalent iodine reagents to
2-Acylbenzothiazoles, synthesis, 205	alkynes, 85, 86
Acylbenzyne, iodonium salts as precursors, 266	from alkynyl(aryl)iodonium salts, 85, 87
2-Acylfurans, polysubstituted, synthesis, 206	Alkenyl(aryl)iodonium triflates, preparation, by
Adamantane, tosylamination, 231	iodonium transfer, 85, 86
Aerobic catalytic system, for C-H functionalization,	Alkenylations, with alkenyliodonium salts
iodine-based, 377	of benzotriazole, 268
Aerobic oxidation of alcohols, catalyzed by PhIO ₂ , Br ₂	of carbon nucleophiles, 269
and NaNO ₂ , 364, 365	of dithiocarbamates and carbonotrithioates, 268
Agelastatins A and B, synthesis via alkynyliodonium salt,	of fluoride, 268,
272, 273	of phosphorothioates and phosphoroselenoates, 268
(–)-Agelastatin, synthesis, 291	of tellurolates and selenolates, 268
(±)-Aiphanol, synthesis, 292	of thioamides, 268
Alcohols, catalytic oxidation, by iodine(V) species, 359,	of thiocyanate, 268
360	transition metal-catalyzed, 269
Aldehydes, coupling with amines to give amides, 232	Alkenyliodonium salts
Alditols, synthesis by alkoxy radical fragmentation, 241	history, 84
Aldoximes, oxidation to nitrile oxides, 217	preparation, 84–88
Alduronic acid, lactone, synthesis, 244	reactions, 267–270
Alkenes	structural studies, 88, 89
azidation to diazides, 222, 226	X-ray structures, 88, 89
diacetoxylation, 178	Alkenyliodonium tetrafluoroborates, preparation
diamination, by (diacetoxyiodo)benzene and	from alkenylboronic acids and
bis-sulfonimides, 230	(diacetoxyiodo)benzene, 85
enantioselective diamination, 230	from alkenylsilanes and iodosylarenes, 85
mechanism of reactions with iodine(III) reagents, 174,	Alkenyl(phenyl)iodonium salts, reactivity and
175	mechanism of reactions, 267
nucleophilic epoxidation of electron-deficient alkenes,	application as alkenylating reagents, 268-270
179	α-elimination leading to alkylidenecarbenes, 269, 270

2-Alkoxyarenols, oxidative dearomatization, 193	Alkynyl cyclopentanols, ring expansions using
2-Alkoxyphenyliodonium ylides	PhI(OH)OTs-I ₂ system, 163
preparation and properties, 100	Alkynyliodonium salts
X-ray structure, 106	as cationic photoinitiators, 429
Alkowy radical fragmentation, 240–244	coupling with organocopper reagents, 270
Alkoxy radicals	mechanism of reactions with nucleophiles, 270
β-fragmentation mechanism, 244	preparation, 89–94
generation from alcohols and iodosylbenzene, 243, 244	properties and structure, 94
generation from alcohols and PhI(OAc) ₂ —iodine	reactions, 270–275
system, 239, 240	X-ray structures, 94, 95
Alkylarenes, oxidative coupling, 200	Alkynyliodonium tosylates
Alkylations, with alkyliodonium salts, 275	polymer-supported, 388
Alkylidene carbenes	preparation from HTIB and terminal alkynes, 90, 91
as intermediates in reactions of alkynyliodonium salts,	Alkynyl(phenyl)iodonium tetrafluoroborates, preparation
270, 271	from iodosylbenzene and alkynylsilanes, 93
intermolecular cyclization, 271	from organotrifluoroborates and (difluoroiodo)arenes,
intramolecular cyclization, 272	93, 94
Alkyliodides	Alkynyl(phenyl)iodonium tosylates, preparation from
oxidation, 248, 249	alkyne and HTIB, 176
oxidatively assisted hydrolysis, 249	Alkynyl(phenyl)iodonium triflates, preparation
Alkyl iodine(III) dichlorides, fluorous, 404	by iodonium transfer, 91, 92
Alkyliodine(III) derivatives, as reactive intermediates,	using 2PhIO·Tf ₂ O, 92
248–250	Alkynyl phosphates, preparation from alkynes and
Alkyliodonium salts, 96–99	[hydroxy(dialkylphosphoryloxy)iodo]benzenes, 176
Alkylperchlorates, synthesis, 249	Allochenodeoxycholic and allocholic acids, synthesis
Alkylperoxybenziodoxoles	using PhIO ₂ /Py ₂ Se ₂ , 286
preparation, 54, 55	Allyl azides, preparation by azidation of
as reagents for oxidation of sulfides, selenides and	allyltrimethylsilanes, 222
phosphines, 219	Allylic amination, 231
Alkylsulfonates, synthesis by oxidatively assisted	Allylic oxidation, of alkenes, with
nucleophilic substitution of iodine, 249	(diacetoxyiodo)benzene and <i>tert</i> -butyl
Alkynes	hydroperoxide, 182 Amberlite resin, 393, 397, 398
oxidative functionalization, 176 reaction with HTIB, 175, 176	(+)-Ambruticin S, synthesis, 298
reaction with	Amidation
[hydroxy(dialkylphosphoryloxy)iodo]benzenes, 176	of allylic silanes, 256
	copper-catalyzed, 255
Alkynyl(aryl)iodonium salts properties, 94	enantioselective, 256
spectra, 94	of hydrocarbons, 256
X-ray structures, 94	Rh(II)-catalyzed, 256
Alkynylations	of saturated C–H bonds, transition metal catalyzed,
with alkynylbenziodoxoles, 273, 274	255, 256
with alkynyl(phenyl)iodonium salts, 270	of silyl enol ethers, 256
transition metal-catalyzed, 271	of steroidal derivatives, catalyzed by
Alkynylbenziodoxathioles, 68	metalloporphyrins, 255
Alkynylbenziodoxaltholes, os Alkynylbenziodoxoles	transition metal catalyzed, 253, 255–260
from benziodoxole triflates and	Amides, azidation, 226, 227
alkynyltrimethylsilanes, 57, 58	Amidobenziodoxoles, 56, 231
preparation, from hydroxybenziodoxole, 57	Amidoiodanes
X-ray structure, 57	derivatives of cyclic imides, 109
Alkynyl cyclohexanols, ring expansions using	as intermediates in Hofmann rearrangement, 109
PhI(OH)OTs=I ₂ system, 163, 165	stable bis(sulfonylimide) derivatives, 109, 110

α -Amination, of aldehydes, using I_2 as a catalyst, 375,	1-Arylcarbazoles, synthesis, 204
376	Aryl(cyano)iodonium fluorides, preparation and X-ray
Amination and hydrazination, of non-functionalized	structure, 96
arenes, 347	Aryl(cyano)iodonium triflates, preparation and
Aminations, using hypervalent iodine reagents,	properties, 94, 95
230–232	Aryl fluorides, synthesis by thermal decomposition of
Amines, oxidation to nitriles, 216, 217	diaryliodonium tetrafluoroborates, 433
Aminobromination	Aryl glycines, synthesis, 244
by (diacetoxyiodo)benzene and bromamine-T, 158,	Aryl(heteroaryl) iodonium salts, preparation by iodonium
159	exchange reaction using aryl(cyano)iodonium
of electron-deficient olefins, 345	triflates, 80
under mechanical milling conditions, 420	Aryl iodides
Amphidinol 3, synthesis, 298	chiral, as catalysts, 341
Angucytcline antibiotics, synthesis, 298	flexible C_2 -symmetric, 353
Anilides	iodooxazoline-based, 342
acetoxylation, 182	spirobiindane-based, 341, 353
etherification, 182	Aryliodine(III) bis(pyridinium) salts
tosyloxylation, 182	preparation and X-ray structure, 109, 110
triflation, 183	oxidative properties, 111
Anilines	Aryliodine(III) carboxylates, 4, 6. See also
ipso-fluorination, 195, 196	[bis(acyloxy)iodo]arenes
oxidation to azobenzenes, 357, 358	Aryliodine(III) diacetate, chiral spirobiindane, for
oxidative dearomatization, 195, 196	enantioselective spirocyclization, 193, 194
in the presence of methanol, 196	Aryliodine(V) fluorides, preparation, properties and
Aporphines, synthesis, 199	X-ray structure, 126, 127
(+)-Aquaticol, synthesis, 293	Aryliodine(III) organosulfonates, 4, 6
Arenes, polycyclic, oxidative functionalization by	Aryliodonium imides
[hydroxy(organosulfonyloxy)iodo]benzenes, 183	ortho-substituted, 111, 112
Aromatic compounds, oxidative functionalization,	preparation, 111
182–183	Aryliodonium salts, 76–84
Aroyl azides, preparation by azidation of aryl aldehydes,	structure in solution, 84
228	X-ray studies, 82
5-Aroyl-pyrrolidinones, synthesis, 203	Aryliodonium ylides
Arsonium-iodonium ylides, reactions, 282	antibacterial activity, 439
Arylation, by diaryliodonium salts, 260–264	history, 99
of aldehydes, catalytic enantioselective α -arylation,	1-Aryl-4-methyl-1,2,4-triazolo[4,3-a]quinoxalines,
266	synthesis, 204
asymmetric α-arylation of prochiral ketones, 262	2-Aryl- <i>N</i> -methoxyethanesulfonamides, hypervalent
of carbon nucleophiles, 262–264	iodine-catalyzed cyclization, 348
of carbon surfaces, 264	Aryl(<i>N</i> -tosylimino)iodanes, X-ray structure, 113
of fluoride anion, 260, 431–440	2-Arylproline derivatives, synthesis, 205
of indoles and pyrroles, 263	1-Arylpyrrolopyrazinones, synthesis, 203
of malonates, 260, 261	Arylselenations, 235, 236
of phenols, 260–262	[(Arylsulfonyl)methyl]iodine(III) bis(trifluoroacetates),
of phosphines, 262	preparation, 38
of silyl enol ethers, 260, 261	(Arylsulfonylmethyl)iodonium salts
of sulfides, 262	preparation and properties, 98
of thiosulfonate anion, 260, 261	reactions with nucleophiles, 275
transition metal-catalyzed, 265	(Arylsulfonylmethyl)iodonium triflate, X-ray structure,
2-Arylbenzimidazoles, synthesis, 206, 420	98
Arylbenziodoxathioles, preparation and X-ray, 68, 69	Aryl-(2-thienyl)iodonium salts, application for
Arylbenziodoxoles, 59, 60	fluoridation, 435, 436

3-Aryl[1,2,4]triazolo[4,3-a]pyridines, synthesis under	Benziodoxoles
solvent-free conditions, 420	as atom-transfer reagents, 50
Asparagine, Hofmann rearrangement of, 214	definition, 50
Azacarbocyclic spirodienones, preparation by oxidative	structure, 51
dearomatization, 193	1,4-Benzodiazepin-2-ones, synthesis, 203
Azadirachtin, synthesis, 298	Benzofurans, synthesis, 205
Azidations, using hypervalent iodine reagents, 222–230	Benzoquinones, preparation from phenols or
of aromatic compounds, 223	hydroquinones, 183
of hydrocarbons, 229	Benzothiazoles, synthesis, 206, 420
by iodosylbenzene/NaN ₃ , 222	2-Benzoxazolones, synthesis, by Hofmann
mechanism, 223, 224	rearrangement, 214
by PhI(OAc) ₂ /TMSN ₃ , 222	Benzyl ethers, deprotection, 244
Azidobenziodoxoles	Benzylic C-H oxidation, by PhIO/KBr in water, 414, 415
chiral, optically active, 56	Benzylic oxidation
preparation and properties, 56	with (diacetoxyiodo)benzene and catalytic I_2 , 181
as radical azidating reagents 56, 228, 229	by IBX, 293
X-ray structure, 56	iodine(V) catalyzed, 363
Azidoidanes	Benzyne
generated in situ, as reagents for azidation, 222–225	arylation of nitriles, 266
generation at low temperature, 108	generation from phenyl benziodoxolone, 266
spectroscopy, 108	generation from
α -Azido ketones, preparation by azidation of ketones,	phenyl[2-(trimethylsilyl)phenyl]iodonium salt, 266
222, 228	O-arylation of carboxylic acids, 266
Azidophenylselenation, 236	trapping with dienes, 266
Aziridination of alkenes	Biaryl coupling, oxidative, 197–201
asymmetric, 253–255	Bifunctional catalysts
copper-catalyzed, 253, 254	bearing RuCl ₃ and iodoarene, 368, 407
hypervalent iodine-catalyzed, 345, 346	bearing TEMPO and iodobenzene, 366
hypoiodite-catalyzed, 373	ion-supported iodoarene–TEMPO, 407–409
mechanism, 255, 373	recyclable, 406–409
transition metal catalyzed, 253–255	SiO ₂ -supported, 368
D 1 10 11 4 11 6 6 7 12 6 200	Bipyrroles, preparation, by oxidative coupling of
Baylis–Hillman adducts, oxidative functionalization, 296	pyrroles, 200, 202
Benzimidazoles, synthesis, 372	Bis(acyloxyiodo)arenes
2-Benzimidazolones, synthesis, by Hofmann	chiral, for enantioselective acetylation, 178, 179
rearrangement, 214	ESI mass spectrometry and ESI-MS/MS, 10
Benziodathiazoles, 50, 51, 69	mechanism of photolysis, 430, 431
Benziodazole oxides, chiral, 122	¹⁷ O NMR spectroscopy, 43
Benziodazoles, 50, 51, 62–64	as photoiniferters, 430
derived from natural amino acids, 62	as photoinitiators, 430
preparation, 62	polymer-supported, preparation, 382–386
rearrangement 3-iminiumbenziodoxoles, 62	preparation, general approaches, 35 preparation, by ligand exchange, 38–41
self-assembly to hypervalent iodine macrocycles, 63 Benziodolium chloride, 74, 75	preparation, by rigand exchange, 38–41 preparation, by oxidation of ArI, 36–38
Benziodoxaboroles, 50, 51, 64–66	properties, 41
preparation, 64	as radical initiators, 236, 237
	reactions in aqueous solutions, 414, 415
self-assembly to macrocycle, 65–67 X-ray structure, 65	structural studies, 41–43
Benziodoxaphospholes, 69	structural studies, 41–45 structure in solution, 43
Benziodoxaphospholes, 69 Benziodoxathioles, 50, 51, 66, 68, 69	X-ray structure, 41–43
Benziodoxacoles, 50, 51	Bis(alkynyliodonium) salts, 92

Bis(azido)iodate, polymer-supported, preparation and preparation using Oxone, 38, 403 reactions, 388 preparation, using peroxytrifluoroacetic acid, 38 (+)-Biscarvacrol, asymmetric total synthesis, 194 1-[Bis(trifluoroacetoxy)iodo]-2,2,2-trifluoroethane, 4,4'-Bis(dichloroiodo)biphenyl X-ray structure, 43 preparation, 28 Bis(trifluoromethyl)benziodoxole oxides, synthesis, as recyclable chlorinating reagent, 153, 393 structure and properties, 122 Bis-iodonium salt [Bmim]BF₄, as a recyclable solvent, 419 bicyclic, 74, 75 [Bmim]Br, as solvent for reactions of IBX, 419 PQQ inhibitors, 443 [Bmim]PF₆, as a recyclable solvent, 419 Bis(methoxycarbonyl)(phenyliodinio)methanide, Bonding in iodine(III) molecules, molecular orbital synthetic applications description, 7 Bonding in iodine(V) molecules, molecular orbital C-H insertion reactions, 278 cyclopropanation of alkenes, 278 description, 7 enantioselective cyclopropanations, 278 (\pm) -Brazilin, synthesis, 292 ortho-methoxy substituted, 278 (+)-Bretonin B, synthesis, 298 Bis(methoxycarbonyl)(phenyliodonium)methanide, Brevioxime, synthesis, 298 preparation, 99, 100 Brominations, using iodine(III) reagents, 158–160 Bis(organosulfonyl)(phenyliodonium)methanides, Bromoacetoxylation, of alkenes, 161 PhIC(SO₂R)₂, preparation, 99 Bromobenziodoxoles Bis(perfluoroalkanesulfonyl)(phenyliodonium)methanides, as brominating agents, 158 preparation, 100 as reagent for allylic and benzylic brominations, 158, Bis(phenyl)iodonium chloride (BPI) as inhibitor of cytochrome P450 reductase, 442 X-ray structure, 52 physiological effects, 442 Bromodecarboxylation, of carboxylic acids, 158, 160 Bis(p-tolyl)iodonium chloride, Tol₂ICl, as food additive Bromoiodinanes, as catalytic species, 342, 343, 346 in growing cattle, 442 Bromolactonization, hypervalent iodine-catalyzed, 342, [Bis(trifluoroacetoxy)iodo]arenes as catalysts for oxidative spirocyclization, 350 4-Bromopyrrolidines, 2-substituted, synthesis, 204 preparation, by oxidation of ArI, 37, 38 [Bis(trifluoroacetoxy)iodo]benzene Cadalenquinone, synthesis, 284 abbreviations, 35 Carbamates, synthesis, by Hofmann rearrangement, common names and abbreviations, 4, 5 215 oxidation of phenols, 184-192 Carbazoles polymer-supported, preparation, 383 organocatalytic preparation, 348 preparation, from (diacetoxyiodo)benzene and synthesis, 206 trifluoroacetic acid, 40 Carbenoid reactions, of iodonium ylides, enantioselective reaction with alkenes, 178 Cu(II) or Rh(II)-catalyzed, 280, 281 reactions in water, 416, 417 Carbon-carbon bond formation, by cyclizations of reagent for Hofmann rearrangement, 213, 215 phenolic precursors, 189, 190 reagent for intramolecular amidation, 203 Carbonyl compounds α-hydroxylation by PhI(OAc)₂/KOH/MeOH, 168 reagent for oxidation of phenol ethers, 198, 199 X-ray, 41 introduction of phosphonate or phosphinate groups, 3-[Bis(trifluoroacetoxy)iodo]benzoic acid preparation, 40 oxidative α-functionalization, 168–171 as recyclable reagent, 398 α-tosyloxylation by [hydroxy(tosyloxy)iodo]benzene, 1-[Bis(trifluoroacetoxy)iodo]-1H,1H-perfluoroalkanes, 169-171 preparation, 38 Carboxylic anhydrides, α -functionalization, 170 [Bis(trifluoroacetoxy)iodo]perfluoroalkanes, Carnosic acid and carnosol, synthesis, 293 $R_fI(CO_2CF_3)_2$ Catalysis, by hypervalent iodine species, 337 as fluorous reagents, 402, 403 Catalysis, by iodine(III) species, 337-358 as oxidant, 403 Catalysis, by iodine(V) species, 358-364

Catalytic cycle	Cyclohexadienimines, 2-alkynyl, by oxidative
for α -acetoxylation of ketones, 338, 339	dearomatization of o-alkynylanilines, 196
for Hofmann rearrangement, 356, 357	Cyclohexa-2,4-dienones, 6,6-disubstituted
involving iodine(V) as active species, 360	as dienes in Diels-Alder cycloaddition, 194, 195
for metalloporphyrin-mediated oxygenations, 366	by oxidative dearomatization, 194
for oxidative bromination, 346, 347	reaction with external dienophiles, 195
Cation–radical intermediates, 15	in synthesis of bacchopetiolone carbocyclic core, 195
in oxidation of phenol ethers, 197	Cyclohexadienones, 3,3-disubstituted, synthesis, by
Cationic polymerization, 427	oxidative dearomatization, 184
C–C bond formation, hypervalent iodine-catalyzed, 354	Cyclohexadienyl(phenyl)iodonium salts, preparation, by Diels-Alder cycloadditions of alkynyliodonium
C-C coupling reactions, Bu ₄ NI-catalyzed, 374, 375	salts, 87, 88
(+)-Cephalostatin, synthesis, 298	1-Cyclohexenyl(phenyl)iodonium salts
Chalcones, rearrangement with	cyclohexyne generation, 268
[bis(trifluoroacetoxy)iodo]benzene, 210	reactions with bases, 267, 268
C-H bond oxidation, using sodium	
2-iodobenzenesulfonate as pre-catalyst and Oxone,	(–)-Decarbamoyloxysaxitoxin, synthesis, 291
363, 364	Decarboxylation, oxidative, with PhI(OAc)2-iodine
Chiral (diacetoxyiodo)binaphthyl, degenerate	system, 239
isomerization, 10	1,2-Dehydrocarborane, 266
Chlorinations, with (dichloroiodo)arenes, 152–158	Deiodination, oxidative, of alkyliodides, 249
of alkenes, 153	(±)-Deoxypreussomerin A, synthesis, 298
of aromatic compounds	15-Deoxyripostatin A, synthesis, 299
enantioselective dichlorination, 154, 155	Dess-Martin periodinane (DMP), 5, 124–126
1-Chloro-1,2-benziodoxol-3-(1 <i>H</i>)-one	dehydration of primary alcohols, 299
preparation, 51 X-ray structure, 51, 52	generation of acyl nitroso compounds, 300
as recyclable reagent for TEMPO-catalyzed	glycol C–C bond cleavage, 298
oxidations, 393, 394	oxidation of alcohols to carbonyl compounds, 296-29
1-Chloro-4-fluoro-1 H -1 λ^3 -benzo[d][1,	oxidation of 1,2-diols, 298
2,3]iodoxoborol-3-ol, 65	oxidation of 2-hydroxyporphyrins and
Chroman derivatives, synthesis by alkoxy radical	2-aminoporphyrins, 300
fragmentation, 241	oxidation of 4-substituted anilides to <i>p</i> -quinones, 301
C–N coupling reactions, hypoiodite-catalyzed, 371–373	oxidations in ionic liquids, 419
C-Nucleosides, preparation, 237	oxidative allylation of Morita-Baylis-Hillman
COX-2 inhibitors, [¹⁸ F]-labeled, 434	adducts, 299
CP-molecules, synthesis, 299	oxidative cleavage of oximes, 300
Cross-coupling reactions, of alkenyliodonium salts, 269,	oxidative cyclization of anilides, 299, 301
270	preparation, 124, 125
Cubyl ammonium salt, synthesis, by Hofmann	properties, 125
rearrangement, 215	as reagent, 282, 283, 296–302
Cyanation, by cyanobenziodoxole, 278	synthesis of 2-substituted benzothiazoles, 302
Cyanobenziodoxole	α-tosyloxylation of ketones, 299
preparation and X-ray structure, 57	X-ray structure, 125, 126
as reagent for cyanation, 277, 278	(Diacetoxyiodo)arenes
Cyanoiodonium salts, 94–96	fluorous, 404
Cyclic ethers, five- to seven-membered, synthesis by	heteroaromatic derivatives, 37
ring-enlargement, 207	ion-supported, 400, 401
Cyclizations, hypervalent iodine-catalyzed, 344, 348	optically active, preparation 36, 37
Cycloalkanols, oxidation to enones, iodine(V) catalyzed,	(Diacetoxyiodo)benzene
363	abbreviations, 35
Cycloetherification, hypoiodite-catalyzed, 370	common names and abbreviations, 4, 5

discovery, 2	Diaryliodonium tetrafluoroborates
ligand exchange with methacrylic acid, 431	preparation, 80
mechanism of palladium-catalyzed reactions, 252	X-ray structures, 84
oxidation of phenols, 184, 185, 192	Diaryliodonium tosylates, preparation
oxidations of alcohols, in the presence of TEMPO, 414, 415	from [hydroxy(tosyloxy)iodo]arenes and arylboronic acids, 78
phenylsulfonate-tagged recyclable reagent, 399	from [hydroxy(tosyloxy)iodo]arenes and
preparation, by oxidation of PhI, 36, 37	aryltributylstannanes, 78
reactions in water, 417	from [hydroxy(tosyloxy)iodo]arenes and
as reagent for Hofmann rearrangement, 213, 214	aryltrimethylsilanes, 78
as terminal oxidant in transition metal-catalyzed	Diaryliodonium triflates, preparation
oxidations, 252	from (diacetoxyiodo)arenes and arenes, 78
tetraarylphosphonium (TAP)-supported, 401	from (diacetoxyiodo)arenes and arylboronic acids, 78
X-ray structure, 41	by one-pot reaction from iodine, arenes and mCPBA,
1-(Diacetoxyiodo)-2-nitrobenzene, X-ray structure, 43	79
(Diacyloxyiodo)arenes	by one-pot reaction from iodoarenes, 78, 79
antibacterial activity, 439	Dibenziodoles, 5-aryl-5 <i>H</i> , 107
decarboxylative fragmentation, 238	Dibenziodolium salts, preparation and X-ray structure,
Diacetoxylation of alkenes, hypervalent iodine-catalyzed,	74, 75
342	Dibenzo[a,c]phenanthridines, synthesis, 203
Diacetoxylations, of alkenes by PhI(OAc) ₂ /BF ₃ ·OEt ₂ ,	Dibenzoheterocyclic compounds, synthesis, by oxidation
syn and anti, 178	of phenol ethers, 198
Diaminopimelic acid analogues, preparation, 238	(Dichloroiodo)arenes, 4, 6
2,3-Diarylbenzo[b]furans, synthesis, 206	fluorous, 404
Diaryl ethers, synthesis, 260–262	preparation, by ligand exchange, 29, 30
Diaryliodonium butyltriphenylborate salts, as	X-ray crystal structure, 30
photoinitiators, 430	(Dichloroiodo)benzene
Diaryliodonium chlorides, preparation using	adduct with tetraphenylphosphonium chloride, 30
(<i>E</i>)-chlorovinyliodine(III) dichloride, 81	chlorination of aminoacetophenone, 155
Diaryliodonium fluorides, <i>ab initio</i> computational studies	chlorination of fullerene, 156
of the extrusion reactions, 12	chlorination of ketones, 153
Diaryliodonium halides, X-ray structures, 84	chlorination of monoterpenes, 153
Diaryliodonium salts	chlorination of norbornene, 153, 154
activity against oral and dental anaerobes, 439	chlorination and oxidation of transition metal
antifungal activity, 439	complexes, 157, 158
benzyne precursors, 266, 267	chlorination of 3-substituted sydnones, 156
biocidal and antimicrobial activity, 439, 440	chlorination at sulfur, phosphorus, selenium, arsenic,
history, 2	antimony, 157
mechanism of photolysis, 427, 428	chlorination of 5,10,15-trisubstituted porphyrins, 155,
mechanism of reactions with nucleophiles, 264,	156
265	discovery, 2
mechanism of solvolysis, 265	electrophilic reactivity of PhICl ₂ , 154
photoinitiators for cationic photopolymerizations,	generation <i>in situ</i> , 157
426–431	mechanism of chlorinations, 152, 153
polymer-supported, 388	preparation, 27, 28, 30
PQQ inhibitors, 443	as reagent for chlorinations, 152
preparation, 76–81	X-ray crystal structure, 30
reactions, 260–267	3-(Dichloroiodo)benzoic acid
reagents for introduction of fluorine-18, 432	preparation, 28
transition metal mediated cross-coupling reactions,	as recyclable reagent, 393
265, 266	as a stoichiometric oxidant, 393, 394

(D) 11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
(Dichloroiodo)methylsulfones, ArSO ₂ CH ₂ ICl ₂ ,	Diphenyliodonium-2-carboxylate, see
preparation, 29	1-phenyl-1,2-benziodoxole-3(1 <i>H</i>)-one
(Dichloroiodo)toluene, as reagent for catalytic	2,3-Diphenylquinoxaline-1-oxide, synthesis, 206
asymmetric chlorinations, 153	Diselenides
(Dicyanoiodo)arenes, ArI(CN) ₂ , 107	oxidation to selenosulfonates, 219, 220
(Dicyanoiodo)benzene, preparation and properties, 107	oxidation to trifluoromethaneselenosulfonates, 220
(Dicyano)iodonium triflate, preparation and properties, 94	Disproportionation, of iodine(III) species, RuCl ₃ -catalyzed, 367, 368
(Difluoroiodo)arenes, 4 electrochemical generation, 146	Ditellurides, oxidation to arenetellurinic mixed anhydrides, 219, 220
preparation, by electrochemical fluorination, 25, 26	Diynyl(phenyl)iodonium triflates, 92
X-ray crystal structure, 26, 27	DOPA, synthesis, 296
(Difluoroiodo)benzene, preparation from PhI(OAc) ₂ and	
anhydrous tetrabutylammonium fluoride, 26	(E)-chlorovinyliodine(III) dichloride, preparation and
(Difluoroiodo)toluene, as fluorinating reagent,	properties, 30
146–152	Elemental iodine, as pre-catalyst, 375–377
electrochemical generation, 146	Enantioselective oxidative spirocyclization, with chiral
fluorination of cyclic ethers, 149	iodoarenes as catalysts, 353
fluorination of thio- and seleno-glycosides, 147	Enantioselective reactions, catalyzed by chiral
fluorinative ring-expansion, 149	iodoarenes, 338, 341, 342, 353, 354
reaction with alkynes, 150	Ent-hyperforin, synthesis, 298
reaction with terminal alkenes, 150	(–)-Epibatidine, synthesis, 262
10,11-Dihydrodibenz[<i>b</i> , <i>f</i>]iodeponium salt, 74, 75	(±)-Epimaritidine, synthesis, 385
2,3-Dihydrofuran derivatives, synthesis, 205	Epoxidation
2,5-Dihydrofurans, synthesis from	of alkenes, transition metal-catalyzed, 251
3-alkoxy-1-alkynyl-(phenyl)iodonium triflates and	of electron-deficient alkenes, 179
sodium benzenesulfinate, 272	enantioselective, catalyzed by chiral transition metal
1,3-Dihydro-1-hydroxy-3-methyl-1,2,3-	complexes, 251
benziodoxaphosphole 2-oxide, 69	organocatalytic, asymmetric, 179
5,6-Dihydroimidazo[2,1- <i>b</i>]thiazoles, 3-substituted,	of steroids, by PhIO ₂ /VO(acac) ₂ , 286
synthesis from alkynyliodonium salts, 271	Epoxides, cleavage, with periodic acid, 303, 304
Dihydrooxazole derivatives, synthesis, 206	Epoxyquinomycin B, synthesis, 301
Dihydropyran, oxidation with iodosylbenzene in water,	Erythromycin B, synthesis, 298
413, 414	Ethynylbenziodoxolones, as acetylene transfer reagents,
2,2'-Diiodo-4,4',6,6'-tetramethylbiphenyl, as catalyst,	273, 274
347, 348, 353	Eudistomin U, synthesis, 296
5,5-Dimethyl-1,3-cyclohexanedione phenyliodonium	Eddistoffill O, Sylfdiesis, 290
	Farglitazar, fluorine-18 labeled, 437
ylide as carbenoid precursor, 278–280	Favorskii rearrangement, of steroidal substrates, 209
ortho-alkoxy substituted, 280	FD-891 16-membered macrolide, synthesis, 298
in the synthesis of five-membered heterocycles, 279,	[18F]DAA1106, PET ligand, synthesis, 437
280	[18F]-fluoridation, with diaryliodonium salts
Dimethyldioxirane (DMDO)	with heteroaromatic iodonium salts, 435, 436
•	
reaction with iodoarenes, 32 reaction with iodocyclohexane, 32	mechanistic and structural studies, 439
reaction with iodocycronexane, 32	microreactor for fluoridation, 440
· · · · · · · · · · · · · · · · · · ·	in the presence of radical scavengers, 434, 435
Diphenyleneiodonium (DPI), 74. See also	regioselectivity, 432–434
dibenziodolium	[18F] fluoride, cyclotron production, 432
Diphenyleneiodonium chloride (DPI)	[¹⁸ F]-fluoroarenes
as a hypoglycaemic agent, 442	azido- or azidomethyl-substituted, 438
physiological effects, 442	meta-substituted, 438

3-[18F]-fluorobenzaldehyde, 439 Furopyrimidinones, synthesis, 204 4-[18F]-fluoroiodobenzene, 434 Furyl carbinols, synthesis, 244 Five-membered iodine heterocycles, benziodoxoles and (–)-Fusarisetin A, synthesis, 299 benziodazoles, 4, 6 [18F]-labeled radiotracers, 432, 434, 435 Gambieric acid A, synthesis, 299 ¹⁸F-labeling, of the aromatic ring, 432 Glycol-cleavage, with periodic acid, 303, 304 Fluorinations, with (difluoroiodo)arenes, see also GM222712, antifungal agent, synthesis, 289 [18F]-fluoridation Gold-catalyzed oxidations, with hypervalent iodine of alkenes, 149, 150 reagents, 253 of dithioketals, 147 Green chemistry, basic principles, 381, 413 of β-ketoesters and β-dicarbonyl compounds, 146 mechanism of fluorination of α -phenylthio esters, 147, (±)-Halichlorine, tricyclic core, synthesis via 148 alkynyliodonium salt, 272, 273 of N-protected anilines, 152 Halichondrin C, synthesis, 299 of organosulfur compounds, 146, 147 Halobenziodoxoles, 51, 52 of steroidal dienes, 150 Haloenones, synthesis by oxidative rearrangement from of terminal alkynes, 150 alkynols, 163, 164 Fluoroalkyl(aryl)iodonium salts, as electrophilic Halogen-ate(I) complexes, polymer-supported, 387, 388 fluoroalkylating reagents, 275, 276 Heliannuols B and D, synthesis, 291 Fluoroalkylation Heptavalent iodine, structural types, 3 of amino acids and peptides, 275 Hetaryliodonium chlorides, preparation using (E)-chlorovinyliodine(III) dichloride, 81 with fluoroalkyliodonium salts, 275, 276 Fluoroalkyliododichlorides, X-ray crystal structures, 30 Heteroaryl(aryl)iodonium tosylates, preparation Fluoroalkyliodonium salts, 96-98 by [3+2] cycloaddition reactions of alkynyliodonium Fluoroalkyliodonium triflimide, preparation and X-ray salts, 81, 82 structure, 98 using [hydroxy(tosyloxy)iodo]arenes, 78 Fluoroalkyl-substituted alcohols, oxidation by catalytic Heteroaryliodonium salts, biocidal and antimicrobial sodium 2-iodobenzenesulfonate and Oxone, 362 activity, 439, 440 Fluoroaryliodine(III) cyanides, preparation, properties Heteroaryliodonium salts, X-ray studies, 82 and X-ray structure, 107 Heteroaryl(phenyl)iodonium salts, as heteroaryl transfer Fluorobenziodoxole, X-ray structure, 52 agents, 262 Fluorodeiodination, of 1-iodonorbornanes, 249 Heteroaryl(phenyl)iodonium triflates, preparation from α-Fluoroketones, preparation from silyl enol ethers and vinyliodonium salt and aryllithiums, 81 difluoroiodotoluene, 146 Heterocycles 3-Fluoropyridine, fluorine-18 labeled, 436, 437 synthesis under solvent-free conditions, 420 3-Fluoroquinoline, fluorine-18 labeled, 436, 437 synthesis using [hydroxy(tosyloxy)iodo]benzene, 170 2-Fluorothiophene, preparation, 436 synthesis using I2 as a catalyst and TBHP as oxidant, Fluorous aryl and alkyl iodine(III) dichlorides, preparation, 28 Heterocyclizations, induced by hypervalent iodine Fluorous techniques, 402 reagents, 203 1*H*-1-Hydroxy-1,2,3-benziodoxathiole 3,3-dioxide, Formylcyclopentane, preparation from cyclohexene and PhIO•BF₃ or (PhIO)₃SO₃, 174 preparation and X-ray, 67, 68 Fragmentation (-)-Hispidospermidin, enantioselective total synthesis, of β -amino alcohols and α -amino acids, 206, 207 255, 256 of tertiary cyclopropyl alcohols, 206, 208 Hofmann rearrangement, 213-216 Fredericamycin A, synthesis, 298 hypervalent iodine-catalyzed, 355 Fullerene C₆₀, azidation, 228 mechanism, 215, 216 Furanones, 5-acetoxy-2(5H), synthesis by using aryl iodides as catalysts and mCPBA, 355, 356 ring-enlargement, 208, 209 using hypervalent iodine species generated in situ, Furfuryl alcohols, 4,5-disubstituted, synthesis, 205 216 Furopyrano-2-isoxazoline derivatives, synthesis, 206 using iodobenzene and Oxone, 355, 357

Homolytic reactions of λ³-iodanes, 14, 15 (–)-Hopeanol and (–)-hopeahainol A, synthesis, 293	preparation from (diacetoxyiodo)arenes and TsOH, 43,
Hydrazination, organocatalytic, 348	preparation, from iodoarenes, mCPBA and
Hydrazones	organosulfonic acids, 45
conversion to alkyltosylates, 212, 213	reactions under microwave irradiation conditions, 421
oxidation to diazo compounds, 218	reagent for Hofmann rearrangement, 213, 215
Hydroxybenziodoxaphosphole	study in aqueous solutions, 47
catalytic activity in hydrolysis of phosphate esters, 69	X-ray structure, 47
X-ray structure, 69	3-[Hydroxy(tosyloxy)iodo]benzoic acid, as recyclable
1-Hydroxy-1,2-benziodoxole-3(1 <i>H</i>)-one	reagent, 398
catalytic activity in the cleavage of phosphates and	Hyperleaving group ability, of ArI group, 260
reactive esters, 53	Hypernucleofuge, definition, 14
as tautomer of 2-iodosylbenzoic acid, 53	Hypervalent
X-ray structure, 53	bond, definition, 1
Hydroxybenziodoxoles, 53, 54	bonding, 4, 7
[Hydroxy((+)-10-camphorosulfonyloxy)iodo]benzene as reagents for enantioselective oxidation of sulfides,	bonding, in trihalide anions, 11
-	12-1-4 square-planar species, 14
220, 221	Hypervalent iodine
[Hydroxy(2,4-dinitrobenzenesulfonyloxy)iodo]benzene	application in organic synthesis, 3
(HDNIB), reactions under microwave irradiation	complexes with nitrogen heterocycles, 109, 110
conditions, 421	computational studies, 11, 12
Hydroxyiodoxolone, pK _a , 70	experimental structural studies, 9
Hydroxylation	history of research, 2, 3
of ketones by [bis(trifluoroacetoxy)iodo]benzene, 169	IUPAC nomenclature, 3, 4
metalloporphyrin-catalyzed, 250, 251	ligand coupling, 1, 13, 337
of substituted acetophenones, 169	ligand exchange, 1, 13, 337
[Hydroxy(mesyloxy)iodo]benzene	oxidative addition, 1, 337
study in aqueous solutions, 47	principles of reactivity, 12
X-ray structure, 47	reactivity, 1, 337
[Hydroxy(organosulfonyloxy)iodo]arenes	reductive elimination, 1, 13
chiral, 175	reviews and books, 2
enantioselective reactions with alkenes, 175, 176	spectroscopic studies, 9
solvent-free preparation, 44, 45, 420	typical coordination patterns, 9
structural studies, 47	X-ray crystal structures, 9
3-Hydroxy-2-oxindoles, synthesis, 205	Hypervalent iodine catalysis, discovery of, 337
Hydroxy(phenyl)iodonium ion, 367	Hypervalent iodine compounds
[Hydroxy(phosphoryloxy)iodo]arenes, solvent-free	calculations of geometry and electronic structure, 12
preparation, 420	computational studies, 7
[Hydroxy(phosphoryloxy)iodo]benzenes, 49	key structural features, 8, 9
[Hydroxy(tosyloxy)iodo]arenes	similarities with organometallic compounds, 7, 13, 337
chiral, 44	synthesis, by ligand exchange, 21
ion-supported, 400, 401	synthesis, by oxidative addition, 21
polymer-supported, preparation, 386–387	typical synthetic applications, 145
[Hydroxy(tosyloxy)iodo]benzene (HTIB)	Hypervalent iodine heterocycles, fused, 73, 74
common names and abbreviations, 4, 5	Hypervalent iodine reagents
ESI-MS and ESI-MS/MS study, 10	chiral, for enantioselective diamination, 230
ionization in solution, 10, 11	definition, 1
in oxidative rearrangements of alkenes, 210	general properties, 3
polymer-supported, 45	nomenclature, 4
potentiometric titrations, 10	reactions under microwave irradiation conditions, 421
preparation, 43–44	solvent-free preparation, 420

Hypervalent molecules, Martin–Arduengo <i>N–X–L</i>	λ ³ -Iodanes, IUPAC recommendations, 3, 4
designation, 3	key structural features, 8
Hypervalent species, definition, 4	Iodanes, μ-oxo-bridged, 47, 48. See also μ-oxo-bridged
Hypervalent twisting, 12	iodanes
Hypoiodite-catalyzed reactions, 370–373	Iodates, inorganic, as second-harmonic generation (SHG) materials, 426
IBX-amides, polymer-supported, 390	metal salts, as second-order nonlinear optical (NLO)
IBX derivatives, as catalysts for oxidation of alcohols,	materials, 426
360, 361	Iodic acid, as reagent, 302, 303
IBX-esters	for dehydrogenation of aldehydes and ketones,
polymer-supported, 390	302
as reagents for oxidation of alcohols, 287	for deprotection of ketoximes and aromatic aldoximes, 302
Imidation	
of complexes of ruthenium(II), osmium(II) and cobalt(III), 260	for deprotection of thioacetals and thioketals, 302 for deprotection of trimethylsilyl, tetrahydropyranyl
copper-catalyzed, 257	and methoxymethyl ethers, 302
enantioselective, of sulfides, 257–259	for oxidation of aromatic amines to quinones, 302
of nitrogen heterocycles, 256, 257	for oxidation of sulfides, 302
of sulfides, 256	for oxidative iodination of aromatic compounds,
of sulfoxides, 256, 257	302
of sulfur, 256	for oxidative rearrangements and couplings, 302
Imidazo[1,2-a]pyridines	Iodide anion, as pre-catalyst, 368–375
synthesis, 372	Iodinanes, general name, 3
2-substituted, synthesis from alkynyliodonium salts,	Iodination, of aromatics, alkenes and alkynes using
271, 272	IBX/I_2 , 296
Imidazoles, synthesis, 206	Iodinations, using iodine(III) reagents, 160–164
Imidazolidinone, chiral catalyst, 179, 181	Iodine
2-Imidazolines, synthesis, 372	in biological organisms, 2
Imidoiodanes, 111–114	bonding, 1
as amidating reagents, 231, 232	coordination of, in [bis(acyloxy)iodo]arenes, 43
soluble derivatives of ortho-alkoxyiodobenzenes,	discovery, 1
112	element, 1
for transfer the imido group to other elements	history, 1, 2
in transition metal catalyzed reactions, 253	oxidation states, 3
X-ray structural studies, 112, 113	pentagonal-planar coordination, 43
α-Iminonitriles, one-pot synthesis by Strecker reaction,	production, from saltpeter, 2
295	production, from underground brine, 2
(±)-Indatraline, total synthesis, 210, 211	in thyroid gland, 2
Indazolone derivatives, synthesis, 203	Iodine(VII)
Indole derivatives, N-substituted, synthesis, 204	structural types, 8
Indole-3-methylamines, synthesis, by Hofmann	synthetic applications, 303–307
rearrangement, 214	Iodine(III), active catalytic species, generation from
Indoline derivatives, synthesis, 203	iodoarenes and Oxone, 350, 357
Indolizidinones, synthesis, 203	Iodine catalytic systems, recyclable, 406–409
Inorganic iodine(III) compounds, 21–23	Iodine(III) compounds, see also hypervalent iodine
Intermolecular interaction in hypervalent iodine	application in organic synthesis, 12, 145
compounds, 9, 11	general classes, 4, 6
Intramolecular C–C cyclization, iodoarene-catalyzed,	general reactivity pattern, 145
354, 355	Iodine(V) compounds
λ^3 -Iodanes, IUPAC recommendations, 3, 4	application in organic synthesis, 13, 282–303
key structural features, 8	heterocycles, 120–126

inorganic reagents, 302, 303 polymer-supported, 389–391 reactions in water, 416, 417 recyclable, nonpolymeric, 405, 406 lodine(VII) fluoride, preparation, properties and structure, 127 lodine fluorides, 22 lodine(III) halides, 22, 23 lodine(III) halides, 22, 23 lodine(III) heterocycles aromatic character, 50 five-membered 50–71 four-membered, 50, 74 six-membered 50, 72, 73 stability, 50 lodine(IIII) oxide, 1 ₂ O ₃ , 23 lodine(III) oxide, 1 ₂ O ₃ ,	Iodine(V) compounds (Continued)	stability, 22
polymer-supported, 389–391 reactions in water, 416, 417 recyclable, nonpolymeric, 405, 406 lodine(VII) compounds, 8, 127, 128 lodine(VII) compounds, 8, 127, 128 lodine(III) moride, preparation, properties and structure, 127 lodine fluorides, 22, 23 lodine(III) halides, 22, 23 lodine(III) halides, 22, 23 lodine(III) halides, 22, 23 lodine(III) recreeycles aromatic character, 50 five-membered, 50, 74 six-membered, 50, 74 six-membered 50, 72, 73 stability, 50 lodine(III) voide, 1,03, 23 lodine-coxygen bond lengths, statistical analysis, 11 lodine pentafluoride as oxidant for carbon monoxide, 426 preparation and properties, 114 structure, 114 lodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 sucuture, 115 for thiocyanation of ketones and aromatic compounds, 303 lodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 lonic-liquid-supported, 400, 401 magnetic annoparticle-supported, 401, 402 polymer-supported, 381–389 lodine(III) rotice preparation and structure, 22 lodoaceatoxylation of alkenes, 161, 162 lodobenzenes ulfonamide, oxidation to lacetoxy-1,2-dibydro-1,3,2-benziodathiazole 3,3-dioxide, 69 2-lodobenzenesulfonamide, oxidation to lacetoxy-1,2-benziodathiazole 3,3-dioxide, 69 2-lodobenzenesulfonamide, va	inorganic, 114, 115	X-ray 22, 23
reactions in water, 416, 417 recyclable, nonpolymeric, 405, 406 lodine(VII) compounds, 8, 127, 128 lodine(VII) fluoride, preparation, properties and structure, 127 lodine fluorides, 22 lodine(III) halides, 22, 23 lodine(III) thereocycles aromatic character, 50 five-membered 50, 72, 73 stability, 50 lodine (IVVII) mixed oxide, 128 lodine(III) voides, 1 ₂ 0, 23 lodine(III) voides, 1 ₂ 0, 23 lodine(III) voides, 1 ₂ 0, 23 lodine(III) oxide, 1 ₂ 03, 23 lodine pentalional studies, 22 properties, 22 stability, 22 X-ray, 22 lodine(III) tris(carboxylates), preparation and structure, 23 lodoacetoxylation of alkenes, 161, 162 lodobenzene as excellent leaving group, 14 magnetic nanoparticle-supported, 402 polystyrene-supported, as catalyst, 406 lodobenzenes, ortho-substituted, as organocatalysts, 342 2-lodobenzenesulfonamide, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 3,3-dioxide, 69 2-lodobenzenesulfonamide, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 1-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 1-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 1-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 1-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 1-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 1-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 1-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 1-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 1-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 1-acetoxy-1,2-dihydro-1,3,2-benziodat	inorganic reagents, 302, 303	
recyclable, nonpolymeric, 405, 406 lodine(VII) compounds, 8, 127, 128 lodine(VII) fluoride, preparation, properties and structure, 127 lodine fluorides, 22 lodine(III) halides, 22, 23 lodine(III) halides, 22, 23 lodine(III) heterocycles aromatic character, 50 five-membered 50, 72, 73 stability, 50 lodine (VVII) mixed oxide, 128 lodine(III) voxides, I ₂ O ₂ , 33 lodine(III) voxides, I ₂ O ₃ , 23 lodine(III) voxides, I ₂ O ₄ and I ₄ O ₉ , 115 lodine pentafluoride as fluorinating reagent, 303 as a industrial fluorinating reagent, 426 as oxidant for carbon monoxide, 426 preparation and properties, 114 structure, 114 lodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 lodine(III) reagents, see also hypervalent iodine reagents fluorosa, 402–404 lodine(III) species with I-N bonds, classification, 107, 10dine tribromide, 23 lodine (III), species with three carbon ligands, 107 lodine irribromide, 23 lodine (III), species with three carbon ligands, 107 lodine tribromide, 23 lodine (III), species with three carbon ligands, 107 lodine tribromide, 23 loxidative das chlorinating reagent, 158	polymer-supported, 389–391	chemical properties, 22
Iodine (VII) compounds, 8, 127, 128 Iodine (VII) fluoride, preparation, properties and structure, 127 Iodine fluorides, 22 Iodine (III) halides, 22, 23 Iodine (III) heterocycles aromatic character, 50 rive-membered 50–71 four-membered, 50, 74 six-membered 50, 72, 73 Istability, 50 Iodine (IVI) mixed oxide, 128 Iodine (IVII) mixed oxide, 128 Iodine (IVII) oxyfluorides, IO ₂ F ₃ and I ₄ O ₉ , 115 Iodine potal luorinating reagent, 303 as industrial fluorinating reagent, 426 as oxidativ for arbon monoxide, 426 preparation and properties, 114 Istructure, 114 Iodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 Iodine (III) reagents, see also hypervalent iodine reagents fluorous, 402–404 Iodine (III) reagents, see also hypervalent iodine reagents fluorous, 402–404 Iodine (III) species with I-N bonds, classification, 107, 108 Iodine (III) species with three carbon ligands, 107 Iodine irrichloride as chlorinating reagent, 158 Iodine (IVII) properties, 101 magnetic nanoparticle-supported, 401, 402 polymer-supported, 340 Iodoenzenes, ortho-substituted, as organocatalysts, 342 Iodobenzenes, ortho-substituted, as organocatalysts, 342 Iodobenzenesulfonamide, oxidation to I-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 3,3-dioxide, 69 2-Iodobenzenesulfonamide, oxidation of alkenes, 161, 162 Iodobenzenes, ortho-substituted, as organocatalysts, 342 3-diodoenzenes, ortho-substituted, as organocatalysts, 342 Iodocarboxylation of alkenes, 161, 162 Iodobenzenes, ortho-substituted, as organocatalysts, 342 2-Iodobenzenesulfonamide, oxidation to I-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 3,3-dioxide, 69 2-Iodobenzenesulfonamide, oxidation of alkenes, 162 2-Iodobenzenesulfonamide, oxidation of alkenes, 162 Iodobenzenesulfonamide, oxidation of alkenes, 162 Iodocarboxylation of alkenes, 162 Iodocarboxylation of alkenes, 162 Iodocarboxylation of alken	reactions in water, 416, 417	computational studies, 22
lodine(VII) fluoride, preparation, properties and structure, 127 lodine fluorides, 22 lodine(III) halides, 22, 23 lodine(III) heterocycles aromatic character, 50 five-membered 50-71 four-membered, 50, 74 six-membered 50, 72, 73 stability, 50 lodine (III) oxides, 120, and 14, 09, 115 lodine (IIII) oxides, 120, and 14, 09, 115 lodine pentafluoride as fluorinating reagent, 303 as industrial fluorinating reagent, 426 as oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 lodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 lodine(IIII) reagents, see also hypervalent iodine reagents fluorous, 402-404 lodine(III) reagents, see also hypervalent iodine reagents fluorous, 402-404 lodine(III) species with I-N bonds, classification, 107, 108 lodine(IIII) species with three carbon ligands, 107 lodine tribromide, 23 lodine(IIII) species with three carbon ligands, 107 lodine tribromide, 23 lodine(IIII) properties, 158 lodine(IIII) species with three carbon ligands, 107 lodine tribromide, 23 lodine (IIII) yackies to fix a stability, 22 X-ray, 22 lodine(IIII) ris(carboxylates), preparation and structure, 15 lodobenzene as excellent leaving group, 14 magnetic nanoparticle-supported, 402 polystyrene-supported, ac catalyst, 406 lodobenzene as excellent leaving group, 14 magnetic nanoparticle-supported, 402 polystyrene-supported, as catalyst, 406 lodobenzene as excellent leaving group, 14 magnetic nanoparticle-supported, 402 polystyrene-supported, as catalyst, 406 lodobenzene as excellent leaving group, 14 magnetic nanoparticle-supported, 402 polystyrene-supported, as catalyst, 406 lodobenzenesulfonamae, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 3,3-dioxide, 69 2-lodobenzenesulfonamae, as pre-catalysts for oxidation of alcones, archo-substituted, as prevalent ionine reagent, 100 lodocarboxylation of	recyclable, nonpolymeric, 405, 406	preparation, 22
lodine(VII) fluoride, preparation, properties and structure, 127 lodine fluorides, 22 lodine(III) halides, 22, 23 lodine(III) heterocycles aromatic character, 50 five-membered, 50, 74 six-membered, 50, 74 six-life supported, 402 social six distinct leaving group, 14 magnetic nanoparticle-supported, 402 social six divides, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole six-diodencemes-sulfonamide, oxidation to l-acetoxy-1,2-dihydro-1,3,	Iodine(VII) compounds, 8, 127, 128	
structure, 127 lodine(III) halides, 22 lodine(III) halides, 22, 23 lodine(III) oxide considered 50, 74 six-membered, 50, 74 six-membered, 50, 74 six-membered 50, 72, 73 stability, 50 lodine (V/VII) mixed oxide, 128 lodine(III) oxide, 1-03, 23 lodine(III) oxide, 1-03, 23 lodine(IVII) oxyfluorides, IO ₂ F ₃ and IOF ₃ , 127 lodine-oxygen bond lengths, statistical analysis, 11 lodine pentalfluoride as oxidatin for carbon monoxide, 426 preparation and properties, 114 structure, 114 lodine pentalfluoride lodine pentalfundine of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 lodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 402 lodine pentalfundine decarboxylation of alkenes, 161, 162 lodobenzene sexcellent leaving group, 14 magnetic nanoparticle-supported, 402 lodobenzenes, ortho-substituted, as organocatalysts, 342 343 2-lodobenzenes-surflonamide, oxidation to loacetoxyl-12-dihydro-1, 3,2-benziodathiazole 3,3-dioxide, 69 2-lodobenzenes-surflonamide, oxidation to loacetoxyl-12-dihydro-1, 3,2-benziodathiazole 3,3-dioxide, 69 2-lodosenzenes-urbox for oxidation of alkenes, 162 lodobenzenes ortho-substituted, as organocatalysts, 342 343 2-lodobenzenes-surflonamide, oxidation to loacetoxyl-12-dihydro-1, 3,2-benziodathiazole 3,3-dioxide, 69 2-lodosenzenes-urbox for oxidation of alkenes, 162 lodosenzenes-urbox for oxidation to loacetoxyl-12-dihydro-1, 3,2-benziodathiazole 3,3-dioxide, 69 2-lodosenzenes-urbox for oxidation of alkenes, 162 lodocarboxylation of alkenes, 162 lodocarbox for oxidation to loacetoxy-1-2-dihydro-1, 3,2-benziodathiazole 3,3-dioxide, 69 2-lodosenzenes orb	Iodine(VII) fluoride, preparation, properties and	
lodine (III) halides, 22, 23 lodine (III) heterocycles aromatic character, 50 five-membered 50–71 four-membered, 50, 74 six-membered 50, 72, 73 six-membered 50, 72, 73 six-membered 50, 72, 73 lodine (IV/VII) mixed oxide, 128 lodine (III) oxide, 1 ₂ O ₃ , 23 lodine (V/VII) mixed oxide, 128 lodine (III) oxides, 1 ₂ O ₄ and 1 ₄ O ₉ , 115 lodine (VIII) mixed oxide, 128 lodine (III) oxides, 1 ₂ O ₄ and 1 ₄ O ₉ , 115 lodine (VIII) mixed oxide, 128 lodine (III) oxides, 1 ₂ O ₄ and 1 ₄ O ₉ , 115 lodine pentafluoride as fluorinating reagent, 303 as industrial fluorinating reagent, 426 as oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 lodine (III) ris(carboxylation of alkenes, 161, 162 lodobenzene as excellent leaving group, 14 magnetic nanoparticle-supported, 402 polystyrene-supported, as catalyst, 406 lodobenzenes, ortho-substituted, as organocatalysts, 342 2-lodobenzenesulfonamide, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 3,3-dioxide, 69 2-lodobenzenesulfonamide, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 3,3-dioxide, 69 2-lodobenzenes as excellent leaving group, 14 magnetic nanoparticle-supported, as catalyst, 406 lodobenzenes, ortho-substituted, as organocatalysts, 342 2-lodobenzenesulfonamide, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 3,3-dioxide, 69 2-lodobenzenes as precuresulfonamide, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole as duchariating reagent, 362 2-lodosophthalic acid, as precursor to recyclable reagent, 396, 397 lodolactonization hypervalent iodine-catalyzed, 344 of pentenoic acids, 205 lodonium imides, 4, 6. See also imidoiodanes lodonium inides, 4, 6. See also imidoiodanes lodonium inimides, 4, 6. See also imidoiodanes lod		-
lodine(III) halides, 22, 23 lodine(III) heterocycles aromatic character, 50 five-membered 50, 71 four-membered 50, 72, 73 stability, 50 lodine (V/VII) mixed oxide, 128 lodine(III/V) oxides, 1 ₂ 0 ₄ and 1 ₄ 0 ₉ , 115 lodine-cycles oxygen bond lengths, statistical analysis, 11 lodine pentafluoride as fluorinating reagent, 303 as industrial fluorinating reagent, 426 as oxidatin for carbon monoxide, 426 preparation and properties, 114 structure, 114 lodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 lodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polystyrene-supported, 381 as a double fleaving group, 14 magnetic nanoparticle, supported, 402 polystyrene-supported, 400, 401 magnetic nanoparticle, supported, 401, 402 polystyrene-supported, 401, 402 polystyrene-supported, 341, 406 lodobenzene, ortho-substituted, as organocatalysts, 342 343 2-lodobenzenesulfonamide, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 3,3-dioxide, 69 2-lodobenzenesulfonamide, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 3,3-dioxide, 69 2-lodobenzenes ulfronamide, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole as fluorious, 402 2-lodoisophthalic acid, as precursor to recyclable reagent, 396, 397 lodonium imides, 4, 6, See also imidoiodanes lodonium indes, 4, 6, See also imidoiodanes lodonium salts lodonium salt, 74, 75 lodonium salts lodonium salt, 74, 75 lodonium salts lodonium salts lodonium salts lodonium salt, 74, 75 lodonium salts lodonium	Iodine fluorides, 22	
lodine(III) heterocycles aromatic character, 50 five-membered 50-71 four-membered, 50, 74 six-membered 50, 72, 73 stability, 50 lodine (V/VII) mixed oxide, 128 lodine(III) oxide, 1 ₂ 0, 23 lodine (V/VII) mixed oxide, 128 lodine(III) oxyfluorides, 1 ₂ 0, and 1 ₄ O ₉ , 115 lodine-oxygen bond lengths, statistical analysis, 11 lodine pentafluoride as fluorinating reagent, 303 as industrial fluorinating reagent, 426 as oxidant for carbon monoxide, 426 preparation and properties, 114 structure, 114 lodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 lodine (III) reagents, see also hypervalent iodine reagents fluorous, 402–404 loinic-liquid-supported, 401, 402 polymer-supported, 3,24 and IoFs, 127 lodine tribromide, 23 lodine tribromide, 23 lodine trichloride as chlorinating reagent, 158 lodosarchoxylation of alkenes, 162 lodobenzene as excellent leaving group, 14 magnetic nanoparticle-supported, 402 polystyrene-supported, as catalyst, 406 lodobenzenes, ortho-substituted, as organocatalysts, 342 3-lodobenzenesulfonamide, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 3,3-dioxide, 69 2-lodobenzenesulfonamide, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 3,3-dioxide, 69 2-lodobenzenesulfonamide, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 3,3-dioxide, 69 2-lodobenzenesulfonamide, oxidation to lodocarboxylation of alkenes, 162 2-lodobenzenes, ortho-substituted, as organocatalysts, 342 3-lodobenzenes, ortho-substituted, as organocatalysts, 342 2-lodobenzenesulfonamide, oxidation to lodocarboxylation of alcenos, ortho-substituted, as organocatalysts, 342 2-lodobenzenesulfonamide, oxidation to lodocarboxylation of alcenos, ortho-substituted, as organocatalysts, 342 2-lodobenzenesulfonamide, oxidation to lodocarboxylation of alcenos, s69 2-lodobenzenesulfonamide, oxidation of alcenos, s69 2-lodobenzen		
aromatic character, 50 five-membered 50, 74 six-membered 50, 72, 73 stability, 50 Iodine (VIVII) mixed oxide, 128 Iodine(III) oxide, 1 ₂ O ₃ , 23 Iodine(III) oxide, 1 ₂ O ₃ , 23 Iodine(III) oxide, 1 ₂ O ₃ , 23 Iodine-oxygen bond lengths, statistical analysis, 11 Iodine pentafluoride as fluorinating reagent, 303 as industrial fluorinating reagent, 426 as oxidant for carbon monoxide, 426 preparation and properties, 114 Istructure, 114 Iodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarrboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 Iodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Iodine(III), species with 1-N bonds, classification, 107, 108 Iodine tribromide, 23 Iodine trichloride as chlorinating reagent, 158 Iodonium in an propertic leaving group, 14 magnetic nanoparticle-supported, 402 polystrene-supported, as catalyst, 406 Iodobenzenes, ortho-substituted, as organocatalysts, 342 Iodobenzenesulfonamide, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole a,3-dioxide, 69 2-lodobenzenesulfonamide, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole as qualtical analysis, 11 lodaceobenzenes, ortho-substituted, as organocatalysts, 342 Iodobenzenesulfonamide, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole as chlorinating reagent, 426 a,3-dioxide, 69 2-lodobenzenesulfonamide, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole as chlorinating reagent, 426 and sidney decarboxylation of alkenes, 162 2-lodoisophthalic acid, as precursor to recyclable reagent, 396, 397 Iodolactonization properties, 101 properties, 101 rearrangement by 1,4 aryl migration, 101 Iodolnium salt, 74, 75 Iodonium salts biological activity, 439–443 cyclic, 74–76 key structural features, 8 ligand coupling mechanism,		Iodoacetoxylation of alkenes, 161, 162
five-membered 50-71 four-membered, 50, 74 six-membered 50, 72, 73 stability, 50 Iodine (WVII) mixed oxide, 128 Iodine(III) oxide, 1 ₂ O ₃ , 23 Iodine(III) oxide, 1 ₂ O ₃ , 23 Iodine(VIVI) oxyfluorides, IO ₂ F ₃ and IOF ₅ , 127 Iodine-oxygen bond lengths, statistical analysis, 11 Iodine pentafluoride as fluorinating reagent, 303 as industrial fluorinating reagent, 426 as oxidant for carbon monoxide, 426 preparation and properties, 114 structure, 114 Iodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 Iodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, as catalyst, 406 Iodobenzenes, ortho-substituted, as organocatalysts, 342 343 2-lodobenzenesulfonamide, oxidation to l-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 3,3-dioxide, 69 2-lodobenzenesulfonamide, 36, 362 Iodocarboxylation of alkenes, 162 2-lodoisophthalic acid, as precursor to recyclable reagent, 396, 397 Iodoline-catalyzed, 344 of pentenoic acids, 205 Iodonium ion, IUPAC rules, 4 Iodonium phenolates preparation from phenols and (diacetoxyiodo)benzene 101 properties, 101 rearrangement by 1,4 aryl migration, 101 Iodolium salt, 74, 75 Iodonium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439		
four-membered, 50, 74 six-membered 50, 72, 73 stability, 50 lodine (VVII) mixed oxide, 128 lodine(III) oxide, 1 ₂ O ₃ , 23 lodine(III) oxyfluorides, 1O ₂ F ₃ and 1OF ₅ , 127 lodine-oxygen bond lengths, statistical analysis, 11 lodine pentafluoride as fluorinating reagent, 303 as industrial fluorinating reagent, 426 as oxidant for carbon monoxide, 426 preparation and properties, 114 structure, 114 lodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 lodine (III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 lodine (III) species with I-N bonds, classification, 107, 108 lodine tribromide, 23 lodine trichloride as chlorinating reagent, 158 magnetic nanoparticle-supported, 402 polymer-supported, 401, 402 polymer-supported, 23 lodine trichloride as chlorinating reagent, 158 magnetic nanoparticle-supported, 402 polymer-supported, 401, 402 polymer-supported, 23 lodine trichloride as chlorinating reagent, 158 magnetic nanoparticle-supported, 402 polymer-supported, 401, 402 polymer-supported, 401	· · · · · · · · · · · · · · · · · · ·	
six-membered 50, 72, 73 stability, 50 Iodine (WVII) mixed oxide, 128 Iodine(III) oxide, 1 ₂ O ₃ , 23 Iodine(IIII) oxide, 1 ₂ O ₃ , 23 Iodine(IIII) oxides, 1 ₂ O ₄ , and 1 ₄ O ₉ , 115 Iodine (VVI) oxides, 1 ₂ O ₄ , and 1 ₄ O ₉ , 115 Iodine (VVI) oxythorides, 10 ₂ F ₃ and IOF ₅ , 127 Iodine oxygen bond lengths, statistical analysis, 11 Iodine pentafluoride as fluorinating reagent, 303 as industrial fluorinating reagent, 426 as oxidant for carbon monoxide, 426 preparation and properties, 114 structure, 114 structure, 114 dodine pentoxide oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 sa reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 Iodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Iodine (III) species with I–N bonds, classification, 107, 108 Iodine (III), species with three carbon ligands, 107 Iodine tribromide, 23 Iodobenzenes, ortho-substituted, as organocatalysts, 342 Iodobenzenesulfonamide, oxidation to I-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 3,3-dioxide, 69 2-Iodobenzenesulfonamide, oxidation to I-acetoxy-1,2-dihydro-1,3,2-benziodathiazole 3,d-dioxide, 69 2-Iodobenzenesulfonamide, oxidation of alcentoxy, 16 2-Iodobenzenesulfonamide, oxidation of alcohols, 361, 362 Iodocarboxylation of alkenes, 162 2-Iodosarboxylation of alkenes, 162 2-Iodosarboxylation of placetoxylation of placetoxyla		
stability, 50 lodine (WVII) mixed oxide, 128 lodine(III) oxide, 1 ₂ O ₃ , 23 lodine(IIII) oxides, 1 ₂ O ₄ and I ₄ O ₉ , 115 lodine(VII) oxyfluorides, IO ₂ F ₃ and IOF ₅ , 127 lodine-oxygen bond lengths, statistical analysis, 11 lodine pentafluoride as fluorinating reagent, 303 as industrial fluorinating reagent, 426 preparation and properties, 114 structure, 114 lodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 lodine (III) reagents. see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 lodine(III) species with I-N bonds, classification, 107, 108 lodine (III), species with three carbon ligands, 107 lodine tribromide, 23 lodine trichloride as chlorinating reagent, 158		
lodine (V/VII) mixed oxide, 128 lodine(III) voide, I ₂ O ₃ , 23 lodine(III) voides, I ₂ O ₄ and I ₄ O ₉ , 115 lodine(III) oxyfluorides, IO ₂ F ₃ and IOF ₅ , 127 lodine-oxygen bond lengths, statistical analysis, 11 lodine pentafluoride as fluorinating reagent, 303 as industrial fluorinating reagent, 426 as oxidant for carbon monoxide, 426 preparation and properties, 114 structure, 114 lodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 lodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 lodine(III) species with I-N bonds, classification, 107, 108 lodine(III), species with three carbon ligands, 107 lodine tribromide, 23 lodine trichloride as chlorinating reagent, 158		
lodine(III) oxide, I ₂ O ₃ , 23 lodine(IIIV) oxides, I ₂ O ₄ and I ₄ O ₉ , 115 lodine(VII) oxyfluorides, IO ₂ F ₃ and IOF ₅ , 127 lodine-oxygen bond lengths, statistical analysis, 11 lodine pentafluoride as fluorinating reagent, 303 as industrial fluorinating reagent, 426 as oxidant for carbon monoxide, 426 preparation and properties, 114 structure, 114 lodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 lodine(III) reagents, see also hypervalent iodine reagents fluorious, 402–404 ionic-liquid-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 lodine(III) species with I–N bonds, classification, 107, 108 lodine tribromide, 23 lodine trichloride as chlorinating reagent, 158 2-Iodobenzenesulfonamide, oxidation to l-acctoxy-1,2-dihydro-1,3,2-benziodathiazole 3,3-dioxide, 69 2-Iodobenzenesulfonates, as pre-catalysts for oxidation of alcohols, 361, 362 lodocarboxylation of alkenes, 162 2-Iodoisophthalic acid, as precursor to recyclable reagent, 396, 397 lodolactonization hypervalent iodine-catalyzed, 344 of pentenoic acids, 205 lodonium inides, 46. See also imidoiodanes lodonium phenolates preparation from phenols and (diacetoxyiodo)benzene 101 properties, 101 rearrangement by 1,4 aryl migration, 101 lodolium salt, 74, 75 lodonium salt, 74, 75 lodoium salt, 74, 75 lodoium alts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structure, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439		
Iodine(III/V) oxides, I ₂ O ₄ and I ₄ O ₉ , 115 I-acetoxy-1,2-dihydro-1,3,2-benziodathiazole Iodine(VII) oxyfluorides, IO ₂ F ₃ and IOF ₅ , 127 Iodine-oxygen bond lengths, statistical analysis, 11 Iodine pentafluoride as fluorinating reagent, 303 as industrial fluorinating reagent, 426 as oxidant for carbon monoxide, 426 preparation and properties, 114 Iodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 Iodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Iodine(III) species with I–N bonds, classification, 107, 108 Iodine tribromide, 23 Iodine trichloride as chlorinating reagent, 158		
Iodine (VII) oxyfluorides, IO ₂ F ₃ and IOF ₅ , 127 Iodine—oxygen bond lengths, statistical analysis, 11 Iodine pentafluoride as fluorinating reagent, 303 as industrial fluorinating reagent, 426 as oxidant for carbon monoxide, 426 preparation and properties, 114 Iodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 Iodine (III) reagents, see also hypervalent iodine reagents fluorous, 402—404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381—389 recyclable, nonpolymeric, 391—404 Iodine (III) species with 1–N bonds, classification, 107, 108 Iodine tribromide, 23 Iodine trichloride as fluorious, 402—404 as chlorinating reagent, 158 3,3-dioxide, 69 2-Iodobenzenesulfonates, as pre-catalysts for oxidation of alkenes, 162 2-Iodoisophthalic acid, as precursor to recyclable reagent, 396, 397 Iodolactonization hypervalent iodine-catalyzed, 344 of pentenoic acids, 205 Iodonium imides, 4, 6. See also imidoiodanes Iodonium phenolates preparation from phenols and (diacetoxyiodo)benzene 101 properties, 101 rearrangement by 1,4 aryl migration, 101 Iodolium salt, 74, 75 Iodonium salts biological activity, 439—443 cyclic, 74—76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260—277 structure, 3 toxicity towards mammals, 439		
Iodine—oxygen bond lengths, statistical analysis, 11 Iodine pentafluoride as fluorinating reagent, 303 as industrial fluorinating reagent, 426 as oxidant for carbon monoxide, 426 preparation and properties, 114 Iodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 preparation and properties, 114 as reagent, 302, 303 Iodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Iodine(III) species with I-N bonds, classification, 107, 108 Iodine(III), species with three carbon ligands, 107 Iodine tribromide, 23 Iodine tribromide, 23 Iodine trichloride as fluorinating reagent, 303 as industrial fluorinating reagent, 426 Iodocarboxylation of alchoes, 361, 362 Iodocarboxylation of alchenes, 162 2-Iodoisenznesulfonates, as pre-catalysts for oxidation of alcohols, 361, 362 Iodocarboxylation of alchenes, 162 2-Iodoisenbhalic acid, as precursor to recyclable reagent, 396, 397 Iodolactonization hypervalent iodine-catalyzed, 344 of pentenoic acids, 205 Iodonium sides, 4, 6. See also imidoiodanes Iodonium phenolates preparation from phenols and (diacetoxyiodo)benzene 101 properties, 101 rearrangement by 1,4 aryl migration, 101 Iodolium salt, 74, 75 Iodonium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439		
lodine pentafluoride as fluorinating reagent, 303 as industrial fluorinating reagent, 426 as oxidant for carbon monoxide, 426 preparation and properties, 114 structure, 114 lodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 lodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 lodine(III) species with I-N bonds, classification, 107, 108 lodine(III), species with three carbon ligands, 107 lodine tribromide, 23 lodine tribromide, 23 lodine trichloride as fluorinating reagent, 303 lodioa oxidation of alkenes, 162 2-Iodoisophthalic acid, as precursor to recyclable reagent, 396, 397 lodolactonization hypervalent iodine-catalyzed, 344 of pentenoic acids, 205 lodonium imides, 4, 6. See also imidoiodanes lodonium phenolates preparation from phenols and (diacetoxyiodo)benzene 101 properties, 101 rearrangement by 1,4 aryl migration, 101 lodolium salt, 74, 75 lodonium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439		
as fluorinating reagent, 303 as industrial fluorinating reagent, 426 as oxidant for carbon monoxide, 426 preparation and properties, 114 Iodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 Iodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Iodine(III) species with I-N bonds, classification, 107, 108 Iodine (III), species with three carbon ligands, 107 Iodine tribromide, 23 Iodocarboxylation of alkenes, 162 2-Iodoisophthalic acid, as precursor to recyclable reagent, 396, 397 Iodolactonization hypervalent iodine-catalyzed, 344 of pentenoic acids, 205 Iodonium imides, 4, 6. See also imidoiodanes Iodonium phenolates preparation from phenols and (diacetoxyiodo)benzene 101 properties, 101 rearrangement by 1,4 aryl migration, 101 Iodolium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439		
as industrial fluorinating reagent, 426 as oxidant for carbon monoxide, 426 preparation and properties, 114 Iddine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 preparation and properties, 114 as reagent, 302, 303 preparation of ketones and aromatic compounds, 303 Iddine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Iodine(III) species with I–N bonds, classification, 107, 108 Iodine(III), species with three carbon ligands, 107 Iodine tribromide, 23 Iodine trichloride as chlorinating reagent, 426 reagent, 396, 397 Iodolactonization hypervalent iodine-catalyzed, 344 of pentenoic acids, 205 Iodonium indies, 4, 6. See also imidoiodanes Iodonium phenolates preparation from phenols and (diacetoxyiodo)benzene 101 rearrangement by 1,4 aryl migration, 101 Iodolium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 Iodonium influes, 4, 6. See also imidoiodanes Iodonium influes, 4, 6. See also Iodonium infl		
as oxidant for carbon monoxide, 426 preparation and properties, 114 Iodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 Iodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Iodine(III) species with I–N bonds, classification, 107, 108 Iodine(III), species with three carbon ligands, 107 Iodine trichloride as chlorinating reagent, 158 reagent, 396, 397 Iodolactonization hypervalent iodine-catalyzed, 344 of pentenoic acids, 205 Iodonium imides, 4, 6. See also imidoiodanes Iodonium phenolates preparation from phenols and (diacetoxyiodo)benzene 101 properties, 101 properties, 101 rearrangement by 1,4 aryl migration, 101 Iodolium salt, 74, 75 Iodonium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439		· · · · · · · · · · · · · · · · · · ·
preparation and properties, 114 structure, 114 lodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 lodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-suppotted, 381–389 recyclable, nonpolymeric, 391–404 lodine(III) species with I–N bonds, classification, 107, 108 lodine(III), species with three carbon ligands, 107 lodine tribromide, 23 lodine trichloride as chlorinating reagent, 158 lodonium imides, 4, 6. See also imidoiodanes lodonium inion, IUPAC rules, 4 lodonium inion, IUPAC rules, 4 lodonium phenolates preparation from phenols and (diacetoxyiodo)benzene 101 properties, 101 rearrangement by 1,4 aryl migration, 101 lodolium salt, 74, 75 lodonium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439		
structure, 114 Idodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 Idodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Idodine(III) species with I–N bonds, classification, 107, 108 Idodine(III), species with three carbon ligands, 107 Idodine tribromide, 23 Idodine trichloride as chlorinating reagent, 158 hypervalent iodine-catalyzed, 344 of pentenoic acids, 205 Idodnium imides, 4, 6. See also imidoiodanes Iodonium in, IUPAC rules, 4 Iodonium phenolates preparation from phenols and (diacetoxyiodo)benzene 101 properties, 101 properties, 101 Iodolium salt, 74, 75 Iodonium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439		
lodine pentoxide oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 Iodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Iodine(III) species with I–N bonds, classification, 107, 108 Iodine(III), species with three carbon ligands, 107 Iodine tribromide, 23 Iodine trichloride as chlorinating reagent, 158 of pentenoic acids, 205 Iodonium imides, 4, 6. See also imidoiodanes Iodonium phenolates preparation from phenols and (diacetoxyiodo)benzene 101 rearrangement by 1,4 aryl migration, 101 Iodolium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439		
oxidation of electron-rich alcohols in water, 303 oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 Iodonium imides, 4, 6. See also imidoiodanes Iodonium inides, 4, 6. See also inidoiodanes Iodonium inides, 4, 6		
oxidative decarboxylation of propiolic acids, 303 preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 Iodonium jhenolates preparation from phenols and (diacetoxyiodo)benzene 101 properties, 101 rearrangement by 1,4 aryl migration, 101 Iodolium salt, 74, 75 Iodonium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 Iodine trichloride as chlorinating reagent, 158 Iodonium ion, IUPAC rules, 4 Iodonium jhenolates preparation from phenols and (diacetoxyiodo)benzene 101 properties, 101 rearrangement by 1,4 aryl migration, 101 Iodolium salt, 74, 75 Iodonium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439	-	
preparation and properties, 114 as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 Iodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Iodine(III) species with I–N bonds, classification, 107, 108 Iodine(III), species with three carbon ligands, 107 Iodine tribromide, 23 Iodine trichloride as chlorinating reagent, 158 Iodonium phenolates preparation from phenols and (diacetoxyiodo)benzene 101 properties, 101 rearrangement by 1,4 aryl migration, 101 Iodolium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439		
as reagent, 302, 303 structure, 115 for thiocyanation of ketones and aromatic compounds, 303 Iodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Iodine(III) species with I–N bonds, classification, 107, 108 Iodine(III), species with three carbon ligands, 107 Iodine tribromide, 23 Iodine trichloride as chlorinating reagent, 158 preparation from phenols and (diacetoxyiodo)benzene 101 properties, 101 rearrangement by 1,4 aryl migration, 101 Iodolium salts, 74, 75 Iodonium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439		
structure, 115 for thiocyanation of ketones and aromatic compounds, 303 Iodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Iodine(III) species with I–N bonds, classification, 107, 108 Iodine(III), species with three carbon ligands, 107 Iodine tribromide, 23 Iodine trichloride as chlorinating reagent, 158 Iodolium salt, 74, 75 Iodolium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439		-
for thiocyanation of ketones and aromatic compounds, 303 Iodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Iodine(III) species with I–N bonds, classification, 107, 108 Iodine(III), species with three carbon ligands, 107 Iodine tribromide, 23 as chlorinating reagent, 158 properties, 101 rearrangement by 1,4 aryl migration, 101 Iodolium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439		preparation from phenols and (diacetoxyiodo)benzene
303 rearrangement by 1,4 aryl migration, 101 Iodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Iodine(III) species with I–N bonds, classification, 107, 108 Iodine(III), species with three carbon ligands, 107 Iodine tribromide, 23 Iodine trichloride as chlorinating reagent, 158 rearrangement by 1,4 aryl migration, 101 Iodolium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439		101
Iodine(III) reagents, see also hypervalent iodine reagents fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Iodine(III) species with I–N bonds, classification, 107, 108 Iodine(III), species with three carbon ligands, 107 Iodine tribromide, 23 Iodine trichloride as chlorinating reagent, 158 Iodolium salt, 74, 75 Iodonium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439	· · · · · · · · · · · · · · · · · · ·	
fluorous, 402–404 ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Iodine(III) species with I–N bonds, classification, 107, 108 Iodine(III), species with three carbon ligands, 107 Iodine tribromide, 23 a schlorinating reagent, 158 Iodonium salts biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439	303	rearrangement by 1,4 aryl migration, 101
ionic-liquid-supported, 400, 401 magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Iodine(III) species with I–N bonds, classification, 107, 108 Iodine(III), species with three carbon ligands, 107 Iodine tribromide, 23 reactions, 260–277 Iodine trichloride as chlorinating reagent, 158 biological activity, 439–443 cyclic, 74–76 definition and classification, 76 history, 76 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439	Iodine(III) reagents, see also hypervalent iodine reagents	Iodolium salt, 74, 75
magnetic nanoparticle-supported, 401, 402 polymer-supported, 381–389 recyclable, nonpolymeric, 391–404 Iddine(III) species with I–N bonds, classification, 107, 108 ligand coupling mechanism, 14 Iddine(III), species with three carbon ligands, 107 Iddine tribromide, 23 reactions, 260–277 Iddine trichloride as chlorinating reagent, 158 cyclic, 74–76 definition and classification, 76 history, 76 key structural features, 8 ligand coupling mechanism, 14 mechanism of photolysis, 427 reactions, 260–277 structure, 3 toxicity towards mammals, 439	fluorous, 402–404	Iodonium salts
polymer-supported, 381–389 definition and classification, 76 recyclable, nonpolymeric, 391–404 lodine(III) species with I–N bonds, classification, 107, 108 ligand coupling mechanism, 14 lodine(III), species with three carbon ligands, 107 mechanism of photolysis, 427 lodine tribromide, 23 reactions, 260–277 lodine trichloride as chlorinating reagent, 158 toxicity towards mammals, 439		biological activity, 439–443
recyclable, nonpolymeric, 391–404 Iodine(III) species with I–N bonds, classification, 107, 108 Iodine(III), species with three carbon ligands, 107 Iodine tribromide, 23 Iodine trichloride 108 Island coupling mechanism, 14 Indine trichloride 108 Indine trichloride	magnetic nanoparticle-supported, 401, 402	cyclic, 74–76
Iodine(III) species with I–N bonds, classification, 107, key structural features, 8 108 ligand coupling mechanism, 14 Iodine(III), species with three carbon ligands, 107 mechanism of photolysis, 427 Iodine tribromide, 23 reactions, 260–277 Iodine trichloride structure, 3 as chlorinating reagent, 158 toxicity towards mammals, 439	polymer-supported, 381–389	definition and classification, 76
108 ligand coupling mechanism, 14 Iodine(III), species with three carbon ligands, 107 mechanism of photolysis, 427 Iodine tribromide, 23 reactions, 260–277 Iodine trichloride structure, 3 as chlorinating reagent, 158 toxicity towards mammals, 439	recyclable, nonpolymeric, 391-404	history, 76
Iodine(III), species with three carbon ligands, 107 mechanism of photolysis, 427 Iodine tribromide, 23 reactions, 260–277 Iodine trichloride structure, 3 as chlorinating reagent, 158 toxicity towards mammals, 439	Iodine(III) species with I–N bonds, classification, 107,	key structural features, 8
Iodine(III), species with three carbon ligands, 107 mechanism of photolysis, 427 Iodine tribromide, 23 reactions, 260–277 Iodine trichloride structure, 3 as chlorinating reagent, 158 toxicity towards mammals, 439	108	ligand coupling mechanism, 14
Iodine tribromide, 23 reactions, 260–277 Iodine trichloride structure, 3 as chlorinating reagent, 158 toxicity towards mammals, 439	Iodine(III), species with three carbon ligands, 107	
Iodine trichloride structure, 3 as chlorinating reagent, 158 toxicity towards mammals, 439		
as chlorinating reagent, 158 toxicity towards mammals, 439	Iodine trichloride	
	as chlorinating reagent, 158	
preparation, 22 X-ray structural data, 76		•
properties, 22, 23 Iodonium 8-I-2 species, in gas phase, 13		

Iodonium transfer reaction, 81	2-Iodosylbenzoic acid, see
Iodonium ylides, 4, 6, 99–106	1-hydroxy-1,2-benziodoxole-3(1 <i>H</i>)-one
as carbene precursors, 278–282	3-Iodosylbenzoic acid
cyclic, 101	as oxidant, 284
1,3-dicarbonyl derivatives, 99, 100	preparation, 32
intramolecular phenyl migration, DFT study, 12	as recyclable reagent, 397–399
monocarbonyl, 105	Iodosyl compounds
preparation, 99–105	polymeric structure, 31
properties, 99	preparation, 31
reactions, 278–282	properties, 31
reactions, catalyzed by rhodium(II) or copper	Iodosyl elimination, in iodosylalkanes, 248, 249
complexes, 280, 281	Iodosyl fluorosulfate, OIOSO ₂ F, preparation and
transylidation reactions, 103	properties, 23
X-ray structural studies 105, 106	Iodosylperfluoroalkanes, preparation, by hydrolysis of
Iodoperfluoroalkanes, fluorous, 402	$C_nF_{2n+1}I(CO_2CF_3)_2$, 32
4-Iodophenoxyacetic acid, as catalyst, for oxidation of	2-Iodosylphenylacetic acid, cyclic tautomer, 72
phenols, 349	2-Iodosylphenylphosphoric acid, cyclic tautomer, 72, 73
Iodophosphoryloxylation of alkynes and alkenes, 162	Iodosyl selenate, (IO) ₂ SeO ₄ , 23
2-Iodopyridines, preparation and properties, 405 X-ray	Iodosyl selenate, (IO) ₂ SeO ₄ , 25 Iodosyl sulfate, (IO) ₂ SO ₄ , preparation and structure, 23
structure, 405 as recyclable reagents, 405	Iodosyl triflate, OIOTf, preparation and properties, 23
2-Iodosobenzoic acid (IBA), 5. See also	Iodotosyloxylation of alkynes, 161
1-hydroxy-1,2-benziodoxole-3(1 <i>H</i>)-one	Iodoxolones
Iodosylalkanes, as reactive intermediates, 248, 249	preparation, by addition of ICl ₃ to
Iodosylarenes	acetylenedicarboxylic acid, 70, 71
NMR and LC-MS study	preparation, by oxidative cyclization of iodofumaric
ortho substituted, solubility, 35	acid, 70, 71
ortho substituted, X-ray crystal structures, 35	X-ray structures, 70
preparation, by hydrolysis of acetates or chlorides, 32,	Iodoxybenzene, see iodylbenzene
33	2-Iodoxybenzenesulfonic acid. <i>See also</i>
preparation, by oxidation of iodoarenes, 31, 32	2-iodylbenzenesulfonic acid
as source of oxygen in transition metal-catalyzed	amides, preparation and X-ray structure, 118
oxidations, 251	preparation and oxidizing properties, 124
structural studies, 33, 34	2-Iodoxybenzoate, isopropyl
Iodosylbenzene	in the metalloporphyrin-catalyzed oxidations, 288
aqua complexes, 13	as reagent, 288
discovery, 2	2-Iodoxybenzoate, methyl ester, 117
disproportionation, 33	2-Iodoxybenzoic acid (IBX)
explosion, 33	amides (IBX-amides), preparation and X-ray structure
in metalloporphyrin-catalyzed oxidations, 250, 251	117
monomer complexes with 18C6 crown ether, 35	analogues of IBX, 122, 123
oligomeric, 33	applications in organic synthesis, 288–296
polymeric structure, 33, 34	as catalyst for oxidation of alcohols, 359, 360
polymer-supported, preparation, 386 reactions, 386	dearomatization of phenols, 292
preparation, by hydrolysis of PhI(OAc) ₂ , 32	esters (IBX-esters), preparation and X-ray structure,
properties, 33	117
reactions in water, 413–416	explosive properties, 121, 122
Iodosylbenzene, activated, cleavage of alkenes, 416, 417	fluorous IBX, 122
Iodosylbenzene, activation by bromide anion, 413, 414	generation of N-centered radicals, 294
Iodosylbenzene sulfate, polymer-supported, preparation	history, 2, 121
and reactions, 386	mechanism of alcohol oxidation, theoretical study, 12
Iodosylbenzene sulfates, 49	non-explosive formulation (SIBX), 122
· · · · · · · · · · · · · · · · · · ·	

2-Iodoxybenzoic acid (IBX) (Continued)	2-Iodylbenzenesulfonic acid (IBS)
o-methyl-substituted IBX (Me-IBX), 122	as catalyst benzylic oxidation, 363
oxidation of alcohols, 288–291	as catalyst for oxidation of alcohols, 361
oxidation of the benzylic carbon, 293, 294	cyclic or pseudocyclic derivatives, 4, 6
oxidation of 1,2-diols, 291	Iodyl fluorosulfate, 115
oxidation of secondary amines to imines, 294	2-Iodylphenol ethers
oxidations in ionic liquids, 419	preparation and X-ray structure, 120
oxidations in non-DMSO solvents, 291, 292	polymer-supported, 391
oxidative cyclization of anilides, 293, 294	2-Iodylphenol, tosyl derivative, preparation, 119
α-oxygenation of carbonyl compounds, 296	2-(o-Iodylphenyl)-oxazolines, chiral, preparation and
pK _a value, 122	properties, 120, 288
polymer-supported, preparation and reactions,	Iodyl sulfate, 115
390	Iodyl triflate, 115
preparation, 282, 283	Iodyltrifluoromethane, preparation, 120
preparation by bromate oxidation, 121	Ion-exchange, recycling methodology, 397
preparation using Oxone, 121	Ionic liquid, [emim] ⁺ [BF4] ⁻ , 400
properties, 283	Ionic liquids, as recyclable solvents, 418, 419
pyridinium salt, 123	Ionic-liquid-supported iodoarenes, as catalysts, 406
six-membered cyclic IBX analogues, 124	Ionic-liquid-supported reagents, 400, 401
soluble analogs (mIBX), 122	Ishihara's cycloetherification, enantioselective, 370, 371
structure, 283	Isoflavone derivatives, synthesis from
synthesis of amino sugars, 293	<i>o</i> -hydroxychalcones, 211
synthesis of heterocycles, 294, 295	Isoindolinone derivatives, synthesis, 203
synthesis of α , β -unsaturated carbonyl systems,	Isoxazoles
293	phenanthro-fused, by oxidative coupling, 199
tetrafluoro-IBX derivative (FIBX), 122	synthesis, 205
water-soluble analogs, 417, 418	Isoxazoline N-oxides, synthesis by oxidation of
X-ray structure, 121, 122	β-hydroxyketoximes, 217, 218
Iodylalkanes, matrix isolation and FTIR spectra,	Isoxazolines, synthesis from aldoximes and olefins, 217
120	·
2-Iodylaniline, tosyl derivative, preparation and X-ray structure, 119	(-)-Jesterone, enantioselective synthesis, 184
Iodylarenes, 115–120	Ketones, macrocyclic, synthesis by alkoxy radical
five-membered heterocyclic benziodoxole derivatives,	fragmentation, 240, 241
4, 6	Ketoximes, deoximation to ketones by HTIB, 212
noncyclic, 4, 6, 115, 116, 283–287	Kita's catalytic spirocyclization, 350
ortho-phosphoryl substituted, X-ray structure, 116	Koser's reagent, 4, 5, 169. See also
preparation and properties, 116	[hydroxy(tosyloxy)iodo]benzene (HTIB)
pseudocyclic, 4, 6, 116–120, 287, 288	Kryptofix (K2.2.2), 433, 434
Iodylbenzene	111) promit (121212), 100, 101
asymmetric oxidation of sulfides, 286, 287	β-Lactam antibiotics, synthesis, 298
cooxidant with 2-pyridineseleninic anhydride, 284	Lactams, synthesis, 204
explosive properties, 116	Lactols, synthesis by oxidative rearrangement reaction of
history, 116	2,3-epoxy alcohols, 208
oxidation of alcohols, 283	Lactones, synthesis, by radical cyclization of carboxylic
oxidation of naphthols, 284	acids, 239
preparation and properties, 116	by alkoxy radical fragmentation, 240
in the presence of VO(acac) ₂ , 286	Lactonization, of 4-phenyl-4-pentenoic acid, 205
X-ray structure, 116	Lagunamycin, synthesis, 291
4-Iodylbenzenesulfonate, potassium salt, preparation and	Lagunamyem, synthesis, 291 Lambda nomenclature, 3, 4
properties, 405, 406	Lamoda nomenciature, 3, 4 L-6-[¹⁸ F]fluoroDOPA, 435
properties, 105, 100	Lol Illinoiopoiri, 155

Ligand coupling, on the iodine atom, 13, 14

Ligand exchange

dissociative and associative mechanism, 13, 14 on the iodine atom, 13, 14

Lycorane Amaryllidaceae alkaloids, enantioselective synthesis of the core, 225

Macrocyclic iodonium salts, 76

Magnetite nanoparticles, 402

- (±)-Magnofargesin, synthesis via alkynyliodonium salt, 272
- (±)-Makaluvamine F, synthesis, 227
- (±)-Maoecrystal V, synthesis, 298
- (+)-Maritidine, synthesis by phenolic spirocyclization, 189
- *m*-chloroperoxybenzoic acid (*m*CPBA), as terminal oxidant in catalytic reactions, 337–356, 366, 369–371, 373, 374
- [Menthyloxy(tosyloxy)iodo]benzenes, chiral, as reagents for enantioselective oxidation of sulfides, 220, 221
- Metalloporphyrin-mediated oxygenations, in the presence of iodobenzene, 366, 367

Metalloporphyrins

adducts with iodosylbenzene, 252

as catalysts in oxidations with iodosylbenzene, 251

- (-)-Metazocine, asymmetric synthesis, 256
- 5-Methoxy[2.2]paracyclophan-4-yl iodonium salts, 439
- 2-Methoxyphenyliodonium

bis(methoxycarbonyl)methanide, X-ray structure, 106

[Methoxy(tosyloxy)iodo]benzene, PhI(OMe)OTs preparation, 45

reagent for oxidative rearrangements, 212

2-Methylphenyl(2-methoxyphenyl)iodonium chloride, conformational structure, 439

mGluR5 ligands, fluorine-18 labeled, 438

m-iodosylbenzoic acid, as source of oxygen in transition metal-catalyzed oxidations, 251, 252

Moenomycin A, synthesis, 291

Monocarbonyl iodonium ylides, generation *in situ* and reactions, 281

Monoperoxysulfate, tetraphenylphosphonium, as terminal oxidant, 359, 360

Morphinandienone alkaloids, synthesis, 200

N-acylnitrenium intermediates, generation from amide and hypervalent iodine, 203

N-alkylsaccharins, preparation by N-radical cyclization, 247

n-butylpyridinium tetrafluoroborate, [BPy]BF₄, as a recyclable solvent, 419

N-hydroxysuccinimide esters, preparation from alcohols, 292

N-(2-iodylphenyl)acylamides (NIPA)

polymer-supported, 390, 391

preparation and X-ray structure, 118, 119

 (\pm) -Nitidanin, synthesis, 291

Nitriles, from aldehydes and (diacetoxyiodo)benzene, in aqueous ammonia, 417

2-Nitrobenzo[b]furans, synthesis, 205

Nitrogen heterocycles, preparation by N-radical cyclization, 247

N,*N*-dimethylarylamines, azidation to *N*-azidomethyl derivatives, 226

N,N-dimethylhydrazides, cleavage to carboxylic acids, 212

Norsesquiterpene, spirolactone/testosterone hybrid, synthesis, 209

N-tosyliminoiodanes, ArINTs

as nitrene precursors in transition metal catalyzed reactions, 253

as reagents in transition metal catalyzed amidation, 255 *N*-tosyliminophenyliodane, PhINTs, preparation, 111

Oligomeric iodosylbenzene perchlorate, X-ray crystal structure, 34

Oligomeric iodosylbenzene sulfate, (PhIO)₃SO₃ ESI-MS

reactions in water, 416

as reagent for oxidation of sulfides, 218

as terminal oxidant in transition metal-catalyzed oxidations, 252

X-ray crystal structures, 34

Organoiodine(III) chlorides

preparation by chlorination of organic iodides, 27 structural studies, 30, 31

Organoiodine(III) compounds, classification by type of ligands, 4, 6

Organoiodine(V) compounds, common classes, 4, 6

Organoiodine(V) compounds, as protein tyrosine phosphatase (PTP) inhibitors, 439

Organoiodine(III) derivatives of phosphoric acid, 49

Organoiodine(III) difluorides, 23–27 structural studies, 26, 27

Organoiodine(V) fluorides, 126, 127

Organoiodine(III) nitrate, PhI(ONO₂)₂, 48

Organoiodine(III) perchlorate, PhI(OClO₃)₂, 48, 49

Organoiodine(III) triflate, PhI(OTf)2, 48

1-Organosulfonyloxy-1,2-benziodoxole-3(1*H*)-ones, preparation, 54

Ortho-effect, in reactions of diaryliodonium salts with nucleophiles, 265

Ortho-phenylation, of phenols, 262, 263	in intramolecular mode, 197–201
1,3,4-Oxadiazoles, 2,5-disubstituted, synthesis, 206	of ketones with carboxylic acids, 369
1,2,4-Oxadiazoles, synthesis, 204	of Michael adducts, by PhIO/Bu ₄ NI in water, 415
Oxazole derivatives, synthesis, 372, 375	mechanism 196, 197
Oxazolines, 2-substituted, synthesis, 206	of non-phenolic aromatic substrates, 200, 202
Oxidation of alcohols	Oxidative cyclizations, 201
by 1-chloro-1,2-benziodoxol-3(1H)-one/TEMPO, 167	Oxidative cycloetherification, of oxo-substituted phenols
1,2-diols, 167	with Bu_4NI/H_2O_2 , 370
with DMP, 296–299	enantioselective, 370
epimerization-free, 298	Oxidative dearomatization
(fluoroalkyl)alkanols, to aldehydes, 166	of anilines, 195, 196
by HTIB under solvent-free microwave irradiation, 167	catalytic, 350
with IBX, 288–291	formation of cyclohexadienones, 195, 196
by iodine(III) reagents, 164–168	in the intramolecular mode, 187
by iodosylbenzene/KBr, 164, 165	mechanism, 196
with Oxone, catalyzed by PhI and RuCl ₃ , 367, 368	of phenols, 183–195
by PhICl ₂ /TEMPO, 167	in the presence of carbon nucleophiles, 187
by PhI(OAc) ₂ /I ₂ , 165	in the presence of fluoride anion, 187
by PhI(OAc) ₂ /NaBr, 165	of 2-substituted phenols, 193–195
by PhI(OAc) ₂ /TEMPO, 165–167	using (diacetoxyiodo)benzene in methanol, 184, 185
with potassium peroxodisulfate, catalyzed by PhI and	Oxidative decarboxylation, of 2-aryl-substituted
TEMPO, 366	carboxylic acids, 206
in the presence of Wittig ylide, 166, 291, 298	Oxidative decarboxylation/iodination of carboxylic acids,
transition metal catalyzed, 167	162
Oxidation of alkylbenzenes, with Oxone, catalyzed by	Oxidative fragmentations, 201, 206
PhI and RuCl ₃ , 367, 368	Oxidative functionalization, hypervalent
Oxidation of anilines, hypervalent iodine-catalyzed, 357	iodine-catalyzed, 342–346
Oxidation of iodoarene, to aryliodine(III) in situ, 337, 338	Oxidative iodination
Oxidations	of alkanes, with the PhI(OAc) ₂ -iodine system, 160
of antimony, 222	of aromatic and heteroaromatic compounds, 160, 161
of bismuth, 222	of electron-deficient heterocyclic systems, 160
of C–H bonds, at the benzylic or allylic position, 181,	of ketones, 162
182	of terminal alkynes, 163
of nitrogen, 216–218	using tosyloxybenziodoxole-iodine system, 161
with PhIO/Bu ₄ NI in water, 415, 416	Oxidative phenolic cyclizations, in the synthesis of
of sulfur, 218	acetylaspidoalbidine, 192
transition metal catalyzed, 250	(–)-cylindricine C and (–)-2-epicylindricine C, 192
using catalytic Bu ₄ NI and TBHP as oxidant, 369	dihydrofuranobenzofurans, 192
Oxidative amination, of arenes, hypervalent	discorhabdin A, 192
iodine-catalyzed, 347, 349	galanthamine, 192
Oxidative bromination, of arenes, hypervalent	platensimycin, 192
iodine-catalyzed, 346	(+)-plicamine, 192
Oxidative brominations, using iodine(III) reagent and	tuberostemonine and didehydrotuberostemonine, 192
bromide, 158, 159	Oxidative rearrangements, 201, 206
Oxidative cleavage, of alkenes and alkynes, hypervalent	Oxidative spirocyclization
iodine-catalyzed, 345	of amides and ketoximes, 187
Oxidative coupling	of catechol, 187
of aromatic substrates, 196–201	enantioselective, 353
of Bodipy monomers, 201, 203	in enantiospecific synthesis of (+)-puupehenone, 187
of cycloalkenylsilanes with silylated nucleobase, 200,	hypervalent iodine-catalyzed, 350–354
203	of N-protected tyrosine, 187

of phenolic substrates, 187 in synthesis of annosqualine, 190 in synthesis of cortistatin class of natural products,	Perfluoroalkyl(phenyl)iodonium triflates (FITS reagents) as electrophilic perfluoroalkylating reagents, 275, 276 preparation, 97
188	Periodate oxidation, application in chemical and
in synthesis of furoquinolinone and angelicin	instrumental analysis, 426
derivatives, 188	Periodates, 128
in synthesis of nitrogen heterocycles, 188	Periodate salts, as oxidizers in pyrotechnic formulations,
in synthesis of (\pm) -stepharine, 190	426
10-Oxidophenothiiodonium salt, 74, 75	Periodic acid
Oximes, oxidation to nitroso compounds, 218	glycol-cleavage, 303, 304
β-Oxoalkyl(phenyl)iodonium salts, generation from silyl	as oxidant, 426
enol ethers, 96, 97, 275	oxidative cleavage of protecting groups, 305
μ-Oxo-bridged iodanes	oxidative iodination of aromatic compounds, 305
mesylate, 47	properties and structure, 128
nitrate, 49	as reagent, 303–305
perchlorate, 48	synthesis of nitriles from alcohols or aldehydes, 305
tetrafluoroborate, 48	transition metal catalyzed oxidations, 304, 305
triflate, 48	Periodinanes, general name, 3
trifluoroacetate, 49, 192, 417	4,5-Phenanthryleneiodonium salts, 74, 75
(9-Oxo-9 <i>H</i> -fluoren-2-yl)phenyliodonium	Phenol ethers
hexafluoroantimonate, as photoinitiator, 42	intramolecular cyclization of azido derivatives 199,
Oxoindoles, 3,3-disubstituted, synthesis, 343	200
(±)-Oxomaritidine, synthesis, 385	nucleophilic substitution using hypervalent iodine,
Oxone (2KHSO ₅ ·KHSO ₄ ·K ₂ SO ₄), in oxidation of	197
iodobenzene, 32	oxidative coupling, hypervalent iodine-catalyzed, 354
Oxone, as terminal oxidant in catalytic reactions, 337,	Phenols
340–343, 349–351, 355–368	oxidation to benzoquinones, hypervalent
α-Oxyacylation, Bu ₄ NI-catalyzed, 369	iodine-catalyzed, 349
Oxygen heterocycles, synthesis by oxidative cyclizations,	oxidation to o-quinones, by IBX, 292
205	oxidation to o-quinones, iodine(V) catalyzed, 364
Oxylactonization, enantioselective	<i>p</i> -substituted, oxidation to <i>p</i> -quinols, 350
with Bu ₄ NI as pre-catalyst and H ₂ O ₂ as oxidant, 369	Phenoxiodonium salt, 74, 75
of ketocarboxylic acids, hypervalent iodine-catalyzed,	Phenylbenziodazole, preparation and X-ray structure, 63
339	1-Phenyl-1,2-benziodoxole-3(1 <i>H</i>)-one, preparation,
of ortho-alkenylbenzoates, 179	59–60
	Phenyl(cyano)iodonium triflate, preparation and
Pachastrissamine, synthesis, 291	application as the iodonium transfer reagent, 95
p-Alkoxyphenols, oxidative dearomatization, by	Phenyl(3-formylphenyl)iodonium salts, synthesis and
(diacetoxyiodo)benzene, 184, 185	¹⁸ F-fluoridation, 438, 439
Palladacycles, reactions with aryliodonium imides, 257	Phenylhydrazones, regeneration of carbonyl, 218
(+)-Pancratistatin, total synthesis, 225	Phenyliodonium imide
Passerini reaction, 295	ortho-methoxy substituted, X-ray structure, 113
Pd-catalyzed oxidations, with hypervalent iodine	ortho-sulfonyl substituted, X-ray structure, 113
reagents, 252	polymeric structure, 113, 114
(+)-Peloruside A, synthesis, 298	Phenyliodonium, leaving group ability, 267
Pentafluoroethylbenziodoxolone, 61	Phenyliodonium ylides, X-ray structures
Pentafluorophenyliododifluoride, C ₆ F ₅ IF ₂ , X-ray and	Phenyl[2-(trimethylsilyl)phenyl]iodonium triflate, as
spectroscopy studies, 27	benzyne precursor, 266
Pentavalent iodine, structural types, 3, 4	Phosphonium-iodonium ylides
Perfluoroalkyliodine(V) fluorides, preparation and X-ray	preparation and X-ray structure, 104
structure, 126	reactions, 282

Phosphoranyl-derived benziodoxoles, preparation and X-ray structure, 60	Positron emission tomography (PET), 431, 432 Potassium iodate
Phosphoryloxylactonization, hypervalent iodine-catalyzed, 344	as dietary supplement and a food additive, 425 as source of iodine in iodized salt, 425
α-Phosphoryloxylation of ketones,	toxicity, 425, 426
iodobenzene-catalyzed, 342	Prins–pinacol, oxidative tandem process, 190, 191
Photoacid generators (PAGs), 427	Pseudocyclic iodylarenes, polymer-supported, 390, 391
Photoinitiators, for cationic photopolymerizations,	(–)-Pseudolaric acid B, synthesis, 298
426	Pseudopterosin and elisabethin, analogs, synthesis, 301
Photopolymerization, 427	Pseudorotation on the iodine atom, 9, 10, 14
Photoresist technology, 426	(±)-Pterocarpans, synthesis, 291
Photosensitivity, of iodonium salt, 427	Pyridine derivatives, synthesis, 375
Physostigmine, synthesis, 344	2-Pyridyldiselenide, as reagent, 285
(–)-Pinidine, asymmetric synthesis, 207	Pyrimidines, phenanthro-fused, by oxidative coupling,
(±)-Platensimycin, synthesis, 298	199
(–)-Platensimycin, synthesis by oxidative Prins–pinacol	Pyrroles, polysubstituted, synthesis, 206
tandem process, 191	Pyrrolidines, 2-substituted, synthesis, 239
<i>p</i> -nitro(dichloroiodo)benzene, chlorination of	Pyrrolidino[60]fullerene, synthesis, 204
pyrrolidine, 156, 157	Pyrrolidinone derivatives, synthesis, 203
Poly[(diacetoxyiodo)styrene]	Pyrrolin-4-ones, synthesis, 204
applications, 383–386	Pyrrolo(benzo)diazepines, synthesis, 204
oxidation of alcohols, 384	Pyrrolo[2,1-c][1,4]diazepines, synthesis, 203
oxidation of glycols, 384	Pyrroloiminoquinone system, by intramolecular coupling
oxidation of phenols, 384, 385	of azido derivatives, 199
oxidation of sulfides, 384	
oxidative iodination, 384	Quaternary ammonium iodide catalyst, chiral, 370
preparation, 382, 383	Quaternary carbon center, by oxidative
synthesis of heterocycles, 385	<i>ipso</i> -rearrangement, 191, 192
synthesis of organyltellurophosphates, 386	Quinazolines, synthesis, 375
Poly[(dichloroiodo)styrene]	Quinolinone derivatives, synthesis, 203
as chlorinating reagent, 381, 382	Q , 200
preparation, 381	Radical alkylation
Poly(ethylene glycol-400), PEG-400, as a recyclable	of alkenes, 237, 238
solvent, 419	of nitrogen heterocycles, 237, 238
Poly[hydroxy(tosyloxy)iodo]styrene, synthetic	Radical cyclizations, 236, 239, 247
applications, 387	Radical fragmentation–phosphorylation, 246
Polymerization, dual photo- and thermal-initiation	Radical fragmentations, 236–248
systems, 427, 428	Radical reactions of λ^3 -iodanes, 14, 15
Polymer-supported reagents, general application, 381	Radical rearrangements, 236, 239–248 Radicals
Polystyrene, azidation and click-type functionalization,	generation from amides and iodosylbenzene, 247, 248
228, 431	nitrogen-centered, generation, 247, 248
Polyvalent iodine, see hypervalent iodine	oxygen-centered, generation, 238–244
Polyvalent iodine compounds, see also hypervalent	Radiofluorination, of diaryliodonium salts, 432
iodine compounds	Reactions in water, 413–418
classification, 3, 4	Recyclable nonpolymeric reagents, separation
commercial availability, 425	techniques, 391
common abbreviations, 4, 5	Recyclable organic solvents, 418–420
common names, 4, 5	Recyclable reagents
IUPAC nomenclature, 3, 4	biphenyl, 394–396
practical applications, 425	derivatives of 1,3,5,7-tetraphenyladamantane, 394–396

terphenyl, 394–396	Spirofurans and benzopyrans, iodobenzene-catalyzed
tetraphenylmethane, 394–396	synthesis, 352
Reductive elimination of PhI, 13, 14	Spiroketalization of phenolic alcohols, DFT calculations,
Resolvin D1, synthesis, 299	12
Resveratrol, diacyl derivatives, hydroxylation, 296	Spirolactams, synthesis
Ring expansion	by catalytic spirocyclization, 353
in reactions of HTIB, with alkenes, 212	by oxidative spirocyclization, 188
reactions, promoted by hypervalent iodine, 207	Spirolactones, preparation from phenolic substrates,
in reactions of 1-vinylcycloalkanol derivatives, 212	350
Ripostatin B, synthesis, 299	Spirooxindoles, synthesis, 205, 355
(+)-Ritterazine K, synthesis, 298	(–)-Spirotryprostatin B, synthesis, 298
, ,	Steroidal derivatives, synthesis by alkoxy radical
Salvinorin A, synthesis, 298	fragmentation, 240, 241, 244
Sceletenone, total synthesis, 191	Steroidal substrates, oxidative transformations, 209, 210
Secondary bonding in hypervalent iodine compounds, 9,	(–)-Subincanadines A and B, synthesis, 291
11	Sulfate esters, cyclic, preparation, 176, 177
Secondary bonds, in [bis(acyloxy)iodo]arenes, 43	Sulfides
Seco-psymberin/irciniastatin A, synthesis, 206	α-azidation, 227
Selenodecarboxylation, 236	enantioselective oxidation, 220
Self-assembly of hypervalent iodine compounds, 9, 11	oxidation to sulfonyl chlorides, 219
SET mechanism, in reactions of λ^3 -iodanes, 15	oxidation to sulfoxides, 218
SET mechanism, in reactions of λ -iodanes, 15	Sulfonylation, allylic, of α-methyl styrenes, 373, 374
(+)-Sieboldine A, synthesis, 299	Sulfonyloxylactonization, hypervalent iodine-catalyzed,
	344
Silica-supported iodoarenes, as catalysts, 406 Silylalkenes, reaction with PhIO•BF ₃ , 175	
	Sulfoxides, chiral, by enantioselective oxidation of
Silyl enol ethers, oxidative functionalization	sulfides, 220, 221
preparation of 1,4-butanediones, 172	T 11 1 1 1 207
preparation of β-keto phosphonates, 172	Tagged hypervalent iodine reagents, 397
preparation of α -methoxylated esters and lactones, 173	Tandem catalytic systems, based on hypervalent iodine,
preparation of oxygen heterocycles, 172	364–368
preparation of α -substituted ketones, 171, 172	TEMPO
preparation of α -trifluoromethanesulfonyloxyketones,	ion-supported, 400
172	as radical scavenger, in reactions of diaryliodonium
α -tosylamination, 231, 232	salts, 434, 435
Single-electron transfer (SET) mechanism, 15	TEMPO-catalyzed oxidations, 166
SiO ₂ -supported RuCl ₃ , as recyclable catalyst, 393	4-Tert-butyl(difluoroiodo)benzene, fluorination of esters
(S)-Ketoprofen, synthesis, 266	of cephalosporin, 146, 148
Sodium metaperiodate, NaIO ₄	Tert-butyl hydroperoxide (TBHP), as terminal oxidant,
for azidoiodination of alkenes, 306	369, 371–375
for carbonylation of benzylamines, 306	1-(<i>Tert</i> -butylperoxy)benziodoxoles, preparation 55 X-ray
for diazidation of alkenes, 306	structure, 55
for glycol-cleavage, 305	1-(Tert-butylsulfonyl)-2-iodylbenzene, preparation and
for oxidation of alcohols to carbonyl compounds, 305	X-ray structure, 116
as reagent, 305–307	Tertiary allylic alcohols, rearrangement to enones,
for transition metal catalyzed oxidations, 306, 307	iodine(V) catalyzed, 362
Solvent-free reactions, 420–422	Tetrachloroiodate salts, as chlorinating reagents, 158
Somamide A, synthesis, 291	(Tetrafluoroiodo)benzene, C ₆ H ₅ IF ₄ , preparation,
Spiculoic acid A, synthesis, 291	properties and X-ray structure, 126
Spirastrellolide A, synthesis, 299	Tetrahydroisoquinolines
Spirocyclization, of 2-substituted phenols,	C-H functionalization, 377
enantioselective, 193, 194	synthesis by Ugi-type multicomponent reaction, 295

Tetrahydropyrans, 2-(<i>N</i> -acylaminal) substituted, synthesis, 206	Trifluoromethylation asymmetric, in the presence of imidazolidinone
(-)-Tetrodotoxin, synthesis using PhIO ₂ /Py ₂ Se ₂ ,	catalyst, 277
285	of C-nucleophiles, 277
1,2,4-Thiadiazoles, synthesis, 204	of β-ketoesters, 276, 277
Thiazoles	of N-nucleophiles, 277
2,4-disubstituted, synthesis from alkynyliodonium	of P-nucleophiles, 277
salts 272	of S-nucleophiles, 277
phenanthro-fused, by oxidative coupling, 199	Trifluoromethylbenziodoxoles, 60–62
Thiazolo-fused quinolinones, synthesis, 203	from 2-iodobenzoic acid, 60
Thienyl iodonium salts, preparation by iodonium transfer using PhI(CN)OTf, 81	from methoxybenziodoxole or acetoxybenziodoxole, 60
Thioacetals, cleavage to carbonyl compounds, 221	as reagents for the trifluoromethylation, 61, 276, 277
Thiocyanation, of indoles, pyrrole and arylamines,	X-ray structures, 60, 61
296	(Trifluoromethyl)iodine dichloride, CF ₃ ICl ₂
Thiocyanations, 232–235	CF ₃ ICl ₂ , X-ray crystal structure, 31
Thioketals, cleavage to carbonyl compounds, 221	preparation, 29
Thiophene dimers, preparation, by oxidative coupling of	Trifluoromethyliododifluoride, CF ₃ IF ₂
3-substituted thiophenes, 200, 202	theoretical studies, 26
Thiosulfonic S-esters, preparation by oxidation of diaryl	X-ray structure, 26, 27
disulfides or thiophenols, 219, 220	Triisopropylsilyl enol ethers, azidation to β-azido
Three–center–four–electron (3c – 4e) bond, 1, 7	adducts, 224, 225
Thyroxine, 2	1-[(Triisopropylsilyl)ethynyl]-1,2-benziodoxol-3(1 <i>H</i>)-
(–)-Tirandamycin C, synthesis, 299	one (TIPS-EBX), 59
Tosyliminoiodane, PhINTs	1,3,5-Trioxocane derivatives, synthesis, 244
aziridination of alkenes, 253	Triphenyl iodine, Ph ₃ I, 107
reagent for Hofmann rearrangement, 215	Tris(alkynyliodonium) salts, 92
α-Tosyloxylation of ketones	Trivalent iodine, structural types, 3, 4
enantioselective, catalyzed by chiral iodoarenes, 341	Twisted IBX, as catalyst for oxidation of alcohols, 361
iodobenzene-catalyzed, 339, 340	α,β -Unsaturated carbonyl compounds, synthesis by IBX
NH ₄ I-catalyzed, 370	oxidation, 293
<i>p</i> -iodotoluene-catalyzed, 340	
in the presence of a catalytic I_2 , 340	Valienamine, synthesis via iodosyl elimination, 249
under solvent-free conditions, 420	Vinyliodonium ions, ab initio MO calculations of
α -Tosyloxyketones	reactivity, 12
from alcohols, by oxidation-tosyloxylation, 170	Vinyl sulfones, chiral, synthesis by alkoxy radical
preparation from ketones and HTIB, 169, 170	fragmentation, 241
Trans influence	
in hypervalent iodine(III) compounds, 7, 11	Wighteone, regioselective synthesis, 210, 211
in hypervalent iodine(V) compounds, 11	
quantitative measurement, 11	Ylides
Trans-2-aminocyclohexanecarboxylic acid, synthesis, by	of 2,4-dihydroxyacetophenones, preparation, 103
Hofmann rearrangement, 215	of hydroxycoumarins, preparation and structure, 101
Triaryliodanes, 107	of hydroxyquinones, preparation, 102 aza analogues,
Triaryl- λ^3 -iodanes, low-temperature NMR studies, 9, 10	102
[1,2,4]Triazolo[4,3- <i>a</i>][1,4]benzodiazepine derivatives, synthesis, 206	Ynamides, by amination of terminal alkynes, 231
1,2,4-Triazolo[4,3-a]pyrimidines, synthesis, 204	Zwitterionic iodonium compounds, 101