

14. O. K. C. Tsui, H. F. Zhang, *Macromolecules* **34**, 9139 (2001).
15. F. Varnik, J. Baschnagel, K. Binder, *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.* **65**, 021507 (2002).
16. D. Long, F. Lequeux, *Eur. Phys. J. E* **4**, 371 (2001).
17. Z. Yang *et al.*, *Appl. Phys. Lett.* **94**, 251906 (2009).
18. O. K. C. Tsui, Y. J. Wang, F. K. Lee, C.-H. Lam, Z. Yang, *Macromolecules* **41**, 1465 (2008).
19. Materials and methods are available as supporting material on Science Online.
20. J.-C. Majeste, J.-P. Montfort, A. Allal, G. Marin, *Rheol. Acta* **37**, 486 (1998).
21. Y. J. Wang, O. K. C. Tsui, *J. Non-Cryst. Solids* **352**, 4977 (2006).
22. J. W. Cahn, *J. Chem. Phys.* **42**, 93 (1965).
23. A. Vrij, J. T. G. Overbeek, *J. Am. Chem. Soc.* **90**, 3074 (1968).
24. S. Peter, H. Meyer, J. Baschnagel, R. Seemann, *J. Phys. Condens. Matter* **19**, 205119 (2007).
25. S. Kawana, R. A. L. Jones, *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.* **63**, 021501 (2001).
26. S. Herminghaus, R. Seemann, K. Landfester, *Phys. Rev. Lett.* **93**, 017801 (2004).
27. J. S. Sharp, J. A. Forrest, *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.* **67**, 031805 (2003).
28. A. Dhinojwala, G. K. Wong, J. M. Torkelson, *J. Chem. Phys.* **100**, 6046 (1994).
29. R. F. Boyer, S. G. Turley, Eds., *Molecular Basis of Transitions and Relaxations* (Gordon and Breach Science Publishers, New York, 1978), p. 33.
30. We thank C. A. Angell for useful discussions and G. B. McKenna for critical reading of this paper. Support from NSF (DMR-0706096 and DMR-0908651) is gratefully acknowledged.

Supporting Online Material

www.sciencemag.org/cgi/content/full/328/5986/1676/DC1
Materials and Methods
Figs. S1 and S2
References

9 November 2009; accepted 18 May 2010
10.1126/science.1184394

The Palladium-Catalyzed Trifluoromethylation of Aryl Chlorides

Eun Jin Cho, Todd D. Senecal, Tom Kinzel, Yong Zhang, Donald A. Watson,* Stephen L. Buchwald†

The trifluoromethyl group can dramatically influence the properties of organic molecules, thereby increasing their applicability as pharmaceuticals, agrochemicals, or building blocks for organic materials. Despite the importance of this substituent, no general method exists for its installment onto functionalized aromatic substrates. Current methods either require the use of harsh reaction conditions or suffer from a limited substrate scope. Here we report the palladium-catalyzed trifluoromethylation of aryl chlorides under mild conditions, allowing the transformation of a wide range of substrates, including heterocycles, in excellent yields. The process tolerates functional groups such as esters, amides, ethers, acetals, nitriles, and tertiary amines and, therefore, should be applicable to late-stage modifications of advanced intermediates. We have also prepared all the putative intermediates in the catalytic cycle and demonstrated their viability in the process.

The introduction of the strongly electron-withdrawing trifluoromethyl group into organic molecules can substantially alter their properties (such as lipophilicity, metabolic stability, and bioavailability) that affect the use of these molecules as pharmaceuticals and agrochemicals (1–3). Additionally, trifluoromethylated organic compounds find applications as materials such as liquid crystals (2). Despite the importance of this substituent, no general catalytic method exists for the introduction of the CF₃ group onto functionalized aromatic intermediates (4).

Structurally simple benzotrifluorides are accessible by radical chlorination of toluene derivatives and subsequent chlorine-fluorine exchange under harsh conditions (5). The replacement of an aromatic halide by a CF₃ group via copper-mediated coupling proceeds under milder reaction conditions but is mainly limited to aryl iodides (6–16). A catalytic version of this process was recently reported, but only aryl iodides with electron-withdrawing substituents and some heterocycles are good substrates (17).

A palladium-catalyzed trifluoromethylation of aryl halides (Fig. 1) has the potential to overcome these limitations: The use of a trifluoromethyl


source as a transmetalating agent obviates the need for harsh reaction conditions that are required to replace individual substituents on benzylic carbon atoms with fluorine. Additionally, because many known ligands promote oxidative addition, even into unactivated aryl chlorides at low temperatures, a wide substrate scope is possible.

Mainly due to the high activation barrier for reductive elimination, the development of such a process has so far been unsuccessful. Several

complexes (4) with bidentate ligands yield either no (18, 19) or only trace amounts (20) of the benzotrifluoride products 5, even after prolonged heating at 130°C. The chelating biphosphine ligands 1,2-Bis(diphenylphosphino)ethane (dppe) and 1,3-Bis(diphenylphosphino)propane (dppp) promote the reductive elimination of 4, only at 145°C, to give PhCF₃ in 10 to 60% yield after 64 hours (19). On the other hand, the feasibility of fast Ar-CF₃ bond formation from a Pd(II) complex under mild conditions was demonstrated by Grushin and Marshall through quantitative conversion of the complex [XantphosPd(Ph)(CF₃)] to PhCF₃ upon heating to 80°C within 3 hours (21). However, the replacement of the Xantphos ligand in 3 with trifluoromethyl ions competes with transmetalation to 4, and consequently, no catalytic system with this system was reported (21, 22).

Complexes 4 are typically prepared from complexes 3, where X = Br or I, by treatment with TMSCF₃ (TMS, trimethylsilyl) and a fluoride source such as CsF, thereby using the formation of a silicon-fluorine bond as driving force (18–20). The challenge of using trifluoromethylsilanes in the presence of fluoride originates in the fluoride-initiated self-decomposition of R₃SiCF₃ to give R₃SiF and difluorocarbene (23). In a catalytic setting, where elevated temperatures are presumably required to promote reductive elimination from 4

Fig. 1. Generalized catalytic cycle for aryl trifluoromethylation (L = ligand; Ar = aryl; X = Cl, Br, I, triflate).



Department of Chemistry, Massachusetts Institute of Technology (MIT), Cambridge, MA 02139, USA.

*Present address: Department of Chemistry and Biochemistry, University of Delaware, Newark, DE 19716, USA.

†To whom correspondence should be addressed. E-mail: sbuchwal@mit.edu

to **5**, transmetalation must be substantially faster than this process.

The oxidation of Pd(II)-CF₃ complexes with a F⁺ reagent provides Pd(IV) complexes that readily reductively eliminate benzotrifluorides (20). The catalytic trifluoromethylation of arenes via C-H activation, oxidation of the Pd(II) intermediate with an electrophilic CF₃⁺ source, and final reductive elimination have recently been reported, but are limited to substrates with specific directing groups (24).

We report the development of a palladium-catalyzed procedure to transform aryl chlorides into their trifluoromethylated analogs using TESCf₃ (TES, triethylsilyl) and KF. High functional-group tolerance under relatively mild conditions is exhibited, and a wide range of substrates, including heteroaryl ones, can be efficiently converted to the desired products. Mechanistic studies suggest that the reaction proceeds via a classical Pd(0)-Pd(II) catalytic cycle, as shown in Fig. 1.

Ligand **6** (BrettPhos) (Fig. 2A) has successfully been employed in challenging amination and fluorination cross-coupling reactions (25, 26). We prepared the oxidative-addition complex **8** and examined numerous trifluoromethyl sources to identify conditions under which both transmetalation and reductive elimination would proceed (table S1) (27). Although most reagents failed to produce a product, the mixture of TESCf₃ and CsF in tetrahydrofuran (THF) at 60°C provided **9** in a promising yield of 28% (Fig. 2B). This result confirms that **6** does indeed promote reductive elimination to form Ar-CF₃ bonds and provides a starting point for the development of a catalytic procedure.

With 3 mole percent (mol %) [(allyl)PdCl]₂ and 12 mol % of ligand **6**, benzotrifluoride **9** was formed in 7% yield from aryl chloride **10** with TESCf₃ and CsF at 110°C. We next investigated several combinations of TMSf₃ or TESCf₃ with simple fluoride salts and found that the highest yield was obtained using TESCf₃ with KF, demonstrating that the catalytic formation of **9** from **10** is possible with these transmetalating agents (Fig. 2C). Full conversion of **10** was achieved by switching the solvent to dioxane and performing the reaction at 120°C, providing **9** in 80% yield.

We studied the performance of other ligands under these conditions and found that **6** was the best ligand for this transformation (table S2). Most other monodentate biaryl phosphine ligands gave lower, but still observable, amounts between 5 and 20% of product **9**. No reaction occurred, however, when Xantphos was used.

The palladium-catalyzed process expands the scope to aryl chlorides and exhibits compatibility with a wide range of functional groups (Fig. 3). Both electron-poor and electron-rich aryl chlorides are suitable substrates and provide the trifluoromethylated products in good yields. More importantly, heteroaromatic substrates such as indoles, carbazoles, quinolines, and benzofurans can be efficiently transformed into their trifluoromethylated analogs. When using ligand **6**, we found that





Fig. 2. (A) Ligands used in this study. Me, methyl; *i*Pr, isopropyl. (B) Best result from a reagent screen to convert complex **8** into benzotrifluoride **9**. *n*Bu, straight-chain butyl. (C) Identification of an optimal combination of trifluoromethyl source and activator for the catalytic conversion of **10** to **9**. R, methyl or ethyl.

Fig. 3. Scope of the palladium-catalyzed trifluoromethylation of aryl and heteroaryl chlorides. Isolated yields are based on an average of at least two runs. Minor amounts (2 to 5%) of reduced starting material (Ar-H) were usually observed. In a typical experiment, a solution of the palladium source and ligand **6** or **7** in dioxane was added to spray-dried KF and the aryl chloride. After addition of TESCf₃, the reaction was stirred at 120 to 140°C for 6 to 20 hours. Because KF is hygroscopic, all reactions were set up in a nitrogen-filled glovebox to prevent the hydrolysis of TESCf₃ in the course of the reaction. Et, ethyl; Hex, hexyl; Ph, phenyl.



* 120 °C; † the compound could not be separated from the reduced Ar-H product. The given yield was estimated by gas chromatography; ‡ 2 mol % [(allyl)PdCl]₂, 6 mol % **6**; § 8 mol % Pd(dba)₂, 12 mol % **6**; || 12 mol % **6**

ortho-substituted substrates gave the corresponding products in low yields only. Switching to the less bulky ligand RuPhos (7) (Fig. 2A) (28) provided the desired ortho-substituted products **11r–v** in excellent yields. Scale-up proved to be straightforward; products **11j** and **11b** were prepared on 2- and 5-mmol scales, respectively, in the same yields as those reported for the 1-mmol-scale reactions.

Esters, acetals, amides, nitriles, ethers, dialkylamines, and a number of heteroaromatic substituents are tolerated. However, substrates bearing aldehydes or ketones are not suitable. Furthermore, substrates cannot contain unprotected OH or NH groups, presumably because of protonation of the CF₃ anion to form fluorosulfonate, reaction at the silicon center of TSCF₃, and/or competing coordination to the palladium center.

To gain insight into the reaction mechanism, we prepared the presumptive Pd–CF₃ intermediates **13** and studied their reductive elimination to yield benzotrifluoride products. Treatment of complexes **12** with TSCF₃/CsF at room temperature in THF allowed the isolation of [6•Pd(Ar)(CF₃)] complexes **13** (Fig. 4A). The compounds exhibit a characteristic quartet in the ³¹P–nuclear magnetic resonance (NMR) spectrum and a doublet in the ¹⁹F–NMR spectrum with a coupling constant of ~45 Hz. The Pd atom in the crystal structures of **13a** (fig. S1) and **13b** (Fig. 4B) is coordinated by the upper-ring methoxy group of the ligand **6** and not by the ipso carbon atom of the lower aromatic ring.

We studied the reductive elimination of complexes **13** in dioxane at 80°C via ¹⁹F–NMR and found first-order decay to give benzotrifluorides **14** in nearly quantitative yield. The rate constants for both the decomposition of **13a** and

13b are almost identical (Fig. 4A and fig. S2 and S4). This surprising result is paralleled by density functional theory calculations that predict an activation energy of ~22 kcal mol^{−1} for both complexes (29, 30). In comparison to the ground states, the calculated Pd–CF₃ distance in the transition states is substantially elongated, whereas the distance between the Pd atom and the aryl ring remains essentially unchanged, suggesting that the main contribution to the activation energy is the breaking of the strong Pd–CF₃ bond. Because the strength of this bond is only minimally influenced by the substituent on the aryl ring, similar rate constants are observed.

When complex **13a** was heated in the presence of excess methyl 4-chlorobenzoate, the oxidative-addition complex **12a** was formed in addition to product **14a**, thus closing the catalytic cycle. The yield and rate of benzotrifluoride formation were identical in the presence or absence of aryl chloride (fig. S3), which implies that reductive elimination affords a Pd(0) species that then undergoes oxidative addition to form **12a**. Therefore, we believe that these reactions proceed via a classical Pd(0)/Pd(II) catalytic cycle, as proposed in Fig. 1.

In preliminary experiments, we have demonstrated that this process is applicable, in somewhat lower yields, to aryl bromides and aryl triflates. We are currently seeking to develop a better understanding of the overall reaction mechanism, as well as to render this process more generally useful and practical. We hope to accomplish this by broadening its substrate scope, lowering the quantity of catalyst necessary, developing milder reaction conditions, and using less expensive and more environmentally friendly trifluoromethylating agents.




Fig. 4. (A) Formation of and reductive elimination from [6•Pd(Ar)(CF₃)] complexes. *t*_{1/2} are the half-lives of the first-order reductive elimination kinetics. (B) X-ray structure of complex **13b**. ORTEP (31) drawing at 50% probability; hydrogen atoms are omitted for clarity.

References and Notes

- D. Cartwright, in *Organofluorine Chemistry*, R. E. Banks, B. E. Smart, J. C. Tatlow, Eds. (Plenum, New York, 1994), pp. 237–262.
- P. Kirsch, *Modern Fluoroorganic Chemistry* (Wiley-VCH, Weinheim, Germany, 2004).
- K. Müller, C. Faeh, F. Diederich, *Science* **317**, 1881 (2007).
- A. M. Thayer, *Chem. Eng. News* **84**, 15 (2006).
- F. Swarts, *Bull. Soc. Chim. Belg.* **24**, 309 (1892).
- V. C. R. McLoughlin, J. Thrower, *Tetrahedron* **25**, 5921 (1969).
- Y. Kobayashi, I. Kumadaki, *Tetrahedron Lett.* **10**, 4095 (1969).
- N. V. Kondratenko, E. P. Vechirko, L. M. Yagupolskii, *Synthesis* **1980**, 932 (1980).
- K. Matsui, E. Tobita, M. Ando, K. Kondo, *Chem. Lett.* **10**, 1719 (1981).
- D. M. Wiemers, D. J. Burton, *J. Am. Chem. Soc.* **108**, 832 (1986).
- G. E. Carr, R. D. Chambers, T. F. Holmes, D. G. Parker, *Perkin Trans. 1* **1988**, 921 (1988).
- J. H. Clark, M. A. McClinton, R. J. Blade, *Chem. Commun.* **1988**, 638 (1988).
- Q. Y. Chen, S. W. Wu, *Chem. Commun.* **1989**, 705 (1989).
- H. Urata, T. Fuchikami, *Tetrahedron Lett.* **32**, 91 (1991).
- F. Cottet, M. Schlosser, *Eur. J. Org. Chem.* **2002**, 327 (2002).
- G. G. Dubinina, H. Furutachi, D. A. Vici, *J. Am. Chem. Soc.* **130**, 8600 (2008).
- M. Oishi, H. Kondo, H. Amii, *Chem. Commun. (Cambridge)* **2009**, 1909 (2009).
- D. A. Culkin, J. F. Hartwig, *Organometallics* **23**, 3398 (2004).
- V. V. Grushin, W. J. Marshall, *J. Am. Chem. Soc.* **128**, 4632 (2006).
- N. D. Ball, J. W. Kampf, M. S. Sanford, *J. Am. Chem. Soc.* **132**, 2878 (2010).
- V. V. Grushin, W. J. Marshall, *J. Am. Chem. Soc.* **128**, 12644 (2006).
- D. Naumann *et al.*, *Z. Anorg. Allg. Chem.* **630**, 746 (2004).
- W. Tyrra *et al.*, *Chemistry* **11**, 6514 (2005).
- X. Wang, L. Truesdale, J. Q. Yu, *J. Am. Chem. Soc.* **132**, 3648 (2010).
- B. P. Fors, D. A. Watson, M. R. Biscoe, S. L. Buchwald, *J. Am. Chem. Soc.* **130**, 13552 (2008).
- D. A. Watson *et al.*, *Science* **325**, 1661 (2009); published online 13 August 2009 (10.1126/science.1178239).
- Supporting material is available on Science Online.
- J. E. Milne, S. L. Buchwald, *J. Am. Chem. Soc.* **126**, 13028 (2004).
- M. J. Frisch *et al.*, Gaussian 03, Revision E.01 (Gaussian, Inc., Wallingford CT, 2004).
- The employed level of theory is B3LYP/6-311++G(2d,p)/LanL2Dz/B3LYP/6-31G(d)/LanL2Dz.
- ORTEP, www.ornl.gov/sci/ortep/ortep.html.
- We thank the NIH for financial support of this project (grant GM46059) and Merck, Nippon Chemical, Boehringer Ingelheim, and BASF for gifts of chemicals and additional funds. T.K. thanks the Alexander von Humboldt Foundation for a Feodor Lynen postdoctoral fellowship. The Varian NMR instrument that we used was supported by the NSF (grants CHE 9808061 and DBI 9729592). We also thank P. Müller (MIT Chemistry Department, X-Ray Diffraction Facility) for obtaining the crystal structures of **13a** and **13b**. Cambridge Crystallographic Data Centre (CCDC) 771779 and 771780 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from the CCDC via www.ccdc.cam.ac.uk/data_request/cif.

Supporting Online Material

www.sciencemag.org/cgi/content/full/328/5986/1679/DC1
SOM Text
Figs. S1 to S5
Tables S1 to S3
References

6 April 2010; accepted 12 May 2010
10.1126/science.1190524

The Palladium-Catalyzed Trifluoromethylation of Aryl Chlorides

Eun Jin Cho, Todd D. Senecal, Tom Kinzel, Yong Zhang, Donald A. Watson and Stephen L. Buchwald

Science **328** (5986), 1679-1681.
DOI: 10.1126/science.1190524

Three Fluorines at Once

The trifluoromethyl (CF₃) group is playing an increasingly important role in the design of pharmaceutical and agrochemical compounds. CF₃ is a powerful attractor of electron density within a given molecular framework, and recently fluorine-hydrocarbon interactions have emerged as a distinct complement to the more traditional hydrophilic/hydrophobic interplay that governs docking between small molecules and proteins. **Cho et al.** (p. 1679) now present an efficient method for appending CF₃ groups to a broad range of aryl substrates. A carefully optimized palladium catalyst was able to speed up a critical elimination step that has plagued previous efforts to realize a general solution to this synthetic challenge.

ARTICLE TOOLS

<http://science.sciencemag.org/content/328/5986/1679>

SUPPLEMENTARY MATERIALS

<http://science.sciencemag.org/content/suppl/2010/06/21/328.5986.1679.DC1>

REFERENCES

This article cites 25 articles, 2 of which you can access for free
<http://science.sciencemag.org/content/328/5986/1679#BIBL>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)