

# BacStop™

## Bactericide / Fungicide

### CAN BE USED FOR ORGANIC PRODUCTION

This product has not been registered by the US Environmental Protection Agency. Anjon AG, Inc. represents that this product qualifies for exemption from registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA)25b.

**DIRECTIONS FOR USE**  
**General Information and Tips for Best Performance**  
BacStop™ is a concentrate and is designed for use both in greenhouses and outdoors. BacStop is intended to be mixed with water and applied with pressurized power operated sprayers or delivered through drip irrigation systems. When diluting, it is recommended to add approximately half the water to the applicator tank, then add the proper amount of BacStop in accordance with the rate table, agitate, then add the remaining amount of water. Then thoroughly agitate the mixture to insure proper suspension. BacStop is to be used as a contact bactericide/fungicide pest control agent.

BacStop can be tank-mixed with other products unless prohibited by the product label. A small volume test mix is recommended to ensure compatibility. BacStop can cause a phytotoxic damage to green foliage in higher concentrations. If tank-mixed with another product a small scale test to ensure lack of phytotoxicity of the combination is recommended.

**DIRECTIONS FOR USE**  
BacStop is a multi-purpose broad spectrum bactericide/fungicide. BacStop can be used for treatment of all food crops and ornamentals. BacStop can also be used for soil applications to aid in reducing the damaging effects of soil borne bacterial and fungal diseases.  
**General Application Notes**  
For corn and wheat pathogens suppression it is recommended to use EF400 Fungicide and BacStop in combination for effective control of bacteria and fungal pathogens. This blend is synergistic and is common that both types of disease are present at the same time. EF400 Fungicide is an ideal partner to the BacStop.

**Companion Blend and Tank Mix**  
Companion blends include EF400 Fungicide, EF300 Insecticide with BacStop.  
**Application Timing**  
Apply diluted spray when high humidity for the development of disease symptoms in crops or at first sign of disease and continue applications at 7 to 10 day intervals or until disease symptoms are gone.  
**Minimum Time for Application to Harvest**  
Spraying or drip application of the product can occur up to the time of harvest on all intended crops.  
**Application Rates for BacStop™**  
For spray application rates see Table.

**Application Method and Equipment**  
BacStop can be sprayed on foliage, the surface of soil or injected into a drip irrigation system at the rates on the appropriate tables. When sprayed on foliage allow 12 hours of a dry period so that irrigation or rain does not wash off application. When applied topically to soil, immediately follow application of BacStop with 1/2 inch of water to wash into soil.  
For drip applications apply to soil that is with humidity level of 70% or more. After BacStop has cleared lines follow with enough water to wash down to root level. Always wash into soil after application with plain water to make sure that BacStop penetrates down to the depth of the root ball. Consult spray nozzle and accessory catalogues for specific information on proper equipment calibration. For best coverage use small droplets.  
**Storage:** Store in original UV protected containers only. Store at temperatures between 41°F to 85°F (5°C to 30°C).  
**Disposal:** Wastes resulting from using this product may be disposed of on site. Dispose of empty jug in a sanitary land fill as allowed by state and local authorities.

# BacStop Bactericide

**Active Ingredients:**Clove 2.0%, Thyme 2.0%, Cinnamon 1.5%, Garlic 1.0%, Peppermint 1.0%-Total (7.5%)  
**Other Ingredients:** Bentonite, Carrageenan, Citrus Pulp, Kaolin, Palm Oil, Purified Water - Total (92.5%)

KEEP OUT OF REACH OF CHILDREN

## CAUTION PRECAUTIONARY STATEMENTS

May causes temporary eye and skin irritation. Do not get in eyes, on skin, or on clothing. Wear protective eyewear such as goggles, face shield, or safety glasses, and chemical resistant gloves. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

### FIRST AID

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. If irritation persists, call a poison control center or doctor for further advice.  
**If on skin:** Take off contaminated clothing, rinse skin immediately with plenty of soap and water for 15-20 minutes. If irritation persists, call a poison control center or doctor for further advice.  
**If swallowed** - Call doctor or poison control center immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by poison control center or doctor. Do not give anything to an unconscious person.  
**If inhaled** - Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible.

### CONDITIONS OF SALE AND WARRANTY

Anjon AG, Inc. warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. Anjon AG, Inc. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR USE. Critical and unforeseeable factors beyond Anjon AG, Inc's. control prevent it from eliminating all risks in connection with the use of this product. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather conditions, presence of other materials, drift to other crops or property, the manner of use or application, or failure to follow label directions will be assumed by the Buyer or User. Handling, storage and use of the product by Buyer and User are beyond the control of Anjon AG, Inc. In no case shall Anjon AG, Inc. be held liable for consequential, special or indirect damages resulting from the use of this product. The limit of Anjon AG, Inc. liability shall be the purchase price for the quantity involved.  
BacStop is a Trademark of Anjon AG, Inc. • Copyright Anjon AG, Inc., 2015

**Anjon AG, Inc.**  
179 Niblick Rd., #405 Paso Robles, CA 93446  
855-434-0300  
www.anjonag.com

**NET CONTENTS**  
**1 GALLON**  
**(3.79L)**

# BacStop Bactericide

Table 1 Specific Crops	Target Diseases	Preventive Rate fl oz	Curative Rate fl oz	Rate per Acre Gallons	Application Directions & Interval
<b>Apple Pear</b>	Apple Scab Fire Blight	6 6	12 12	60-100	Begin applications from 1/2 in. tip to harvest, 7 day intervals.
<b>Blueberry</b>	Mummy Berry Anthracnose Fruit Rot Bacteria Soft Rot	9 1 9	12 16 12	30-50	Spray at green tip, early pink bud, pink bud, early bloom, mid bloom, petal fall and fruit set
<b>Citrus</b>	Citrus Canker	8	12	60-100	
<b>Cucurbits</b>	Bacterial Leaf Spot	6	12	25-30	Preventative - make application every 7 days. After first symptoms observed, apply every 3-5 days until control is established. Recommended to use in combination with EF400.
Cucumber, squash, pumpkin, melon, pickle	Bacteria Wilt	8	12	25-40	
	Bacteria Fruit Blotch	6	10	30-40	
	Downey Mildew	6	8	30-40	
	Powdery Mildew	6	8	25-40	
<b>Cotton</b>	Bacteria Blight	10	14	10-15	Apply in combination with EF400
<b>Dry Edible Beans and Green Beans</b>	Common Blight Halo Blight Brown Spot	6 6 6	9 9 8	20 20 20	Apply 3rd trifoliolate, R1, R4, R7. Recommended use in combination with EF400 to reduce fungal problems.
<b>Field Corn</b>	Anthracnose Goss's Wilt	8 12	12 16	20 20	Anthracnose - Spray at 4th leaf and continue every 10-14 days or 7-10 days under high pressure conditions. For Goss's Wilt apply at V-4 stage and row tassel. Apply every 7-10 days until disease is suppressed from stalk.
<b>Grape</b>	Botrytis Bunch Rot Phomopsis cane & Leaf Spot	6 8	8 12	40-50 40-50	Spray at full bloom, berry touch and veraizon.
<b>Onion</b>	Anthracnose - onion smudge	8	12	25-40	Begin shortly after emergence and spray every 7 days
<b>Peppers</b>	Bacteria Leaf Spot	6	8	25-40	If first symptoms are observed apply every 5-7 days until control is attained.
<b>Potatoes</b>	Common Scab Bacteria Wilt Early Blight Late Blight	6-8 6 6 6	6-8 6 6 8	25-30 25-30 25-30 25-30	For soil treatment for Scab use in combination with EF400. After application water into soil to reach root area. For topical spray use as recommended. Begin spraying at first sign of disease.
<b>Potato Seed Preplant</b>	To help stop the spread of Scab and other bacterial and fungal disease	BacStop 3	EF400 10	20	Use in combination with EF400. Use fine spray and allow to dry before planting. Good coverage is critical.
<b>Soybeans</b>	Bacteria Blight	6	8	20	Application schedule: R1, R4, R-6
<b>Strawberry</b>	Bacteria Blight Bacteria Wilt Botrytis Powdery Mildew	4 4 4 4	8 8 8 4	25-40 25-40 25-40 25-40	Product should be applied on a preventative program. For soil borne diseases can applied through drip irrigation system. Use in combination with EF400
<b>Tomatoes</b>	Bacteria Leaf Spot Bacterial Canker Bacteria Wilt Early Blight Late Blight	4 4 4 4 4	6 6 6 6 6	25-40 25-40 25-40 25-40	Begin spraying at first signs of disease onset and maintain a 7-10 day interval during periods of high infection
<b>Wheat</b>	Fusarium Head Blight Bacteria Head Blight	4 6	8 8	20 20	4th to 5th leaf followed at 14 days later
<b>Ornamentals</b>	Bacterial Leaf Spot	4	6	25-50	Preventative - apply every 7 days Curative - Apply at first symptoms and continue every 5 day until control established.

0516VT