

# Probability

---

Many events can't be predicted with certainty. We use the tools of probability to quantify how likely they are to happen. To reason about probabilities, we'll use a data type called a *probability space*. We'll build up this data type over the course of this set of lecture notes.

Probability builds on top of set theory and logic. We can think of logic as a limiting case of probability, where everything is either completely certain or completely uncertain.

## Atomic events

---

We start from a *universe* or *sample space*, a set  $U$  of mutually exclusive and exhaustive possible outcomes. The elements of  $U$  are called *simple* or *atomic* events. Often these simple events will be interpretable: e.g., the number of spots on the top face after we roll a die (like this:  $\boxplus$ ), or the card we draw from a deck (like this:  $7 \spadesuit$ ). We assign probabilities to atomic events:  $P(a)$  is the probability that  $a \in U$  will occur, and  $P(\bar{a}) = 1 - P(a)$  is the probability that it will not.

*For our purposes,  $U$  will always be finite. If we want to deal with continuous outcomes, a good way to think about them is to discretize: pretend that there are a very large but finite number of events, spaced out so that one is always very close to any possible continuous outcome. Since this viewpoint is for understanding rather than computation, there's no limit on how big we can make  $U$ : e.g., we could have one event for every 64-bit floating point number, or even every tuple of 100 64-bit numbers. Discretization is actually better in many cases than trying to deal directly with continuous outcomes: probability on continuous spaces holds a lot of mathematical traps, such as [hidden infinities](#) and [counterintuitive behaviors](#).*

We'll take the viewpoint that probabilities are *subjective*: I can assign any probabilities I want, so long as I follow the rules described here for manipulating them. And I can assign probabilities to any outcomes that I can imagine measuring, no matter whether I think of the process that determines the outcomes as being random.

*What does the statement  $P(a) = \frac{1}{3}$  mean? One common interpretation is that, if I reset the world and ran it forward many times, measuring whether event  $a$  happened in each trial, I'd see it happen in a third of them. This interpretation makes sense for something like a biased coin that we can flip many times. But it is not necessarily helpful in general: first, it's not really possible to reset the world. Second, we might want to assign a probability to an event that is deterministic but unknown to us, so that repeated trials don't yield independent measurements. Instead, in*

*such cases we interpret the statement as a measure of my subjective confidence in  $a$ : I'd be willing to take 2 : 1 odds in a bet on  $a$ , which is the same odds that would have me break even if betting on a biased coin with  $P(H) = \frac{1}{3}$ .*

Probabilities are always nonnegative; an impossible event has probability 0. Probabilities always sum to 1 over the universe:

$$\sum_{e \in U} P(e) = 1$$

That means that each individual event has probability at most 1; an event with probability 1 is certain to happen.

*In continuous spaces, we measure probability density instead of probability. Probability density is nonnegative, but can be greater than 1.*

## Compound events

---

A compound event is a subset of the universe,  $E \subseteq U$ . The probability of  $E$  is the sum of the probabilities of atomic events in  $E$ :

$$P(E) = \sum_{a \in E} P(a)$$

We are certain to get some event in the universe:

$$P(U) = \sum_{a \in U} P(a) = 1$$

We'll sometimes abuse notation and conflate  $a$  with  $\{a\}$ : that is, we'll interchange an atomic event and a compound event with only one element.

Since events are sets, we can combine them using set operations: given events  $A$  and  $B$ ,

- $A \cup B$  means that we get some atomic event in the union of  $A$  and  $B$ ; that is,  $A \cup B$  is the event that *either*  $A$  or  $B$  happens
- $A \cap B$  is the event that *both*  $A$  and  $B$  happen
- $A \setminus B$  is the event that  $A$  happens but  $B$  does not happen
- $A^C = U \setminus A$  is the event that  $A$  does not happen

As you can see from the examples above, set operations correspond to logical

combinations of events. So we'll sometimes use logical notation as well:

- $A \vee B$  is the same as  $A \cup B$
- $A \wedge B$  is the same as  $A \cap B$
- $\neg A$  or  $\bar{A}$  is the same as  $A^C$

We can make the correspondence with logical notation precise:

- Logical predicates correspond to events. For example, in our die-roll example, the predicate `even` corresponds to the set of even-valued outcomes.
- Logical functions can be used to build predicates. For example, if we have a function `suit` that extracts the suit of a card, then the expression `suit = ♠` corresponds to the set  $\{a \mid \text{suit}(a) = \spadesuit\}$ , or

$$\{A\spadesuit, 2\spadesuit, 3\spadesuit, 4\spadesuit, 5\spadesuit, 6\spadesuit, 7\spadesuit, 8\spadesuit, 9\spadesuit, 10\spadesuit, J\spadesuit, Q\spadesuit, K\spadesuit\}$$

Note that we have omitted the argument  $a$  to the function `suit`: by convention, the atomic event is implicitly an argument of every function or predicate.

- General expressions correspond to the set of atomic events that satisfy them. For example, the expression `suit = ♠  $\wedge$  even` corresponds to the set


$$\{2\spadesuit, 4\spadesuit, 6\spadesuit, 8\spadesuit, 10\spadesuit, Q\spadesuit\}$$

which is  $\{a \mid \text{suit}(a) = \spadesuit \wedge \text{even}(a)\}$ .

Exercise: what is the set notation for the logical event  $\neg(A \rightarrow B)$ ?

## Venn diagrams

We can visualize our universe and the probabilities of various events using a *Venn diagram*: we picture the universe as a large rectangle, and events as shapes that are subsets of the rectangle. Every point in the rectangle corresponds to a different atomic event. The overlap between shapes tells us which compound events intersect, that is, which can occur simultaneously.



For example, if we are picking a random food to eat for dinner, the universe is the set of all possible foods we might pick, corresponding to the whole rectangle. The event that we pick something yummy is the larger circle, and the event that we pick a vegetable is the smaller circle. The overlap between the two circles is the set of yummy vegetables, which includes the atomic event *zucchini* — at least, according to your professor. The area outside the two circles corresponds to the set of outcomes which are neither yummy nor vegetables.


Often we try to make the areas of shapes within the Venn diagram proportional to their probability. But sometimes we can't or don't want to — for example, if we don't know those probabilities yet.

Exercise: where in the above diagram does the event that we pick pizza fit? What about Brussels sprouts? What about escargot? (We realize that the answers are subjective.)

## Experiments

We can use probabilities to describe *experiments* that we are thinking of conducting. We make a probability space that describes what we think might happen; atomic events represent possible outcomes of the experiment, and the universe contains all possible outcomes.

For example, suppose we flip a coin twice. We can make a probability space that says that each sequence of heads and tails is equally likely:



Here the universe is  $\{HH, HT, TH, TT\}$ , and we've assigned probability  $\frac{1}{4}$  to each of these atomic events.

In a realistic experiment, we might not observe the distinction between all atomic events. For example, above we might just observe the total number of heads instead of the exact sequence of coin flips. We can describe what we observe using compound events: in this case,  $\{HH\}$ ,  $\{HT, TH\}$ , and  $\{TT\}$ .

## Uniform distributions

If all atomic events are equally likely, our probability space represents a *uniform distribution*. This distribution is often a default choice for an initial model. But we also often need *nonuniform distributions*, those in which different atomic events have different probabilities.

For example, a fair die is modeled well by a uniform distribution:  $P(1) = P(2) = P(3) = P(4) = P(5) = P(6) = \frac{1}{6}$ . But a weighted die needs a non-uniform distribution. If rolling a 6 is four times as likely as any other outcome, we have  $P(1) = P(2) = P(3) = P(4) = P(5) = \frac{1}{9}$  and  $P(6) = \frac{4}{9}$ .

# Working with probabilities

---

In small probability spaces we can just list out all the atomic events, and calculate probabilities of compound events by the sum rule. But it's common to have so many atomic events that we have no hope of listing them all: for example, there are more possible shuffles of a 52-card deck than there are atoms in the Earth. So, we need tools to help us do the calculations without explicitly enumerating everything. There are *lots* of such tools; but in the rest of these notes we'll cover a few of the most useful.

## Disjoint union

---

If  $A$  and  $B$  are disjoint events, meaning that  $A \cap B = \emptyset$ , then

$$P(A \cup B) = P(A) + P(B)$$

This follows directly from our definitions:

$$P(A \cup B) = \sum_{e \in A \cup B} P(e) = \sum_{e \in A} P(e) + \sum_{e \in B} P(e) = P(A) + P(B)$$

Disjoint events are also called *mutually exclusive*.

The disjoint union rule also works for multiple sets: if the sets  $A_i$  are mutually exclusive, meaning they are pairwise disjoint, then

$$P(\bigcup_i A_i) = \sum_i P(A_i)$$

A useful generalization of the disjoint union rule is the *union bound*: for any  $A$  and  $B$ , disjoint or not, we have


$$P(A \cup B) \leq P(A) + P(B)$$

We'll prove this in the next section.

## Non-disjoint union

---

We can use the tools above to figure out what happens when we take a union of sets that overlap, like in our food example from before.



Let  $A$  be the event that we pick a vegetable, and  $B$  be the event that we pick a yummy food. How can we calculate  $P(A \cup B)$ ?

We can split the set  $A \cup B$  into three disjoint parts:  $A \setminus B$ ,  $B \setminus A$ , and  $A \cap B$ . So,

$$P(A \cup B) = P(A \setminus B) + P(B \setminus A) + P(A \cap B)$$

We can also split  $A$  and  $B$  into two pieces each:

$$P(A) = P(A \setminus B) + P(A \cap B)$$

$$P(B) = P(B \setminus A) + P(A \cap B)$$

Adding these together,

$$P(A) + P(B) = P(A \setminus B) + P(B \setminus A) + 2P(A \cap B)$$

Since  $P(A \cap B) \geq 0$ , we see that  $P(A) + P(B) \geq P(A \cup B)$ ; this is the union bound we described above. By subtracting the above expressions for  $P(A) + P(B)$  and  $P(A \cup B)$  we can figure out the exact difference:

$$P(A) + P(B) - P(A \cup B) = P(A \cap B)$$

Exercise: Suppose we stand on a street corner in Pittsburgh and survey people. We ask two questions: do you like computer science, and do you watch the Steelers. You find that 67% of passers-by respond yes to the first question, and 90% respond yes to the second. Furthermore, 65% respond yes to both questions. What is the

## Logical rules

We've already talked about the relationship between logical and set-based descriptions of events. This relationship means that there is a probability inference rule for every logical inference rule. For example, De Morgan's laws:

$$P(\neg(A \wedge B)) = P(\neg A \vee \neg B)$$

$$P(\neg(A \vee B)) = P(\neg A \wedge \neg B)$$

Perhaps more subtly, if one logical statement implies another,  $\psi \rightarrow \phi$ , then  $P(\phi) \geq P(\psi)$ . For example, if being human implies being mortal, then  $P(\text{human}) \geq P(\text{mortal})$ . This follows since the event  $\psi$  has to be a subset of the event  $\phi$ .

## Combinatorics

If we start from a uniform distribution over our entire probability space, then calculating the probabilities of events boils down to counting: how many atomic events satisfy a given property? For this purpose, combinatoric tools can be very useful. The general strategy is to describe a compound event using a sequence of simpler choices, then count the ways to make these choices, so that we can get an exhaustive description of all the atomic events that make up a compound event. The key difficulty is to make sure to count each atomic event exactly once.

For example, what is the probability of drawing three of a kind in a five-card hand? The atomic events here are the possible five-card hands:  $52!/47!$  of them. To get a three of a kind, we must pick what rank to have three of (13 choices), which card of that rank to exclude (4 choices), and what two other cards to include in the hand ( $48 \cdot 47$  choices). Finally, we need to decide what order we draw these cards ( $5!$  choices). Every way of making these choices describes a different three-of-a-kind hand, and all such hands can be constructed with some sequence of choices. So the total number of atomic events that correspond to three-card hands is  $13 \cdot 4 \cdot 48 \cdot 47 \cdot 5!$ , and the probability of getting one is

$$\frac{13 \cdot 4 \cdot 48 \cdot 47 \cdot 5!}{52 \cdot 51 \cdot 50 \cdot 49 \cdot 48}$$



# Extensions

---

Probability spaces are extremely expressive, but they don't cover every possible situation. Here are a few examples of situations that might need additional tools.

First, what if I think that I'm not the only agent in the universe? That is, I believe there are other intelligent beings who predict my actions and try to optimize their own actions in response. It seems hard to assign probabilities to the actions of these agents.

For another example, suppose two people disagree on the probabilities of some outcomes. They can each build a separate probability space; but how should they talk with each other about the probabilities of events that they both can observe? How should they interpret each other's behavior?

For a third example, exact probability calculations can be very expensive. What should I do if I can't afford the necessary computation? How should I predict the behavior of a computationally limited agent?

For a final example, the conclusions I reach by probabilistic reasoning can be very sensitive to my starting model. Can I recognize when one of my assumptions causes a large change in my conclusions? Can I design a more stable reasoning process?