

MolView manual

This is the manual for MolView version 2.2

MolView is a free web application hosted on molview.org

You can contact info@molview.org if you have any questions

Table of Contents

1.	Drawing structural formulas	3
	Top toolbar	
	Left toolbar	
	Right toolbar	
2.	Finding structures	
	Tools	
	Link	
	Export	
	Information card	
	Spectroscopy	5
	3D model resource	
	Advanced search	6
4.	Spectroscopy	7
	Export data	7
5.	3D model	
	Reset	8
	Representation	8
	Background	8
	Engines	8
	Model transformation	9
	Crystallography	9
	Fog and clipping	9
6.	Protein display	.10
	Show bio assembly	.10
	Chain representation	.10
	Chain coloring	.10
· ·		.11
	Clear	
	High Quality	.11
	Calculations	.11
	Moacurement	11

1. Drawing structural formulas

You can draw structural formulas using the Sketcher component.

Top toolbar

The top toolbar contains all general editing tools. The *2D to 3D* tool converts the structural formula into a 3D molecule which is displayed in the model window. If you don't want that implicit hydrogen atoms are stripped from the structural formula after loading, you have to toggle the hydrogen stripping button off (third from right)



Top toolbar, from left to right: delete all, erase, undo, redo, move atoms/bonds, center view, rectangle select, lasso select, clear selection, toggle hydrogen stripping, clean structural formula, 2D to 3D conversion

Left toolbar


In the left toolbar, you can select a tool you want to use in order to modify or extend the structural formula. You can draw a carbon chain using the chain tool (11th button in image) Just click a start point or atom and drag a chain.



Left toolbar (rotated), from left to right: single bond, double bond, triple bond, up/down bond, benzene, cyclopropane, butane, pentane, hexane, heptane, chain, charge+, charge-

Right toolbar

The right toolbar contains some common elements and a periodic table tool in case you want to pick another element. You can select an atom and click an existing atom in order to replace it with the selected atom. Note that you can only replace existing atoms. In order to add an atom, you will first have to add a new bond using the tools from the left toolbar.




Right toolbar (rotated), from left to right: hydrogen, carbon, nitrogen, oxygen, sulfur, phosphorus, fluorine, iodine, chlorine, bromine, table of elements, info (about dialog)

2. Finding structures

You can load molecules from large databases like PubChem and RCSB using the search form located on the left side of the menu-bar. Just type what you are looking for and a list of available molecules will appear.

You can also click on the dropdown button next to the search field to select a specific database. This will perform a more extensive search on the selected database. Currently, three big databases are supported:

- PubChem
- The RCSB Protein Data Bank
- The Crystallography Open Database



The search form

3. Tools

The **Tools** menu contains several utility functions which are listed below.

Link

You can embed or share a specific compound, macromolecule or crystal using the provided URL or HTML code. Note that the linked structure is the one which is currently displayed in the model window. You can also copy the URL from the address bar in order to link to the current structure.

Export

Export options in the Export menu:

- Structural formula image: sketcher snapshot (PNG with alpha channel)
- **3D model image:** model snapshot (PNG, alpha channel in GLmol and ChemDoodle)
- **SDF file:** MDL Molfile from the 3D model (common molecules)
- PDB file: Protein Data Bank file from the 3D model (macromolecules)
- CIF file: Crystallographic Information File from the 3D model (crystal structures)

Information card

This collects and displays information about the current structural formula (loaded from the Sketcher) You can also find a database reference for the current 3D model.

Spectroscopy

This shows a new layer where you can view molecular spectra of the current structural formula (loaded from the Sketcher) More details are covered in the Spectroscopy chapter.

3D model resource

This redirects you to the web-page for the current 3D model on the website of its source database (except when the model is resolved using the Chemical Identifier Resolver)

Advanced search

These functions allow you to perform some advanced searches through the PubChem database using the structural formula from the sketcher.

- 1. **Similarity search:** search for compounds with a similar structural formula
- 2. **Substructure search:** search for compounds with the current structure as subset
- 3. **Superstructure search:** search for compounds with the current structure as superset

4. Spectroscopy

You can open the Spectroscopy view via **Tools > Spectroscopy**. You can view three kinds of molecular spectra.

- 1. Mass spectrum
- 2. IR spectrum
- 3. H1-NMR prediction

Export data

You can also export different kinds of data from the currently selected spectrum.

- 1. **PNG image:** snapshot from interactive spectrum
- 2. **JCAMP file:** JCAMP-DX file of the current spectrum

5. 3D model

The **Model** menu contains some general functions for the 3D model.

Reset

This function sets the model position, zoom and rotation back to default.

Representation

You can choose from a list of different molecule representations including; ball and stick, stick, van der Waals spheres, wireframe and lines. Macromolecules are automatically drawn using ribbons.

Background

You can switch between a black, gray or white background. The default background is black (exported images from GLmol or ChemDoodle have a transparent background)

Engines

You can choose from three different render engines: **GLmol**, **Jmol** and **ChemDoodle**. GLmol is used as default render engine. GLmol and ChemDoodle are based on WebGL, a browser technology to support 3D graphics. If WebGL is not available in your browser, Jmol wil be used for all rendering.

MolView automatically switches to:

- 1. **Jmol** if you execute functions from the Jmol menu
- 2. **GLmol** if you load macromolecules and WebGL is available (due to significant higher performance)
- 3. **ChemDoodle** if you load a crystal structure and WebGL is available (GLmol cannot render crystal structures)

You might want to switch back to GLmol when you do no longer need Jmol or ChemDoodle since GLmol has a better performance.

Note that macromolecules are drawn slightly different in each engine. ChemDoodle provides the finest display. You should, however, avoid using ChemDoodle for very large macromolecules.

Model transformation

You can rotate, pan and zoom the 3D model. Use the right button for rotation, the middle button for translation (except for ChemDoodle) and the scrollwheel for zooming. On touch devices, you can rotate the model with one finger and scale the model using two fingers.

Crystallography

You can load an array of crystal cells (2x2x2 or 1x3x3) or a single unit cell when viewing crystal structures.

Fog and clipping

When you are viewing large structures, like proteins, it can be useful to hide a certain part using fog or a clipping plane. GLmol offers a few options to do this.

- Fog: you can move the fog forward by dragging the mouse up while holding CTRL + SHIFT (drag in the opposite direction to move the fog backward)
- 2. **Clipping plane:** you can move a frontal clipping plane into the structure by dragging the mouse to the **left** while holding **CTRL** + **SHIFT** (drag in the opposite direction to move the clipping plane back)

6. Protein display

The **Protein** menu offers a number of protein display settings including different color schemes and different chain representations.

Show bio assembly

When loading a protein structure, MolView shows the asymmetric unit by default. This function allows you to view the full biological unit instead.

Chain representation

You can choose from four different chain representations. You can also view the full chain structure by enabling the **Bonds** option.

- 1. **Ribbon:** draws ribbon diagram (default representation)
- 2. **Cylinder and plate:** solid cylinders for α -helices and solid plates for β -sheets
- 3. **B-factor tube:** tube were thickness is calculated from the B-factor *(thermal motion)*
- 4. **C-alpha trace:** lines between central carbon atom in amino-acids (*very fast rendering*)

Chain coloring

You can choose from six chain color schemes.

- 1. **Secondary structure:** different colors for α -helices, β -sheets, etc.
- 2. **Spectrum:** color spectrum (*rainbow*)
- 3. **Chain:** each chains gets a different color
- 4. **Residue:** all amino-acid residues are colored differently
- 5. Polarity: colors polar amino-acids red and non polar amino-acids white
- 6. **B-factor:** blue for low B-factor and red for high B-factor (if provided)

7. Advanced Jmol tools

The **Jmol** menu offers some awesome Jmol-only functions and calculations.

Clear

Clears all executed calculations and measurements.

High Quality

Enables High Quality rendering in Jmol (enabled by default on fast devices) When turned off, anti-aliasing is disabled and the model is drawn using lines while transforming it.

Calculations

You can perform the following Jmol calculations in Jmol:

- MEP surface lucent/opaque: calculates and projects molecular electrostatic potential on a translucent or opaque on van der Waals surface
- Charge: calculates and draws the atomic charge as labels
- Bond dipoles: calculates and draws the individual bond dipoles
- Overall dipole: calculates and draws the net bond dipole
- **Energy minimization:** executes an interactive MMFF94 energy minimization (note that this function only executes a maximum of 100 minimization steps at a time)

Measurement

You can measure distance, angle and torsion using Jmol. You can activate and deactivate one of these measurement types via the Jmol menu.

- **Distance:** distance between two atoms in nm
- Angle: angle between two bonds in degrees
- **Torsion:** torsion between four atoms in degrees

Note that the resolved 3D model is only an approach of the real molecule, this means you have to execute an **Energy minimization** in order to do reliable measurements.