

Proof of Master Theorem

Geoffrey Matthews

April 17, 2018

1 Divide and Conquer Recurrence

Suppose that $a \geq 1$ and $b \geq 1$ are constants, let $f(n)$ be a function, and let $T(n)$ be defined on natural numbers with the recurrence:

$$T(n) = aT\left(\frac{n}{b}\right) + f(n) \quad (1)$$

Let's assume n is a power of b , so that

$$n = b^k \quad (2)$$

$$k = \log_b(n) \quad (3)$$

Now let's see what we get by trying to spell out terms and then cancel:

$$T(b^k) = aT(b^{k-1}) + f(b^k) \quad (4)$$

$$aT(b^{k-1}) = a^2T(b^{k-2}) + af(b^{k-1}) \quad (5)$$

$$a^2T(b^{k-2}) = a^3T(b^{k-3}) + a^2f(b^{k-2}) \quad (6)$$

$$\dots \quad (7)$$

$$a^{k-1}T(b^1) = a^kT(1) + a^{k-1}f(b) \quad (8)$$

Adding up both sides while cancelling terms, we get

$$T(b^k) = a^k T(1) + \sum_{i=1}^k a^{k-i} f(b^i) \quad (9)$$

$$= a^{\log_b(n)} T(1) + \sum_{i=1}^{\log_b(n)} a^{k-i} f(b^i) \quad (10)$$

$$= n^{\log_b(a)} T(1) + \sum_{i=1}^{\log_b(n)} a^{k-i} f(b^i) \quad (11)$$

Since $T(1)$ will be a constant, the complexity of the whole thing depends on whether

$$n^{\log_b(a)} \quad (12)$$

grows faster or slower than

$$\sum_{i=1}^{\log_b(n)} a^{k-i} f(b^i) \quad (13)$$

Note that expression 13 can be rewritten as

$$\sum_{i=1}^{\log_b(n)} a^{k-i} f(b^i) = \sum_{i=0}^{\log_b(n)-1} a^i f(b^{k-i}) \quad (14)$$

$$= \sum_{i=0}^{\log_b(n)-1} a^i f\left(\frac{b^k}{b^i}\right) \quad (15)$$

$$= \sum_{i=0}^{\log_b(n)-1} a^i f\left(\frac{n}{b^i}\right) \quad (16)$$

Let's see what we can find out about this expression, using either 13 or 16, and how it compares to 12.

$$\mathbf{2} \quad f(n) = \Theta(n^{\log_b(a)})$$

Let's first suppose

$$f(n) = \Theta(n^{\log_b(a)}) \tag{17}$$

then

$$f(b^i) = \Theta((b^i)^{\log_b(a)}) \tag{18}$$

$$= \Theta(b^{i \log_b(a)}) \tag{19}$$

$$= \Theta\left(\left(b^{\log_b(a)}\right)^i\right) \tag{20}$$

$$= \Theta(a^i) \tag{21}$$

Plugging this back in our expression 13, gives

$$\sum_{i=1}^{\log_b(n)} a^{k-i} f(b^i) = \sum_{i=1}^{\log_b(n)} a^{k-i} \Theta(a^i) \tag{22}$$

$$= \sum_{i=1}^{\log_b(n)} \Theta(a^k) \tag{23}$$

$$= \sum_{i=1}^{\log_b(n)} \Theta(a^{\log_b(n)}) \tag{24}$$

$$= \sum_{i=1}^{\log_b(n)} \Theta(n^{\log_b(a)}) \tag{25}$$

$$= (\log_b(n)) \Theta(n^{\log_b(a)}) \tag{26}$$

$$= \Theta(\lg(n) n^{\log_b(a)}) \tag{27}$$

Recall the expression for the complexity of $T(n) = T(b^k)$ in equation 11:

$$T(b^k) = n^{\log_b(a)} T(1) + \sum_{i=1}^{\log_b(n)} a^{k-i} f(b^i) \tag{11, revisited}$$

Putting 27 into 11, the complexity of $T(n)$, shows that if

$$f(n) = \Theta \left(n^{\log_b(a)} \right) \quad (17, \text{ revisited})$$

then

$$T(n) = \Theta \left(\lg(n) n^{\log_b(a)} \right) \quad (28)$$

$$\mathbf{3} \quad f(n) = O\left(n^{\log_b(a)-\epsilon}\right)$$

Now let's suppose that $f(n)$ has some slightly lower complexity. Suppose $0 < \epsilon < \log_b(a)$ is some (small) positive real number, and

$$f(n) = O\left(n^{\log_b(a)-\epsilon}\right) \tag{29}$$

then

$$f\left(\frac{n}{b^i}\right) = O\left(\left(\frac{n}{b^i}\right)^{\log_b(a)-\epsilon}\right) \tag{30}$$

$$= O\left(\frac{n^{\log_b(a)-\epsilon}}{b^{i \log_b(a)-i\epsilon}}\right) \tag{31}$$

$$= O\left(\left(\frac{n^{\log_b(a)}}{n^\epsilon}\right)\left(\frac{b^{i\epsilon}}{b^{i \log_b(a)}}\right)\right) \tag{32}$$

$$= O\left(\frac{b^\epsilon}{b^{\log_b(a)}}\right)^i \tag{33}$$

$$= O\left(\frac{b^\epsilon}{a}\right)^i \tag{34}$$

Plugging this back into expression 16 gives a geometric series which can be solved:

$$\sum_{i=0}^{\log_b(n)-1} a^i f\left(\frac{n}{b^i}\right) = O\left(\sum_{i=0}^{\log_b(n)-1} a^i \left(\frac{b^\epsilon}{a}\right)^i\right) \quad (35)$$

$$= O\left(\sum_{i=0}^{\log_b(n)-1} (b^\epsilon)^i\right) \quad (36)$$

$$= O\left(\frac{(b^\epsilon)^{\log_b(n)} - 1}{b^\epsilon - 1}\right) \quad (37)$$

$$= O\left(\frac{(n)^{\log_b(b^\epsilon)} - 1}{b^\epsilon - 1}\right) \quad (38)$$

$$= O\left(\frac{n^\epsilon - 1}{b^\epsilon - 1}\right) \quad (39)$$

$$= O(n^\epsilon) \quad (40)$$

$$= O\left(n^{\log_b(a)}\right) \quad (41)$$

Since b and ϵ are constants, and $0 < \epsilon < \log_b(a)$.

Recall again the expression for the complexity of $T(n) = T(b^k)$ in equation 11:

$$T(b^k) = n^{\log_b(a)} T(1) + \sum_{i=1}^{\log_b(n)} a^{k-i} f(b^i) \quad (11, \text{revisited})$$

Plugging 41 into 11 shows that if

$$f(n) = O\left(n^{\log_b(a)-\epsilon}\right) \quad (29, \text{revisited})$$

then

$$T(n) = \Theta\left(n^{\log_b(a)}\right) \quad (42)$$

$$4 \quad f(n) = \Omega \left(n^{\log_b(a)+\epsilon} \right)$$

We leave it as an exercise to show that if f has some slightly higher complexity,

$$f(n) = \Omega \left(n^{\log_b(a)+\epsilon} \right) \tag{43}$$

and $af\left(\frac{n}{b}\right) \leq cf(n)$ for some c and large enough n , then

$$T(n) = \Theta(f(n)) \tag{44}$$