

RESEARCH ARTICLE

Distinct, direct and climate-mediated environmental controls on global particulate and mineral-associated organic carbon storage

Paige M. Hansen^{1,2}  | Rebecca Even¹  | Alison E. King¹  | Jocelyn Lavalée^{1,2,3}  |
Meagan Schipanski^{1,2}  | M. Francesca Cotrufo^{1,2} 

¹Department of Soil and Crop Sciences, Colorado State University, Fort Collins, Colorado, USA

²Graduate Degree Program in Ecology, Colorado State University, Fort Collins, Colorado, USA

³Environmental Defense Fund, New York, New York, USA

Correspondence

Paige M. Hansen, Department of Soil and Crop Sciences, Colorado State University, Fort Collins, CO, USA.

Email: paige.m.hansen@colostate.edu

Funding information

National Science Foundation Division of Environmental Biology, Grant/Award Number: 2016003

Abstract

Identifying controls on soil organic carbon (SOC) storage, and where SOC is most vulnerable to loss, are essential to managing soils for both climate change mitigation and global food security. However, we currently lack a comprehensive understanding of the global drivers of SOC storage, especially with regards to particulate (POC) and mineral-associated organic carbon (MAOC). To better understand hierarchical controls on POC and MAOC, we applied path analyses to SOC fractions, climate (i.e., mean annual temperature [MAT] and mean annual precipitation minus potential evapotranspiration [MAP-PET]), carbon (C) input (i.e., net primary production [NPP]), and soil property data synthesized from 72 published studies, along with data we generated from the National Ecological Observatory Network soil pits ($n=901$ total observations). To assess the utility of investigating POC and MAOC separately in understanding SOC storage controls, we then compared these results with another path analysis predicting bulk SOC storage. We found that POC storage is negatively related to MAT and soil pH, while MAOC storage is positively related to NPP and MAP-PET, but negatively related to soil % sand. Our path analysis predicting bulk SOC revealed similar trends but explained less variation in C storage than our POC and MAOC analyses. Given that temperature and pH impose constraints on microbial decomposition, this indicates that POC is primarily controlled by SOC loss processes. In contrast, strong relationships with variables related to plant productivity constraints, moisture, and mineral surface availability for sorption indicate that MAOC is primarily controlled by climate-driven variations in C inputs to the soil, as well as C stabilization mechanisms. Altogether, these results demonstrate that global POC and MAOC storage are controlled by separate environmental variables, further justifying the need to quantify and model these C fractions separately to assess and forecast the responses of SOC storage to global change.

KEYWORDS

climate, environmental controls, primary productivity, soil organic carbon fractions, soil properties

1 | INTRODUCTION

Maintaining and increasing soil organic carbon (SOC) stores is crucial to mitigating climate change and to enhancing the soil's capacity to provide essential ecosystem services, including nutrient recycling and maintenance of plant productivity. As such, it is imperative that we develop an advanced understanding of the environmental conditions under which SOC is most vulnerable to loss, as well as its responses to deliberate management practices aimed at increasing SOC storage. Although we can now identify how drivers of SOC storage, including climate (Jobbágy & Jackson, 2000), carbon (C) inputs, and soil properties (Hassink, 1997; Six et al., 2002) operate on a variety of spatial scales (Wiesmeier et al., 2019), we lack a quantitative framework of hierarchical controls on SOC storage that incorporates its conceptualization into separate fractions. Such knowledge could give us a more specific understanding of how different forms of SOC can be gained or can be lost across multiple ecosystem types, allowing us to better target “multi-pool management” practices (Angst et al., 2023) towards increasing or maintaining SOC storage in different fractions.

Conceptualizing SOC into particulate (POC) and mineral-associated organic carbon (MAOC) fractions can be particularly useful as a first step in building a more comprehensive understanding of global controls on SOC formation and persistence (Cotrufo & Lavelle, 2022; Lavelle et al., 2020). Particulate organic C is formed via fragmentation of structural plant inputs and is primarily protected by physical occlusion in aggregates (or in the case of pyrogenic C, through chemical recalcitrance). Limited protection from decomposition results in POC having relatively short mean residence times (von Lützow et al., 2007), except in locations where decomposition is limited by physical or physiological constraints on microbial activity (Cotrufo & Lavelle, 2022) where POC is found to accumulate (Herndon et al., 2017). In contrast, MAOC is formed via sorption of microbial necromass and decomposition products, as well as soluble plant inputs, to soil mineral surfaces (Haddix et al., 2016; Kallenbach et al., 2016; Liang et al., 2019). These mineral bonds make MAOC relatively inaccessible to microbial decomposition, leading to, on average, longer mean residence times than POC (von Lützow et al., 2007). Given their distinct mechanisms of formation and persistence, it is likely that global POC and MAOC storage are controlled by contrasting environmental variables.

Climate may be the primary driver of both POC and MAOC storage because of its effects on factors that influence the amount of C that enters (i.e., C inputs) and leaves the soil (i.e., C outputs; Cotrufo et al., 2021). For instance, both temperature and moisture are important controls on annual C inputs to soil (i.e., net primary production [NPP]; Churkina & Running, 1998), with increased C inputs often leading to greater SOC storage (Lajtha et al., 2014; Luo et al., 2017). While we lack an understanding of how C inputs separately affect POC and MAOC storage, MAOC can be expected to be more reflective of C inputs than POC, due to POC's lack of soil matrix protection from decomposition. Independent of inputs, climate may act as an additional control on SOC storage through its effects on microbial

activity constraints (Cotrufo & Lavelle, 2022). For instance, POC may be particularly sensitive to climate-driven variations in decomposition (Conant et al., 2011). Temperature and moisture are key controls on microbial activity, with higher temperatures leading to accelerated decomposition (Schimel, 2018), especially for complex structural compounds that require higher activation energy to decompose (Davidson & Janssens, 2006) and when optimum soil moisture levels increase microbial access to substrates through increased diffusion (Schimel, 2018). As such, we may expect lower POC storage in warm compared to cold ecosystems, especially when microbial access to substrates is limited by low diffusion under dry conditions. Additionally, moisture may serve as a direct control on MAOC storage through its effects on dissolved organic carbon (DOC) leaching. As plant-derived DOC can make up nearly half of MAOC storage (Angst et al., 2021), leaching of DOC due to high moisture may contribute to greater MAOC stores (Haddix et al., 2020), especially in ecosystems where the majority of MAOC is plant-derived. However, little work has focused on how temperature, moisture, and C inputs interact with one another to control POC and MAOC storage, especially at large spatial scales.

In addition to climate, soil physicochemical properties may act as a further control on POC and MAOC storage. For instance, strong associations with soil mineral surfaces may mean that the capacity of the soil to form mineral bonds with organic C molecules is an additional control on MAOC storage, with higher silt and clay contents leading to greater storage of MAOC (Hassink, 1997; Six et al., 2002). Other physicochemical properties, including exchangeable calcium (Ca) and iron (Fe) and aluminum (Al) hydroxides, may operate as additional controls on MAOC storage (King et al., 2023; Kirsten et al., 2021; Rowley et al., 2021), whose importance to C storage may be mediated by soil moisture and pH (Rasmussen et al., 2018). pH may also control POC storage through its inhibitory effects on microbial activity. Like cold temperatures, low pH can slow microbial decomposition (Rousk et al., 2009) and may contribute to greater POC storage in acidic soils. However, little work has investigated the extent to which these soil properties interact with our hypothesized overarching climatic controls in determining global patterns of POC and MAOC storage.

To identify hierarchical controls on global C storage, we synthesized POC and MAOC fraction data from 72 existing studies and databases, along with SOC fraction data we generated from the National Ecological Observatory Network (NEON) “megapit” soils (Hinckley et al., 2016). We included studies conducted in a diversity of climate and land cover types, with the goal of identifying controls on POC and MAOC that are generalizable across a variety of ecosystems. We aimed to assess the extent to which climate, NPP, and soil properties directly versus indirectly control C storage in POC, MAOC, and bulk SOC. Based on the observations described above, we broadly hypothesized that POC and MAOC storage would be governed by distinct, climate-mediated environmental controls (Cotrufo et al., 2021), with POC being primarily controlled by C loss processes related to microbial activity constraints, and MAOC being primarily controlled by a combination

of C input limitations and soil properties related to C stabilization. Additionally, we hypothesized that linking environmental variables to POC and MAOC would exhibit greater utility in understanding global patterns of C storage than linking the same variables to bulk SOC alone.

2 | MATERIALS AND METHODS

2.1 | Global synthesis of particulate and MAOC data

We constructed an observational dataset of POC and MAOC by conducting a comprehensive search on Google Scholar using various combinations of keywords including “soil carbon,” “soil organic matter,” “particulate and mineral-associated,” “light and heavy fractions,” “density fractionation,” and “size fractionation,” then collating all relevant data from published studies and existing databases (Figure 1). We chose to use Google Scholar as it may have greater geographic representation than other search engines (Harzing & Alakangas, 2016; Martín-Martín et al., 2018). For studies that fractionated soil by size or density only, we defined POC as $>53\text{--}63\mu\text{m}$ in size or $<1.65\text{--}1.85\text{g cm}^{-3}$ in density, and MAOC as $<53\text{--}63\mu\text{m}$ or $>1.65\text{--}1.85\text{g cm}^{-3}$. For studies that fractionated soil by both size and density, we considered POC to be $<1.65\text{--}1.85\text{g cm}^{-3}$, and MAOC

to be $>1.65\text{--}1.85\text{g cm}^{-3}$ and $<53\text{--}63\mu\text{m}$. Heavy coarse fraction ($>1.65\text{--}1.85\text{g cm}^{-3}$ and $>53\text{--}63\mu\text{m}$; sensu Leuthold et al., 2022) C and occluded SOC fraction (in the cases in which POC was separated by density before dispersion) data were retained for calculations of total SOC (see Section 2.4, below), but excluded from analyses performed on fraction data only. This was done to maintain as much consistency in SOC fraction definitions as possible, as the fractionation schemes of the studies included in our synthesis varied with both study and ecosystem type. While we focused our synthesis primarily on observational studies, we also included data from control plots (i.e., not receiving experimental treatments) of relevant field studies involving experimental N deposition, CO_2 enrichment, climate manipulations, etc. However, for experimental studies in agricultural systems where treatment represented a modified management practice, we included all reported data to account for the fact that management practices vary across the landscape, and may generate different environmental controls (e.g., NPP, pH, etc.) on POC and MAOC storage.

2.2 | NEON data generation

We supplemented our synthesis with additional SOC fraction data we generated by fractionating “megapit” mineral soil samples from the NEON (Hinckley et al., 2016). Soil samples were taken from megapits




FIGURE 1 Geographic location of all data points included in our synthesis ($n=901$). Different colors represent the land cover type assigned to each point based on author-reported plant community composition and management information. For croplands, $n=312$; for forests, $n=233$; for grasslands, $n=301$; for shrublands, $n=35$; for tundra, $n=7$; and for wetlands, $n=10$. The distribution of our included data across Köppen-Geiger Climate Classification Zones (Beck et al., 2018) can be found in Figure S1. Map lines delineate study areas and do not necessarily depict accepted national boundaries.

or large, temporary soil pits dug to either 2 m depth or bedrock at each of NEON's 47 terrestrial research sites, divided by horizon, then homogenized and characterized according to standard NEON protocols. A subsample of each of the 2-mm sieved, air-dried megapit soil samples was shipped to Colorado State University, where they were de-quarantined for 16 h at 116°C according to APHIS regulation. After de-quarantining, we fractionated all soil samples by both density (1.85 g cm^{-3}) and size ($53 \mu\text{m}$) after mechanical dispersion as described in Zhang et al., 2021. Briefly, 5.5–6.0 g subsamples were first shaken in deionized (DI) water for 15 min, then centrifuged at 3400 rpm. Dissolved organic matter was decanted over a $20 \mu\text{m}$ nylon filter and stored at -20°C . Any particulate material on the filter was set aside as part of the light particulate organic matter fraction ($<1.85 \text{ g cm}^{-3}$). Twelve glass beads and 1.85 g cm^{-3} sodium polytungstate (SPT) were added to the soil residue and shaken for 18 h to disperse aggregates. Soils were then centrifuged for 30 min at 3400 rpm, and the remaining light particulate organic matter was aspirated onto a $20 \mu\text{m}$ nylon filter. The pellet ($>1.85 \text{ g cm}^{-3}$) was rinsed multiple times to remove any remaining SPT before wet-sieving to separate it into heavy coarse organic matter ($>53 \mu\text{m}$), and mineral-associated organic matter ($<53 \mu\text{m}$). All solid fractions were oven-dried at 60°C , then finely ground to ensure homogenization before being analyzed for %C on an elemental analyzer (Costech ECS 4010; Valencia, CA, USA). Samples that contained carbonates according to the NEON database were acid-fumigated to remove carbonates (Harris et al., 2001), then re-analyzed to obtain organic C values. For this study, we used the light POC and MAOC values from topsoil samples only (see Section 2.4, below), for a total of 47 observations for each of the fractions, as well as bulk SOC.

2.3 | Auxiliary data incorporation

In addition to SOC fraction data, we incorporated auxiliary data including mean annual temperature (MAT; $^\circ\text{C}$), mean annual precipitation (MAP; mm), potential evapotranspiration (PET; mm), aboveground NPP ($\text{Mg Cha}^{-1}\text{year}^{-1}$) as a proxy for annual soil C inputs, soil texture and pH, bulk density, sampling depth, plant community composition, and any applicable management practices into our synthesized dataset. In all cases, we used the site data as reported in publications or contained within the NEON database. However, if specific site data types such as MAT, MAP, PET, and NPP were not available, we used reported geographic coordinates to extract equivalent 20-year average MAT and MAP values from WorldClim 2.0 (Fick & Hijmans, 2017), and 20-year average PET and NPP values from MODIS (data products MOD16A3GF.061 and MOD17A3HGF.006, respectively; Running et al., 2021; Running & Zhao, 2019). We subtracted PET from MAP to create a metric of effective moisture (MAP-PET; mm; sensu Kramer & Chadwick, 2018). Additionally, we assigned one of six broad land cover types to each soil profile (i.e., cropland, forest, grassland, shrubland, tundra, wetland; Figure 1) based on reported plant community composition and management information.

2.4 | Data processing

After compiling both synthesis and NEON megapit data, we used reported SOC stocks, soil depth, and bulk density values to convert all SOC fraction stocks to SOC fraction concentrations ($\text{g fraction-C kg soil}^{-1}$). As mentioned above, we did not include organic C in heavy coarse organic matter or in occluded SOC fractions in our downstream analyses of fraction data. However, we retained and used any heavy coarse organic matter or occluded SOC fraction data in calculations of g SOC kg soil^{-1} (i.e., sum of all fraction C values), as well as calculations of MAOC relative to bulk SOC storage (i.e., f_{MAOC} ; g MAOC kg soil^{-1} / g SOC kg soil^{-1}). While our original intent was to test for variations in SOC storage drivers throughout the soil profile, there was not enough subsoil data available at the global scale to include subsoils in our downstream statistical analyses. As such, we chose to focus our analysis specifically on topsoil. We considered the first sampled depth increment or horizon reported in each study to be representative of topsoil, and applied data from that layer only in downstream analyses. Depending on study site and sampling design, topsoils included both O and A horizons, and spanned different soil depths (Figure S2). This was done to minimize biases associated with variation in maximum topsoil sample depth across our synthesized dataset, as methods of normalizing data to a standardized depth (e.g., Abdalla et al., 2018; Hou et al., 2020; Ogle et al., 2005) across all our data points diluted the effects of driving variables known to change along the soil profile less than, or differently from, SOC (data not shown). After filtering our data so that it contained only topsoil, we averaged all data collected at the plot level such that we had one data point per site. At the end of all data processing, our final dataset included 901 total data points (i.e., $n=901$ for each SOC fraction as well as bulk SOC) from 72 studies (Figure 1; Figure S1; Table S1; see Data Sources, below).

2.5 | Statistical analyses

We first assessed broad relationships and patterns among our data using simple linear regressions. Specifically, we used linear regressions to test for relationships of MAT and MAP-PET with NPP, as well as relationships of MAT, MAP-PET, NPP, soil pH, and % sand with POC, MAOC, bulk SOC, and f_{MAOC} . We did not include multiple factors or any interactive effects in these simple regressions, such that we tested for the effects of only one explanatory variable per regression. We then used results from these analyses and literature-supported hypotheses to conduct path analyses investigating direct and indirect controls on C storage in POC, MAOC, and SOC using the R package 'lavaan' (Rosseel, 2012). Given that many known drivers of soil C storage interact with one another, we chose to use path analyses because of their ability to test for mediation, as well as direct and indirect effects of variables on outcomes of interest (Shipley, 2016). We created two separate path analyses, one which predicted POC and MAOC storage and one which predicted storage in bulk SOC. The fit of different iterations of these path analyses was assessed

using Chi square, the comparative fit index ($CFI > .9$), the root mean square error of approximation ($RMSEA < .08$), and the standardized root mean square residual ($SRMR < .08$; Hooper et al., 2008). Our final path analysis structures that exhibited the best fits allowed for direct effects of MAT, MAP-PET, NPP, pH, and % sand, as well as indirect effects of MAT and MAP-PET via NPP, on POC and MAOC storage, or bulk SOC storage. We originally included topsoil sampling depth as a covariate in our analyses to account for differences in it within and across studies but removed it due to poor fit. We assessed effects of our environmental variables (i.e., MAT, MAP-PET, NPP, pH, and % sand) by assigning coefficients to individual pathways, then multiplying coefficients to calculate indirect effects (e.g., for indirect effects of MAP-PET, we multiplied the effect of MAP-PET on NPP by the effect of NPP on MAOC). We summed direct and indirect path coefficients to estimate total effects of each variable on POC, MAOC and bulk SOC storage. In addition to determining controls on SOC storage, we compared our POC and MAOC fraction path analysis with our bulk SOC analysis to assess the utility of separating SOC into POC and MAOC in understanding global patterns of SOC formation and persistence. We also assessed differences in f_{MAOC} among land cover types using Welch's one-way analysis of variance to account for uneven sample size among land cover types, and further investigated our significant test result using Games-Howell post-hoc pairwise comparisons using the R package 'rstatix' (Kassambra, 2023). All analyses were carried out in R version 4.1.1 (R Core Team, 2021).

3 | RESULTS

3.1 | Broad relationships between climate, NPP, soil properties, and POC, MAOC, and bulk SOC storage

We first explored global patterns of SOC formation and persistence by conducting linear regressions between environmental variables and POC, MAOC, and SOC storage, as well as relationships between climate and NPP. We found significant, positive relationships between NPP and both MAT and MAP-PET (Figure S3a,b; Table S2), reflecting expected associations between climate and NPP. We also found significant relationships between POC and MAT, MAP-PET, NPP, and pH. MAP-PET and NPP were positively associated with POC, while MAT and pH were negatively associated with POC storage (Figure S3c–g; Table S2). Environmental controls on MAOC included MAP-PET, NPP, and % sand, with MAP-PET and NPP being positively related, and % sand being negatively related to global MAOC storage (Figure S3h–i; Table S2). All studied environmental variables were associated with bulk SOC. Specifically, MAP-PET and NPP were positively associated with SOC, while MAT, pH, and % sand were negatively associated with bulk SOC storage (Figure S3m–q; Table S2). We note that the significance of the above relationships may be driven in part by high sample size, as many relationships had relatively small r^2 values (i.e., .0111–.221).

3.2 | Direct and indirect controls on global POC, MAOC, and bulk SOC storage

To assess direct and indirect effects of the above environmental variables on soil C, we constructed separate path analyses to predict C storage in POC and MAOC and bulk SOC (Figures 2 and 3). Our fraction path analysis fit the data well ($\chi^2_{(2,n=649)} = 1.239, p = .538$; $CFI = 1.0$; $RMSEA = .068$; $SRMR = .007$; Hooper et al., 2008), and explained 35.8% of the variation in NPP, 12.3% of the variation in POC, and 20.5% of the variation in MAOC (Figure 2; Table S3). Consistent with our linear regression analyses, NPP was positively predicted by both MAT and MAP-PET. Global POC storage was directly predicted by MAT and pH, both of which had a negative effect on POC. Contrary to our linear regressions, however, neither MAP-PET nor NPP emerged as a significant predictor of POC, potentially because path analysis calculates partial regression coefficients that account for correlations and covariance between multiple factors, which likely reduced the already weak relevance (i.e., low r^2) of our linear regression coefficients. Global MAOC storage had direct, positive relationships with MAP-PET and NPP (and as such, indirect relationships with MAT and MAP-PET), as well as a direct, negative relationship with % sand (Figure 2; Table S3).

Our SOC path analysis revealed similar trends in global C storage. Like our fraction analysis, our SOC path analysis fit our data well ($\chi^2_{(2,n=649)} = 1.239, p = .538$; $CFI = 1.0$; $RMSEA = .0$; $SRMR = .007$; Hooper et al., 2008) and explained 35.8% of the variation in NPP, with both MAT and MAP-PET once again having direct, positive effects on NPP (Figure 3; Table S4). However, this path analysis explained only 12.3% of the variation in global SOC storage, which was predicted by all included exogenous variables. MAT, pH, and % sand had direct, negative effects on SOC storage, while MAP-PET and NPP had direct, positive effects on SOC. In addition, MAT and MAP-PET had positive, indirect effects on SOC via NPP (Figure 3; Table S4).

3.3 | Relationships between f_{MAOC} , environmental variables, and land cover type

We next aimed to understand the extent to which the above environmental controls are reflected in the fraction of C stored in MAOC relative to total bulk SOC (i.e., f_{MAOC}), as well as across land cover types that are representative of those controls. MAT, NPP, and soil pH were positively related to f_{MAOC} while soil % sand was negatively related (Table S5; Figure S4), though these relationships were relatively weak (i.e., in all cases, $r^2 < .1$). We did not find any relationships between MAP-PET and f_{MAOC} (Table S5; Figure S4). Additionally, land cover was a significant predictor of f_{MAOC} ($F_{5,37,901} = 58.024$; $p < 2e-16$; Figure 4), with croplands having higher f_{MAOC} than all other land cover types (Table S6). In addition, both shrublands and grasslands had higher f_{MAOC} than forests, wetlands, and tundra (Table S6). Notably, most land cover types spanned




FIGURE 2 Results from path analysis showing direct and indirect effects of mean annual temperature (MAT; °C), mean annual precipitation minus potential evapotranspiration (MAP-PET; mm), net primary production (NPP; $\text{Mg C ha}^{-1} \text{ year}^{-1}$), soil pH, and % sand on global particulate (POC; g C kg soil^{-1}) and mineral-associated organic carbon (MAOC; g C kg soil^{-1}) storage (a). Green and red arrows indicate significant positive and negative effects, respectively, of the variables described above on POC and MAOC storage. Grey, dashed arrows indicate non-significant paths. The widths of the arrows correspond to standardized path coefficients, shown in numbers above each arrow. Standardized total (i.e., direct plus indirect) effects of the variables described above on POC and MAOC storage are shown in (b) and (c). In all cases, n.s. indicates non-significant, $***p < .001$. Full results and output are provided in [Table S3](#).




FIGURE 3 Results from our path analysis showing direct and indirect effects of mean annual temperature (MAT; °C), mean annual precipitation minus potential evapotranspiration (MAP-PET; mm), net primary production (NPP; $\text{tons C ha}^{-1} \text{ year}^{-1}$), soil pH, and % sand on global bulk soil organic carbon (SOC; g C kg soil^{-1}) storage (a). Green and red arrows indicate significant positive and negative effects of the variables described above on bulk SOC storage. The widths of the arrows correspond to standardized path coefficients, shown in numbers above each arrow. Standardized total (i.e., direct plus indirect) effects of the variables described above on bulk SOC storage are shown in (b). In all cases, n.s. indicates non-significant, $*p < .05$, $**p < .01$, and $***p < .001$. Full results and output are provided in [Table S4](#).




FIGURE 4 Comparison of the fraction of C stored in mineral-associated organic carbon (MAOC) relative to bulk soil organic carbon (SOC), represented as f_{MAOC} (%; (g MAOC kg soil⁻¹/g SOC kg soil⁻¹) × 100), across land cover types (cropland $n=312$; forest $n=233$; grassland $n=301$; shrubland $n=35$; tundra $n=7$; wetland $n=10$). Solid black dots inside each box indicate group means, and letters above bars indicate significant differences between groups. Full post-hoc testing outputs are provided in Table S6. Additional information on the distribution of our mean annual temperature, mean annual precipitation minus potential evapotranspiration, net primary production, soil pH, and % sand data across land cover types is provided in Figure S5 and Table S7.

large ranges of f_{MAOC} , which declined from >75% in croplands to <50% in tundra (Figure 4).

4 | DISCUSSION

In this study, we sought to identify hierarchical controls on global POC, MAOC, and bulk SOC storage by applying path analyses to a large dataset synthesized from published papers and data we generated from NEON megapit soils. We also evaluated the utility of C fractionation in understanding global SOC storage patterns by comparing our POC and MAOC fraction path analysis to our bulk SOC analysis. Overall, climate and soil properties (i.e., pH and texture) emerged as the primary controls on SOC, with climate exerting both direct as well as indirect controls on SOC through controlling plant productivity. Separating SOC into POC and MAOC improved our ability to identify a hierarchy of controls on SOC storage, as demonstrated by identification of distinct environmental drivers of POC and MAOC storage, along with greater % variation explained in our SOC fraction path analysis compared to our bulk SOC analysis. We found that MAT and soil pH were dominant controls on POC storage, while MAP-PET, NPP, and % sand were dominant controls on MAOC storage. These results are echoed in our bulk SOC path analysis, which revealed a similar structure of controls. These results were also echoed in our analysis of f_{MAOC} across land cover types, with land cover types that tend to experience weaker constraints on SOC decomposition (i.e., croplands) having a greater fraction of SOC

stored in MAOC than types that experience stronger constraints (i.e., wetlands and tundra). Altogether, these results demonstrate that POC and MAOC are controlled by separate environmental variables, with POC storage being primarily controlled by decomposition (i.e., C output) limitations, and MAOC being primarily controlled by climate-driven plant productivity (i.e., C input) limitations, as well as SOC stabilization.

Consistent with our original hypotheses, we found that global POC storage is controlled primarily by factors that impose limitations on C loss processes (Figure 3; Table S3). The two variables most closely related to POC storage, temperature and pH, are strong controls on microbial activity, with both lower pH (Rousk et al., 2009) and lower temperatures (Schimel, 2018) contributing to slower decomposition. These findings are supported by studies conducted at smaller spatial scales, with low pH contributing to greater POC storage across the United States (Yu et al., 2022) and Europe (Lugato et al., 2021), as well as increased temperatures leading to greater POC compared to MAOC losses in lab incubations (Benbi et al., 2014). Lower POC storage under conditions that promote microbial decomposition is consistent with this SOC fraction's turnover time (von Lützow et al., 2007) and sensitivity to disturbance (e.g., Lobe et al., 2011; Poeplau et al., 2017; Song et al., 2014; Thaysen et al., 2017; Wu et al., 2023), whereby little to no protection from microbial attack makes POC reflective of microbial physical and physiological access constraints (Cotrufo & Lavelle, 2022). This shows that POC may be most vulnerable to C losses from the soil, despite being formed primarily from structural plant inputs, and therefore represents the SOC fraction that requires protection from climate change and management disturbance, such as tillage or draining of wetlands (Ashagrie et al., 2007; Bouajila & Tahar, 2010; Lavelle et al., 2019).

We found that global MAOC storage is controlled by a different set of environmental controls from POC, including % sand, NPP, and MAP-PET (Figure 3; Table S3). This indicates that MAOC is primarily controlled by climate-driven limitations on C inputs, moisture, and stabilization potential. Soil % sand was the largest control on MAOC. This relationship is broadly supported by the literature, with lower sand (and therefore greater silt and clay) contents contributing to greater MAOC storage (Hassink, 1997; Six et al., 2002). Our additional observed relationship between MAOC storage and MAP-PET is also consistent with previous findings, whereby MAP-PET was found to control the abundance of C retained by reactive minerals (Kramer & Chadwick, 2018). This association with effective moisture may be reflective of the direct sorption pathway of MAOC formation, with increased moisture leading to greater leaching of soluble plant inputs that are capable of sorbing directly to soil mineral surfaces (Haddix et al., 2020). Though large portions of the MAOC pool can be formed from microbial decomposition products and necromass (Huang et al., 2019; Kallenbach et al., 2016; Liang et al., 2019), this direct pathway of MAOC formation from plant compounds is increasingly recognized as a significant pathway of MAOC formation, especially in wet environments (Angst et al., 2021; Yu et al., 2022). Limitation on C inputs (i.e., NPP) represented another constraint

on global MAOC storage. We interpret this relationship between MAOC and plant C inputs as the result of MAOC's protection from microbial access, making it less vulnerable to types of disturbance that accelerate decomposition and therefore more representative of C inputs to the soil than POC. Additionally, there could be interactions between the soil matrix and plant productivity not explored in this study that contribute to greater MAOC storage. For instance, recent work indicates that increased soil mineral capacity index (MCI) leads to both greater plant productivity and MAOC storage (King et al., 2023). As Fe and Ca, key components of this MCI, are essential plant nutrients and important for mineral stabilization, factors that promote plant productivity may also promote MAOC stabilization and storage (King et al., 2023; but see Fuhrmann & Zuberer, 2021; Ramos et al., 2018). Similarly, interactions between increased silt and clay content, soil moisture, and plant productivity resulted in higher MAOC storage in dryland soils (Mao et al., *in review*).

To our knowledge, this analysis is one of the first to assess global, hierarchical controls acting on both POC and MAOC storage (though recent work has investigated the roles of climate and soil mineralogy in determining MAOC storage and saturation; Georgiou et al., 2022). However, the structure of controls we identified, particularly with regards to climate as an overarching driver of POC and MAOC storage, is generally consistent with studies conducted at smaller spatial scales. For instance, climate and C inputs exerted the greatest influence on total bulk SOC and POC:MAOC ratio, as well as change in bulk SOC storage, in Australian croplands (Luo et al., 2017). Increased moisture has also been associated with greater MAOC persistence at continental (Heckman et al., 2023) and global scales (Heckman et al., 2022), though in contrast to recent work on global patterns of MAOC saturation, we do not find that MAT is a significant control on global MAOC storage (Georgiou et al., 2022), potentially due to differences in data analysis or in distribution of data points among climate zones (Figure S1) between our studies. Our work moves beyond the above studies by expanding our understanding of SOC storage controls through explicit incorporation of both POC and MAOC, the former of which is relatively understudied in comparison to the latter, as well as by including a global distribution of climate, soil, and vegetation types.

In addition to identifying global controls on POC and MAOC storage, we compared our SOC fraction to our bulk SOC path analysis. While we were able to identify similar effects of climate, C inputs, and soil physicochemical properties on SOC storage when we considered SOC as a single pool, the effects of these variables on SOC were stronger in our fraction path analysis than in our bulk SOC analysis. Additionally, our bulk SOC path analysis explained less variation in SOC storage than our fraction path analysis (12.3% for bulk SOC vs. 20.5% for MAOC; Figures 3 and 4; Tables S3 and S4). This demonstrates the utility of separating SOC into distinct fractions to better understand controls on its formation and storage (Lavallee et al., 2020). In this study, separating SOC into POC and MAOC not only conferred greater explanatory power, but also revealed that these separate fractions are controlled by distinct environmental

variables—a finding that could not be realized when considering SOC as a single pool. Building upon the findings of previous syntheses that demonstrate that SOC is not created equal, and that its component fractions respond differently to global change and management (Heckman et al., 2022; Prairie et al., 2023; Rocci et al., 2021), this study indicates that separating SOC into POC and MAOC improves our ability to understand and quantify environmental controls on SOC storage at the global scale. As such, we continue to advocate that SOC is not studied as a unique pool. In particular, as the field has gained robust knowledge of SOC dynamics from its separation into POC and MAOC, we advocate for further separation of SOC into, for example, free versus occluded POC and exchangeable versus stable MAOC, as well as further analyses on how fractionation scheme may influence predictions of SOC fraction formation and persistence (e.g., Leuthold et al., *in review*; Poeplau et al., 2018). Doing so could advance our understanding of how processes like aggregate inclusion as well as sorption and desorption of DOC to mineral surfaces impact soil C storage, thus improving our ability to predict responses of SOC to global change.

Like our bulk SOC path analysis, our comparisons of SOC storage in MAOC relative to total bulk SOC (i.e., f_{MAOC}) across our included environmental variables and across land cover types dovetail our proposed understanding of POC as controlled by C loss processes. We found that land cover types thought to experience weaker constraints on microbial decomposition, such as croplands (e.g., due to the effects of tillage, Balesdent et al., 2000; Lupwayi et al., 2004 and optimal nutrient status, Parihar et al., 2019) had a greater proportion of their total SOC pool stored in MAOC than types that tend to experience stronger decomposition constraints, such as wetlands or tundra (i.e., due to anaerobic conditions (Huang et al., 2020) or cold temperatures (Frøseth & Bleken, 2015; Figure 4; Table S5). This is mirrored in our linear regressions between MAT, pH, and f_{MAOC} , whereby lower temperatures and lower pH confer greater proportions of SOC stored as POC (Figure S3; Table S5). These findings build upon Sokol et al., 2022 through more explicit incorporation of land cover types including wetlands and tundra, who also found that croplands and polar regions tend to have higher and lower f_{MAOC} , respectively, than both grasslands and forests. Given the large variance in f_{MAOC} within land cover types, especially in types that span a wide range of climatic conditions (e.g., grasslands and forests), this trend suggests that constraints on C inputs and losses from the soil may be a larger control on SOC fraction storage than C input quality. However, the large overlap in f_{MAOC} between forests and grasslands may also be explained by large differences in input quality within land cover types, especially for forests (Krishna & Mohan, 2017; Pérez-Harguindeguy et al., 2000). While more work is needed to confirm the effects of plant input quality on f_{MAOC} , these findings generally support the hypothesis that climate, and its effects on C inputs and decomposition-related SOC losses, represent overarching controls on SOC formation and persistence at the global scale (Cotrufo et al., 2021).

Moreover, within the context of ongoing calls to focus research efforts on building new, stable SOC to meet climate change

mitigation goals (Bradford et al., 2019; Rumpel et al., 2018; Vermeulen et al., 2019), our findings highlight the importance of not just accruing new SOC, but also preventing losses of existing SOC, particularly C that is stored in POC. Our analyses support the notion that POC is highly sensitive to loss (Lobe et al., 2011; Poeplau et al., 2017; Song et al., 2014; Thaysen et al., 2017; Wu et al., 2023), and demonstrate that ecosystems that are especially vulnerable to disturbances associated with global change (e.g., wetlands, tundra) harbor SOC stores that are mostly comprised of POC (e.g., Herndon et al., 2017; Mirabito & Chambers, 2023; Sousa et al., 2015; Xu et al., 2009). Given that C loss processes exert a much greater control over POC than C inputs, and that the soils with the highest C contents (e.g., wetlands and tundra) have high POC stores, large amounts of SOC may be very difficult to rebuild if lost, particularly under a warming climate. As such, a focus on building new SOC, especially in the form of MAOC, is not enough (*sensu* Angst et al., 2023)—we claim that developing incentives that maintain current SOC stores in natural lands is equally essential to preventing the negative effects of climate change.

Despite our above findings, we note that our path analyses explained only 12.3%, 20.5%, and 12.3% of the variation in POC, MAOC, and bulk SOC storage, respectively (Figure 3; Table S3), indicating that the climate, C input, and soil property variables we included in our analyses were not sufficient to describe global patterns of SOC storage. Several recent studies demonstrate that additional soil physicochemical properties not included in this study, especially exchangeable Ca and Fe- and Al-hydroxides, are better predictors of MAOC storage than % sand alone (King et al., 2023; Kirsten et al., 2021; Rowley et al., 2021). Microbial traits including mycorrhizal type (Craig et al., 2018; Horsch et al., 2023; Keller et al., 2021) and transformation efficiency may be additional modulators of both SOC fraction storage and f_{MAOC} , though the majority of studies on microbial transformations have focused primarily on their effects on microbial-derived MAOC (e.g., Craig et al., 2022; Ernakovich et al., 2021; Kallenbach et al., 2016; Liang et al., 2019). While satellite-based NPP data is often used as a proxy for C inputs to the soil (e.g., Chen et al., 2021; Ecclesia et al., 2016; He et al., 2023), it may not be as applicable to agricultural systems, where depending on management strategy, plant residues are often removed from the soil. Additionally, though land cover type may be an indirect indicator of C input quality, specific litter quality traits have demonstrated effects on C storage in POC and MAOC (e.g., Córdova et al., 2018; Craig et al., 2022; Haddix et al., 2016), and may represent a secondary control on SOC fractions, after climate-driven controls. Despite their relevance to this study, there was not enough exchangeable Ca, Fe- and Al-hydroxide, microbial, or litter quality data available in conjunction with C fraction data at the global scale to include them in our synthesis, prohibiting us from testing their effects on global C storage. Furthermore, there was relatively low representation of land cover types including tundra, wetlands, and shrublands in our dataset. In addition to potentially contributing to the relatively low explanatory power of our path analyses, this limited our ability to run more comprehensive analyses of land cover-specific controls

on C storage. In combination with an additional lack of standardized data reporting, particularly with regards to SOC and N fraction data, the above prevented us from more robustly testing several of the hypotheses presented in the In-N-Out framework (Cotrufo et al., 2021), as was our original intent. Given the difficulties of compiling large datasets for synthesis and meta-analysis projects, we echo recent calls in advocating for more standardized collection and reporting of fraction and ancillary data in SOC studies (Todd-Brown et al., 2022). Successful harmonization of SOC fraction data is an important next step in enabling more comprehensive testing of hypotheses related to SOC storage and N recycling, including the broad, global controls we aimed to test in this work, as well as ecosystem-specific controls that aid in development of site-specific strategies to prevent C loss. Both types of knowledge are critical to realizing the potential of soils to mitigate climate change and alleviate food insecurity.

In conclusion, our analyses revealed that global POC and MAOC storage are driven by separate environmental variables. Specifically, global POC storage is controlled primarily by C loss processes, while global MAOC storage is primarily controlled by constraints on C inputs and C stabilization mechanisms. These resulted in land cover types that tend to experience more rapid decomposition having a greater portion of their total soil C pool stored in MAOC. Despite these findings and demonstrating the effectiveness of fractionation in understanding global patterns of SOC storage, we were only able to explain 12.3% and 20.5% of the variation in POC and MAOC storage. As such, our work highlights the need for increased measurement of variables, including exchangeable Ca, Fe-, and Al-hydroxides, various types of microbial data, as well as fraction N, when conducting studies. Additionally, we invite the soils community to begin separating SOC into fractions beyond POC and MAOC, with the goal of improving our understanding of the likely contrasting behavior of free versus occluded POC and exchangeable versus stable MAOC. We hope that this improved measurement, combined with advancing soils data quality and reporting standards, will aid in effective data harmonization that can enable the field to build upon our findings. Doing so will allow us to develop the robust understanding of controls on soil C and N cycling needed to mitigate climate change and ensure the soil can continue to provide essential ecosystem services well into the future.

AUTHOR CONTRIBUTIONS

Paige M. Hansen: Conceptualization; data curation; formal analysis; investigation; methodology; project administration; validation; visualization; writing – original draft; writing – review and editing. **Rebecca Even:** Data curation; investigation; writing – original draft; writing – review and editing. **Alison E. King:** Data curation; formal analysis; investigation; writing – original draft; writing – review and editing. **Jocelyn Lavalée:** Conceptualization; data curation; formal analysis; funding acquisition; investigation; methodology; validation; writing – original draft; writing – review and editing. **Meagan Schipanski:** Conceptualization; formal analysis; funding acquisition; writing – original draft; writing – review and editing. **M. Francesca Cotrufo:**

Conceptualization; data curation; formal analysis; funding acquisition; investigation; methodology; project administration; supervision; validation; writing – original draft; writing – review and editing.

ACKNOWLEDGEMENTS

We thank members of the Cotrufo Soil Innovation Lab (SoIL) for their helpful suggestions on our analyses and manuscript, as well as Ann Hess for her suggestions on improving our path analyses. Soil samples were provided from the National Ecological Observatory Network (NEON) Megapit Soil Archive. NEON is a program sponsored by the National Science Foundation (NSF) and operated under cooperative agreement by Battelle. This material is based in part upon work supported by the NSF through the NEON Program, and the NSF-DEB award #2016003.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are openly available in Zenodo at <https://doi.org/10.5281/zenodo.10182813>.

ORCID

Paige M. Hansen  <https://orcid.org/0000-0002-9838-5601>

Rebecca Even  <https://orcid.org/0009-0005-0506-4841>

Alison E. King  <https://orcid.org/0000-0002-4155-577X>

Jocelyn Lavalley  <https://orcid.org/0000-0002-3028-7087>

Meagan Schipanski  <https://orcid.org/0000-0002-1661-9858>

M. Francesca Cotrufo  <https://orcid.org/0000-0002-6191-8953>

REFERENCES

- Abdalla, M., Hastings, A., Chadwick, D. R., Jones, D. L., Evans, C. D., Jones, M. B., Rees, R. M., & Smith, P. (2018). Critical review of the impacts of grazing intensity on soil organic carbon storage and other soil quality indicators in extensively managed grasslands. *Agriculture, Ecosystems & Environment*, 253, 62–81. <https://doi.org/10.1016/j.agee.2017.10.023>
- Angst, G., Mueller, K. E., Castellano, M. J., Vogel, C., Wiesmeier, M., & Mueller, C. W. (2023). Unlocking complex soil systems as carbon sinks: Multi-pool management as the key. *Nature Communications*, 14(1), 2967. <https://doi.org/10.1038/s41467-023-38700-5>
- Angst, G., Mueller, K. E., Nierop, K. G. J., & Simpson, M. J. (2021). Plant- or microbial-derived? A review on the molecular composition of stabilized soil organic matter. *Soil Biology and Biochemistry*, 156, 108189. <https://doi.org/10.1016/j.soilbio.2021.108189>
- Ashagrie, Y., Zech, W., Guggenberger, G., & Mamo, T. (2007). Soil aggregation, and total and particulate organic matter following conversion of native forests to continuous cultivation in Ethiopia. *Soil and Tillage Research*, 94(1), 101–108. <https://doi.org/10.1016/j.still.2006.07.005>
- Balesdent, J., Chenu, C., & Balabane, M. (2000). Relationship of soil organic matter dynamics to physical protection and tillage. *Soil and Tillage Research*, 53(3), 215–230. [https://doi.org/10.1016/S0167-1987\(99\)00107-5](https://doi.org/10.1016/S0167-1987(99)00107-5)
- Beck, H. E., Zimmermann, N. E., McVicar, T. R., Vergopolan, N., Berg, A., & Wood, E. F. (2018). Present and future Köppen-Geiger climate classification maps at 1-km resolution. *Scientific Data*, 5(1), 180214. <https://doi.org/10.1038/sdata.2018.214>
- Benbi, D. K., Boparai, A. K., & Brar, K. (2014). Decomposition of particulate organic matter is more sensitive to temperature than the mineral associated organic matter. *Soil Biology and Biochemistry*, 70, 183–192. <https://doi.org/10.1016/j.soilbio.2013.12.032>
- Bouajila, A., & Tahar, G. (2010). Land use effect on soil and particulate organic carbon, and aggregate stability in some soils in Tunisia. *African Journal of Agricultural Research*, 5(8), 764–774. <https://doi.org/10.5897/AJAR10.183>
- Bradford, M. A., Carey, C. J., Atwood, L., Bossio, D., Fenichel, E. P., Gennet, S., Fargione, J., Fisher, J. R. B., Fuller, E., Kane, D. A., Lehmann, J., Oldfield, E. E., Ordway, E. M., Rudek, J., Sanderman, J., & Wood, S. A. (2019). Soil carbon science for policy and practice. *Nature Sustainability*, 2(12), 1070–1072. <https://doi.org/10.1038/s41893-019-0431-y>
- Chen, L., Fang, K., Wei, B., Qin, S., Feng, X., Hu, T., Ji, C., & Yang, Y. (2021). Soil carbon persistence governed by plant input and mineral protection at regional and global scales. *Ecology Letters*, 24(5), 1018–1028. <https://doi.org/10.1111/ele.13723>
- Churkina, G., & Running, S. W. (1998). Contrasting climatic controls on the estimated productivity of global terrestrial biomes. *Ecosystems*, 1(2), 206–215. <https://doi.org/10.1007/s100219900016>
- Conant, R. T., Ryan, M. G., Ågren, G. I., Birge, H. E., Davidson, E. A., Eliasson, P. E., Evans, S. E., Frey, S. D., Giardina, C. P., Hopkins, F. M., Hyvönen, R., Kirschbaum, M. U. F., Lavalley, J. M., Leifeld, J., Parton, W. J., Megan Steinweg, J., Wallenstein, M. D., Martin Wetterstedt, J. Å., & Bradford, M. A. (2011). Temperature and soil organic matter decomposition rates—Synthesis of current knowledge and a way forward. *Global Change Biology*, 17(11), 3392–3404. <https://doi.org/10.1111/j.1365-2486.2011.02496.x>
- Córdova, S. C., Olk, D. C., Dietzel, R. N., Mueller, K. E., Archontoulis, S. V., & Castellano, M. J. (2018). Plant litter quality affects the accumulation rate, composition, and stability of mineral-associated soil organic matter. *Soil Biology and Biochemistry*, 125, 115–124. <https://doi.org/10.1016/j.soilbio.2018.07.010>
- Cotrufo, M. F., & Lavalley, J. M. (2022). Soil organic matter formation, persistence, and functioning: A synthesis of current understanding to inform its conservation and regeneration. In *Advances in agronomy* (Vol. 172, pp. 1–66). Elsevier. <https://doi.org/10.1016/bs.agron.2021.11.002>
- Cotrufo, M. F., Lavalley, J. M., Zhang, Y., Hansen, P. M., Paustian, K. H., Schipanski, M., & Wallenstein, M. D. (2021). I n-N-O ut: A hierarchical framework to understand and predict soil carbon storage and nitrogen recycling. *Global Change Biology*, 27(19), 4465–4468. <https://doi.org/10.1111/gcb.15782>
- Craig, M. E., Geyer, K. M., Beidler, K. V., Brzostek, E. R., Frey, S. D., Stuart Grandy, A., Liang, C., & Phillips, R. P. (2022). Fast-decaying plant litter enhances soil carbon in temperate forests but not through microbial physiological traits. *Nature Communications*, 13(1), 1229. <https://doi.org/10.1038/s41467-022-28715-9>
- Craig, M. E., Turner, B. L., Liang, C., Clay, K., Johnson, D. J., & Phillips, R. P. (2018). Tree mycorrhizal type predicts within-site variability in the storage and distribution of soil organic matter. *Global Change Biology*, 24(8), 3317–3330. <https://doi.org/10.1111/gcb.14132>
- Davidson, E. A., & Janssens, I. A. (2006). Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature*, 440(7081), 165–173. <https://doi.org/10.1038/nature04514>
- Eclesia, R. P., Jobbagy, E. G., Jackson, R. B., Rizzotto, M., & Piñeiro, G. (2016). Stabilization of new carbon inputs rather than old carbon decomposition determines soil organic carbon shifts following woody or herbaceous vegetation transitions. *Plant and Soil*, 409(1–2), 99–116. <https://doi.org/10.1007/s11104-016-2951-9>
- Ernakovich, J. G., Baldock, J., Creamer, C., Sanderman, J., Kalbitz, K., & Farrell, M. (2021). A combined microbial and ecosystem metric of carbon retention efficiency explains land cover-dependent soil microbial biodiversity–ecosystem function relationships. *Biogeochemistry*, 153(1), 1–15. <https://doi.org/10.1007/s10533-020-00736-w>

- Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*, 37(12), 4302–4315. <https://doi.org/10.1002/joc.5086>
- Frøseth, R. B., & Bleken, M. A. (2015). Effect of low temperature and soil type on the decomposition rate of soil organic carbon and clover leaves, and related priming effect. *Soil Biology and Biochemistry*, 80, 156–166. <https://doi.org/10.1016/j.soilbio.2014.10.004>
- Fuhrmann, J. J., & Zuberer, D. A. (2021). Carbon transformations and soil organic matter formation. In *Principles and applications of soil microbiology* (pp. 327–361). Elsevier. <https://doi.org/10.1016/B978-0-12-820202-9.00013-7>
- Georgiou, K., Jackson, R. B., Vinduřková, O., Abramoff, R. Z., Ahlström, A., Feng, W., Harden, J. W., Pellegrini, A. F. A., Polley, H. W., Soong, J. L., Riley, W. J., & Torn, M. S. (2022). Global stocks and capacity of mineral-associated soil organic carbon. *Nature Communications*, 13(1), 3797. <https://doi.org/10.1038/s41467-022-31540-9>
- Haddix, M. L., Gregorich, E. G., Helgason, B. L., Janzen, H., Ellert, B. H., & Francesca Cotrufo, M. (2020). Climate, carbon content, and soil texture control the independent formation and persistence of particulate and mineral-associated organic matter in soil. *Geoderma*, 363, 114160. <https://doi.org/10.1016/j.geoderma.2019.114160>
- Haddix, M. L., Paul, E. A., & Cotrufo, M. F. (2016). Dual, differential isotope labeling shows the preferential movement of labile plant constituents into mineral-bonded soil organic matter. *Global Change Biology*, 22(6), 2301–2312. <https://doi.org/10.1111/gcb.13237>
- Harris, D., Horwath, W. R., & van Kessel, C. (2001). Acid fumigation of soils to remove carbonates prior to total organic carbon or CARBON-13 isotopic analysis. *Soil Science Society of America Journal*, 65(6), 1853–1856. <https://doi.org/10.2136/sssaj2001.1853>
- Harzing, A.-W., & Alakangas, S. (2016). Google scholar, Scopus and the web of science: A longitudinal and cross-disciplinary comparison. *Scientometrics*, 106(2), 787–804. <https://doi.org/10.1007/s11192-015-1798-9>
- Hassink, J. (1997). The capacity of soils to preserve organic C and N by their association with clay and silt particles. *Plant and Soil*, 191(1), 77–87. <https://doi.org/10.1023/A:1004213929699>
- He, X., Abramoff, R., Abs, E., Georgiou, K., Zhang, H., & Goll, D. S. (2023). Contribution of carbon inputs to soil carbon accumulation cannot be neglected. *bioRxiv*. <https://doi.org/10.1101/2023.07.17.549330>
- Heckman, K., Hicks Pries, C. E., Lawrence, C. R., Rasmussen, C., Crow, S. E., Hoyt, A. M., Fromm, S. F., Shi, Z., Stoner, S., McGrath, C., Beem-Miller, J., Berhe, A. A., Blankinship, J. C., Keiluweit, M., Marin-Spiotta, E., Monroe, J. G., Plante, A. F., Schimel, J., Sierra, C. A., ... Wagai, R. (2022). Beyond bulk: Density fractions explain heterogeneity in global soil carbon abundance and persistence. *Global Change Biology*, 28(3), 1178–1196. <https://doi.org/10.1111/gcb.16023>
- Heckman, K. A., Possinger, A. R., Badgley, B. D., Bowman, M. M., Gallo, A. C., Hatten, J. A., Nave, L. E., SanClements, M. D., Swanston, C. W., Weiglein, T. L., Wieder, W. R., & Strahm, B. D. (2023). Moisture-driven divergence in mineral-associated soil carbon persistence. *Proceedings of the National Academy of Sciences of the United States of America*, 120(7), e2210044120. <https://doi.org/10.1073/pnas.2210044120>
- Herndon, E., AlBashaireh, A., Singer, D., Roy Chowdhury, T., Gu, B., & Graham, D. (2017). Influence of iron redox cycling on organo-mineral associations in Arctic tundra soil. *Geochimica et Cosmochimica Acta*, 207, 210–231. <https://doi.org/10.1016/j.gca.2017.02.034>
- Hinckley, E.-L. S., Bonan, G. B., Bowen, G. J., Colman, B. P., Duffy, P. A., Goodale, C. L., Houlton, B. Z., Marin-Spiotta, E., Ogle, K., Ollinger, S. V., Paul, E. A., Vitousek, P. M., Weathers, K. C., & Williams, D. G. (2016). The soil and plant biogeochemistry sampling design for The National Ecological Observatory Network. *Ecosphere*, 7(3), e01234. <https://doi.org/10.1002/ecs2.1234>
- Hooper, D., Coughlan, J., & Mullen, M. R. (2008). Structural equation modelling: Guidelines for determining model fit. *Electronic Journal of Business Research Methods*, 6(1), 53–60.
- Horsch, C. C. A., Antunes, P. M., Fahey, C., Grandy, A. S., & Kallenbach, C. M. (2023). Trait-based assembly of arbuscular mycorrhizal fungal communities determines soil carbon formation and retention. *New Phytologist*, 239(1), 311–324. <https://doi.org/10.1111/nph.18914>
- Hou, G., Delang, C. O., Lu, X., & Gao, L. (2020). A meta-analysis of changes in soil organic carbon stocks after afforestation with deciduous broadleaved, sempervirent broadleaved, and conifer tree species. *Annals of Forest Science*, 77(4), 1–13. <https://doi.org/10.1007/s13595-020-00997-3>
- Huang, W., Ye, C., Hockaday, W. C., & Hall, S. J. (2020). Trade-offs in soil carbon protection mechanisms under aerobic and anaerobic conditions. *Global Change Biology*, 26(6), 3726–3737. <https://doi.org/10.1111/gcb.15100>
- Huang, Y., Liang, C., Duan, X., Chen, H., & Li, D. (2019). Variation of microbial residue contribution to soil organic carbon sequestration following land use change in a subtropical karst region. *Geoderma*, 353, 340–346. <https://doi.org/10.1016/j.geoderma.2019.07.028>
- Jobbágy, E. G., & Jackson, R. B. (2000). The vertical distribution of soil organic carbon and its relation to climate and vegetation. *Ecological Applications*, 10(2), 423–436. [https://doi.org/10.1890/1051-0761\(2000\)010\[0423:TVDOSO\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[0423:TVDOSO]2.0.CO;2)
- Kallenbach, C. M., Frey, S. D., & Grandy, A. S. (2016). Direct evidence for microbial-derived soil organic matter formation and its ecophysiological controls. *Nature Communications*, 7(1), 13630. <https://doi.org/10.1038/ncomms13630>
- Kassambra, A. (2023). *rstatix: Pipe-friendly framework for basic statistical tests*. <https://CRAN.R-project.org/package=rstatix>
- Keller, A. B., Brzostek, E. R., Craig, M. E., Fisher, J. B., & Phillips, R. P. (2021). Root-derived inputs are major contributors to soil carbon in temperate forests, but vary by mycorrhizal type. *Ecology Letters*, 24(4), 626–635. <https://doi.org/10.1111/ele.13651>
- King, A. E., Amsili, J. P., Córdova, S. C., Culman, S., Fonte, S. J., Kotcon, J., Liebig, M., Masters, M. D., McVay, K., Olk, D. C., Schipanski, M., Schneider, S. K., Stewart, C. E., & Cotrufo, M. F. (2023). A soil matrix capacity index to predict mineral-associated but not particulate organic carbon across a range of climate and soil pH. *Biogeochemistry*, 165(1), 1–14. <https://doi.org/10.1007/s10533-023-01066-3>
- Kirsten, M., Mikutta, R., Vogel, C., Thompson, A., Mueller, C. W., Kimaro, D. N., Bergsma, H. L. T., Feger, K.-H., & Kalbitz, K. (2021). Iron oxides and aluminous clays selectively control soil carbon storage and stability in the humid tropics. *Scientific Reports*, 11, 5076. <https://doi.org/10.1038/s41598-021-84777-7>
- Kramer, M. G., & Chadwick, O. A. (2018). Climate-driven thresholds in reactive mineral retention of soil carbon at the global scale. *Nature Clim Change*, 8, 1104–1108. <https://doi.org/10.1038/s41558-018-0341-4>
- Krishna, M. P., & Mohan, M. (2017). Litter decomposition in forest ecosystems: A review. *Energy, Ecology and Environment*, 2(4), 236–249. <https://doi.org/10.1007/s40974-017-0064-9>
- Lajtha, K., Townsend, K. L., Kramer, M. G., Swanston, C., Bowden, R. D., & Nadelhoffer, K. (2014). Changes to particulate versus mineral-associated soil carbon after 50 years of litter manipulation in forest and prairie experimental ecosystems. *Biogeochemistry*, 119(1), 341–360. <https://doi.org/10.1007/s10533-014-9970-5>
- Lavallee, J. M., Conant, R. T., Haddix, M. L., Follett, R. F., Bird, M. I., & Paul, E. A. (2019). Selective preservation of pyrogenic carbon across soil organic matter fractions and its influence on calculations of carbon mean residence times. *Geoderma*, 354, 113866. <https://doi.org/10.1016/j.geoderma.2019.07.024>
- Lavallee, J. M., Soong, J. L., & Cotrufo, M. F. (2020). Conceptualizing soil organic matter into particulate and mineral-associated forms to address global change in the 21st century. *Global Change Biology*, 26(1), 261–273. <https://doi.org/10.1111/gcb.14859>
- Leuthold, S. J., Haddix, M. L., Lavallee, J., & Cotrufo, M. F. (2022). Physical fractionation techniques. In *Reference Module in Earth Systems and*

- Environmental Sciences*. Elsevier. <https://doi.org/10.1016/B978-0-12-822974-3.00067-7>
- Leuthold, S. J., Lavalley, J. Haddix, M., & Cotrufo, M. (in review). *Contrasting properties of soil organic matter fractions isolated by different physical separation methodologies*. *Geoderma*.
- Liang, C., Amelung, W., Lehmann, J., & Kästner, M. (2019). Quantitative assessment of microbial necromass contribution to soil organic matter. *Global Change Biology*, 25(11), 3578–3590. <https://doi.org/10.1111/gcb.14781>
- Lobe, I., Sandhage-Hofmann, A., Brodowski, S., du Preez, C. C., & Amelung, W. (2011). Aggregate dynamics and associated soil organic matter contents as influenced by prolonged arable cropping in the South African Highveld. *Geoderma*, 162(3), 251–259. <https://doi.org/10.1016/j.geoderma.2011.02.001>
- Lugato, E., Lavalley, J. M., Haddix, M. L., Panagos, P., & Cotrufo, M. F. (2021). Different climate sensitivity of particulate and mineral-associated soil organic matter. *Nature Geoscience*, 14(5), 295–300. <https://doi.org/10.1038/s41561-021-00744-x>
- Luo, Z., Feng, W., Luo, Y., Baldock, J., & Wang, E. (2017). Soil organic carbon dynamics jointly controlled by climate, carbon inputs, soil properties and soil carbon fractions. *Global Change Biology*, 23(10), 4430–4439. <https://doi.org/10.1111/gcb.13767>
- Lupwayi, N. Z., Clayton, G. W., O'Donovan, J. T., Harker, K. N., Turkington, T. K., & Rice, W. A. (2004). Decomposition of crop residues under conventional and zero tillage. *Canadian Journal of Soil Science*, 84(4), 403–410. <https://doi.org/10.4141/S03-082>
- Mao, H., Cotrufo, M. F., Hart, S. C., Sullivan, B. W., Zhu, X., Zhang, J., Liang, C., & Zhu, M. (In review). *Soil Biology & Biochemistry*.
- Martín-Martín, A., Orduna-Malea, E., Thelwall, M., & Delgado López-Cózar, E. (2018). Google Scholar, Web of Science, and Scopus: A systematic comparison of citations in 252 subject categories. *Journal of Informetrics*, 12(4), 1160–1177. <https://doi.org/10.1016/j.joi.2018.09.002>
- Mirabito, A. J., & Chambers, L. G. (2023). Quantifying mineral-associated organic matter in wetlands as an indicator of the degree of soil carbon protection. *Geoderma*, 430, 116327. <https://doi.org/10.1016/j.geoderma.2023.116327>
- Ogle, S. M., Breidt, F. J., & Paustian, K. (2005). Agricultural management impacts on soil organic carbon storage under moist and dry climatic conditions of temperate and tropical regions. *Biogeochemistry*, 72(1), 87–121. <https://doi.org/10.1007/s10533-004-0360-2>
- Parihar, C. M., Singh, A. K., Jat, S. L., Ghosh, A., Dey, A., Nayak, H. S., Parihar, M. D., Mahala, D. M., Yadav, R. K., Rai, V., Satyanaryana, T., & Jat, M. L. (2019). Dependence of temperature sensitivity of soil organic carbon decomposition on nutrient management options under conservation agriculture in a sub-tropical Inceptisol. *Soil and Tillage Research*, 190, 50–60. <https://doi.org/10.1016/j.still.2019.02.016>
- Pérez-Harguindeguy, N., Díaz, S., Cornelissen, J. H. C., Vendramini, F., Cabido, M., & Castellanos, A. (2000). Chemistry and toughness predict leaf litter decomposition rates over a wide spectrum of functional types and taxa in central Argentina. *Plant and Soil*, 218(1/2), 21–30.
- Poepplau, C., Don, A., Six, J., Kaiser, M., Benbi, D., Chenu, C., Cotrufo, M. F., Derrien, D., Gioacchini, P., Grand, S., Gregorich, E., Griepentrog, M., Gunina, A., Haddix, M., Kuzyakov, Y., Kühnel, A., Macdonald, L. M., Soong, J., Trigalet, S., ... Nieder, R. (2018). Isolating organic carbon fractions with varying turnover rates in temperate agricultural soils—A comprehensive method comparison. *Soil Biology and Biochemistry*, 125, 10–26. <https://doi.org/10.1016/j.soilbio.2018.06.025>
- Poepplau, C., Kätterer, T., Leblans, N. I. W., & Sigurdsson, B. D. (2017). Sensitivity of soil carbon fractions and their specific stabilization mechanisms to extreme soil warming in a subarctic grassland. *Global Change Biology*, 23(3), 1316–1327. <https://doi.org/10.1111/gcb.13491>
- Prairie, A. M., King, A. E., & Cotrufo, M. F. (2023). Restoring particulate and mineral-associated organic carbon through regenerative agriculture. *Proceedings of the National Academy of Sciences of the United States of America*, 120(21), e2217481120. <https://doi.org/10.1073/pnas.2217481120>
- R Core Team. (2021). *R: A language and environment for statistical computing* (4.1.1) [Computer software]. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Ramos, F. T., Dores, E. F. D. C., Weber, O. L. D. S., Beber, D. C., Campelo, J. H., & Maia, J. C. D. S. (2018). Soil organic matter doubles the cation exchange capacity of tropical soil under no-till farming in Brazil. *Journal of the Science of Food and Agriculture*, 98(9), 3595–3602. <https://doi.org/10.1002/jsfa.8881>
- Rasmussen, C., Heckman, K., Wieder, W. R., Keiluweit, M., Lawrence, C. R., Berhe, A. A., Blankinship, J. C., Crow, S. E., Druhan, J. L., Hicks Pries, C. E., Marin-Spiotta, E., Plante, A. F., Schädel, C., Schimel, J. P., Sierra, C. A., Thompson, A., & Wagai, R. (2018). Beyond clay: Towards an improved set of variables for predicting soil organic matter content. *Biogeochemistry*, 137(3), 297–306. <https://doi.org/10.1007/s10533-018-0424-3>
- Rocci, K. S., Lavalley, J. M., Stewart, C. E., & Cotrufo, M. F. (2021). Soil organic carbon response to global environmental change depends on its distribution between mineral-associated and particulate organic matter: A meta-analysis. *Science of the Total Environment*, 793, 148569. <https://doi.org/10.1016/j.scitotenv.2021.148569>
- Rosseel, Y. (2012). lavaan: An R package for structural equation modeling. *Journal of Statistical Software*, 48(2), 1–36. <https://doi.org/10.18637/jss.v048.i02>
- Rousk, J., Brookes, P. C., & Bååth, E. (2009). Contrasting soil pH effects on fungal and bacterial growth suggest functional redundancy in carbon mineralization. *Applied and Environmental Microbiology*, 75(6), 1589–1596. <https://doi.org/10.1128/AEM.02775-08>
- Rowley, M. C., Grand, S., Spangenberg, J. E., & Verrecchia, E. P. (2021). Evidence linking calcium to increased organo-mineral association in soils. *Biogeochemistry*, 153(3), 223–241. <https://doi.org/10.1007/s10533-021-00779-7>
- Rumpel, C., Amiraslani, F., Koutika, L.-S., Smith, P., Whitehead, D., & Wollenberg, E. (2018). Put more carbon in soils to meet Paris climate pledges. *Nature*, 564(7734), 32–34. <https://doi.org/10.1038/d41586-018-07587-4>
- Running, S., Mu, Q., Zhao, M., & Moreno, A. (2021). MODIS/Terra net evapotranspiration gap-filled yearly L4 global 500m SIN grid V061. NASA EOSDIS Land Processes DAAC. <https://doi.org/10.5067/MODIS/MOD16A3GF.061>
- Running, S., & Zhao, M. (2019). MOD17A3HGF MODIS/Terra net primary production gap-filled yearly L4 global 500 m SIN grid V006. NASA EOSDIS Land Processes DAAC. <https://doi.org/10.5067/MODIS/MOD17A3HGF.006>
- Schimel, J. P. (2018). Life in dry soils: Effects of drought on soil microbial communities and processes. *Annual Review of Ecology, Evolution, and Systematics*, 49(1), 409–432. <https://doi.org/10.1146/annurev-ecolsys-110617-062614>
- Shipley, B. (2016). *Cause and correlation in biology: A user's guide to path analysis, structural equations and causal inference with R* (2nd ed.). Cambridge University Press. <https://doi.org/10.1017/CBO9781139979573>
- Six, J., Conant, R. T., Paul, E. A., & Paustian, K. (2002). Stabilization mechanisms of soil organic matter: Implications for C-saturation of soils. *Plant and Soil*, 241, 155–176.
- Sokol, N. W., Whalen, E. D., Jilling, A., Kallenbach, C., Pett-Ridge, J., & Georgiou, K. (2022). Global distribution, formation and fate of mineral-associated soil organic matter under a changing climate: A trait-based perspective. *Functional Ecology*, 36(6), 1411–1429. <https://doi.org/10.1111/1365-2435.14040>
- Song, B., Niu, S., Li, L., Zhang, L., & Yu, G. (2014). Soil carbon fractions in grasslands respond differently to various levels of nitrogen

- enrichments. *Plant and Soil*, 384(1), 401–412. <https://doi.org/10.1007/s11104-014-2219-1>
- Sousa, R. F. D., Brasil, E. P. F., Figueiredo, C. C. D., & Leandro, W. M. (2015). Soil organic matter fractions in preserved and disturbed wetlands of the Cerrado biome. *Revista Brasileira de Ciência Do Solo*, 39(1), 222–231. <https://doi.org/10.1590/01000683rbcbs20150048>
- Thaysen, E. M., Reinsch, S., Larsen, K. S., & Ambus, P. (2017). Decrease in heathland soil labile organic carbon under future atmospheric and climatic conditions. *Biogeochemistry*, 133(1), 17–36. <https://doi.org/10.1007/s10533-017-0303-3>
- Todd-Brown, K. E. O., Abramoff, R. Z., Beem-Miller, J., Blair, H. K., Earl, S., Frederick, K. J., Fuku, D. R., Guevara Santamaria, M., Harden, J. W., Heckman, K., Heran, L. J., Holmquist, J. R., Hoyt, A. M., Klings, D. H., LeBauer, D. S., Malhotra, A., McClelland, S. C., Nave, L. E., Rocci, K. S., ... Younger, M. L. (2022). Reviews and syntheses: The promise of big diverse soil data, moving current practices towards future potential. *Biogeosciences*, 19(14), 3505–3522. <https://doi.org/10.5194/bg-19-3505-2022>
- Vermeulen, S., Bossio, D., Lehmann, J., Luu, P., Paustian, K., Webb, C., Augé, F., Bacudo, I., Baedeker, T., Havemann, T., Jones, C., King, R., Reddy, M., Sunga, I., Von Unger, M., & Warkner, M. (2019). A global agenda for collective action on soil carbon. *Nature Sustainability*, 2(1), 2–4. <https://doi.org/10.1038/s41893-018-0212-z>
- von Lützow, M., Kögel-Knabner, I., Ekschmitt, K., Flessa, H., Guggenberger, G., Matzner, E., & Marschner, B. (2007). SOM fractionation methods: Relevance to functional pools and to stabilization mechanisms. *Soil Biology and Biochemistry*, 39(9), 2183–2207. <https://doi.org/10.1016/j.soilbio.2007.03.007>
- Wiesmeier, M., Urbanski, L., Hobbey, E., Lang, B., von Lützow, M., Marin-Spiotta, E., van Wesemael, B., Rabot, E., Ließ, M., Garcia-Franco, N., Wollschläger, U., Vogel, H.-J., & Kögel-Knabner, I. (2019). Soil organic carbon storage as a key function of soils—A review of drivers and indicators at various scales. *Geoderma*, 333, 149–162. <https://doi.org/10.1016/j.geoderma.2018.07.026>
- Wu, J., Zhang, H., Pan, Y., Cheng, X., Zhang, K., & Liu, G. (2023). Particulate organic carbon is more sensitive to nitrogen addition than mineral-associated organic carbon: A meta-analysis. *Soil and Tillage Research*, 232, 105770. <https://doi.org/10.1016/j.still.2023.105770>
- Xu, C., Guo, L., Ping, C.-L., & White, D. M. (2009). Chemical and isotopic characterization of size-fractionated organic matter from cryoturbated tundra soils, northern Alaska. *Journal of Geophysical Research: Biogeosciences*, 114(G3), G03002. <https://doi.org/10.1029/2008JG000846>
- Yu, W., Huang, W., Weintraub-Leff, S. R., & Hall, S. J. (2022). Where and why do particulate organic matter (POM) and mineral-associated organic matter (MAOM) differ among diverse soils? *Soil Biology and Biochemistry*, 172, 108756. <https://doi.org/10.1016/j.soilbio.2022.108756>
- Zhang, Y., Lavalley, J. M., Robertson, A. D., Even, R., Ogle, S. M., Paustian, K., & Cotrufo, M. F. (2021). Simulating measurable ecosystem carbon and nitrogen dynamics with the mechanistically defined MEMS 2.0 model. *Biogeosciences*, 18(10), 3147–3171. <https://doi.org/10.5194/bg-18-3147-2021>
- some tropical soils. *Geoderma*, 143(1–2), 14–25. <https://doi.org/10.1016/j.geoderma.2007.10.003>
- Błońska, E., Lasota, J., Tullus, A., Lutter, R., & Ostonen, I. (2019). Impact of dead-wood decomposition on soil organic carbon sequestration in Estonian and Polish forests. *Annals of Forest Science*, 76(4), 102. <https://doi.org/10.1007/s13595-019-0889-9>
- Caravaca, F., Lax, A., & Albaladejo, J. (1999). Organic matter, nutrient contents and cation exchange capacity in fine fractions from semiarid calcareous soils. *Geoderma*, 93(3–4), 161–176. [https://doi.org/10.1016/S0016-7061\(99\)00045-2](https://doi.org/10.1016/S0016-7061(99)00045-2)
- Conen, F., Zimmermann, M., Leifeld, J., Seth, B., & Alewell, C. (2008). Relative stability of soil carbon revealed by shifts in $\delta^{15}\text{N}$ and C:N ratio. *Biogeosciences*, 5(1), 123–128. <https://doi.org/10.5194/bg-5-123-2008>
- Cui, J., Li, Z., Liu, Z., Ge, B., Fang, C., Zhou, C., & Tang, B. (2014). Physical and chemical stabilization of soil organic carbon along a 500-year cultivated soil chronosequence originating from estuarine wetlands: Temporal patterns and land use effects. *Agriculture, Ecosystems & Environment*, 196, 10–20. <https://doi.org/10.1016/j.agee.2014.06.013>
- Dietterich, L. H., Karpman, J., Neupane, A., Ciochina, M., & Cusack, D. F. (2021). Carbon content of soil fractions varies with season, rainfall, and soil fertility across a lowland tropical moist forest gradient. *Biogeochemistry*, 155(3), 431–452. <https://doi.org/10.1007/s10533-021-00836-1>
- Doetterl, S., Stevens, A., Six, J., Merckx, R., Van Oost, K., Casanova Pinto, M., Casanova-Katny, A., Muñoz, C., Boudin, M., Zagal Venegas, E., & Boeckx, P. (2015). Soil carbon storage controlled by interactions between geochemistry and climate. *Nature Geoscience*, 8(10), 780–783. <https://doi.org/10.1038/ngeo2516>
- Dutta, K., Schuur, E. A. G., Neff, J. C., & Zimov, S. A. (2006). Potential carbon release from permafrost soils of Northeastern Siberia. *Global Change Biology*, 12(12), 2336–2351. <https://doi.org/10.1111/j.1365-2486.2006.01259.x>
- Feller, C., Albrecht, A., & Tessier, D. (1996). Aggregation and organic matter storage in kaolinitic and smectitic tropical soils. In M. R. Carter & B. A. Stewart (Eds.), *Structure and organic matter storage in agricultural soils* (1st ed., pp. 309–359). CRC Press. <https://doi.org/10.1201/9781003075561-12>
- Georgiou, K., Jackson, R. B., Vinduškova, O., Abramoff, R. Z., Ahlström, A., Feng, W., Harden, J. W., Pellegrini, A. F. A., Polley, H. W., Soong, J. L., Riley, W. J., & Torn, M. S. (2022). Global stocks and capacity of mineral-associated soil organic carbon. *Nature Communications*, 13(1), 3797. <https://doi.org/10.1038/s41467-022-31540-9>
- Gong, W., Yan, X., Wang, J., Hu, T., & Gong, Y. (2009). Long-term manure and fertilizer effects on soil organic matter fractions and microbes under a wheat–maize cropping system in northern China. *Geoderma*, 149(3–4), 318–324. <https://doi.org/10.1016/j.geoderma.2008.12.010>
- Guggenberger, G., Christensen, B. T., & Zech, W. (1994). Land-use effects on the composition of organic matter in particle-size separates of soil: I. Lignin and carbohydrate signature. *European Journal of Soil Science*, 45(4), 449–458. <https://doi.org/10.1111/j.1365-2389.1994.tb00530.x>
- Guibert, H., Fallavier, P., & Roméro, J. (1999). Carbon content in soil particle size and consequence on cation exchange capacity of Alfisols. *Communications in Soil Science and Plant Analysis*, 30(17–18), 2521–2537. <https://doi.org/10.1080/00103629909370392>
- Gutiérrez del Arroyo, O., & Silver, W. L. (2018). Disentangling the long-term effects of disturbance on soil biogeochemistry in a wet tropical forest ecosystem. *Global Change Biology*, 24(4), 1673–1684. <https://doi.org/10.1111/gcb.14027>
- Hall, S. J., Berhe, A. A., & Thompson, A. (2018). Order from disorder: Do soil organic matter composition and turnover co-vary with iron phase crystallinity? *Biogeochemistry*, 140(1), 93–110. <https://doi.org/10.1007/s10533-018-0476-4>
- Hall, S. J., McNicol, G., Natake, T., & Silver, W. L. (2015). Large fluxes and rapid turnover of mineral-associated carbon across topographic gradients in a humid tropical forest: Insights from paired $\delta^{13}\text{C}$ analysis. *Biogeosciences*, 12(8), 2471–2487. <https://doi.org/10.5194/bg-12-2471-2015>
- Harden, J. W. (1987). *Soils developed in granitic alluvium near Merced, California*. USGPO. <https://doi.org/10.3133/b1590A>
- Harden, J. W., Fries, T. L., & Pavich, M. J. (2002). Cycling of beryllium and carbon through hillslope soils in Iowa. *Biogeochemistry*, 60(3), 317–336. <https://doi.org/10.1023/A:1020308729553>
- Hatton, P.-J., Kleber, M., Zeller, B., Moni, C., Plante, A. F., Townsend, K., Gelhaye, L., Lajtha, K., & Derrien, D. (2012). Transfer of litter-derived N to soil mineral–organic associations: Evidence from decadal ^{15}N tracer experiments. *Organic Geochemistry*, 42(12), 1489–1501. <https://doi.org/10.1016/j.orggeochem.2011.05.002>

DATA SOURCES

- Balabane, M., & Plante, A. F. (2004). Aggregation and carbon storage in silty soil using physical fractionation techniques. *European Journal of Soil Science*, 55(2), 415–427. <https://doi.org/10.1111/j.1351-0754.2004.0608.x>
- Balesdent, J., Besnard, E., Arrouays, D., & Chenu, C. (1998). The dynamics of carbon in particle-size fractions of soil in a forest-cultivation sequence. *Plant and Soil*, 201(1), 49–57. <https://doi.org/10.1023/A:1004337314970>
- Barthès, B. G., Kouakoua, E., Larré-Larrouy, M.-C., Razafimbelo, T. M., de Luca, E. F., Azontonde, A., Neves, C. S. V. J., de Freitas, P. L., & Feller, C. L. (2008). Texture and sesquioxide effects on water-stable aggregates and organic matter in

- Heckman, K., Lawrence, C. R., & Harden, J. W. (2018). A sequential selective dissolution method to quantify storage and stability of organic carbon associated with Al and Fe hydroxide phases. *Geoderma*, 312, 24–35. <https://doi.org/10.1016/j.geoderma.2017.09.043>
- Heckman, K. A. (2010). *Pedogenesis & carbon dynamics across a lithosequence under ponderosa pine*. <https://doi.org/10.5281/ZENODO.1486081>
- Herndon, E., AlBashaireh, A., Singer, D., Roy Chowdhury, T., Gu, B., & Graham, D. (2017). Influence of iron redox cycling on organo-mineral associations in Arctic tundra soil. *Geochimica et Cosmochimica Acta*, 207, 210–231. <https://doi.org/10.1016/j.gca.2017.02.034>
- Jindaluang, W., Kheoruenromne, I., Suddhiprakarn, A., Singh, B. P., & Singh, B. (2013). Influence of soil texture and mineralogy on organic matter content and composition in physically separated fractions soils of Thailand. *Geoderma*, 195–196, 207–219. <https://doi.org/10.1016/j.geoderma.2012.12.003>
- Keller, A. B., Borer, E. T., Collins, S. L., DeLancey, L. C., Fay, P. A., Hofmockel, K. S., Leakey, A. D. B., Mayes, M. A., Seabloom, E. W., Walter, C. A., Wang, Y., Zhao, Q., & Hobbie, S. E. (2022). Soil carbon stocks in temperate grasslands differ strongly across sites but are insensitive to decade-long fertilization. *Global Change Biology*, 28(4), 1659–1677. <https://doi.org/10.1111/gcb.15988>
- Khomo, L., Trumbore, S., Bern, C. R., & Chadwick, O. A. (2017). Timescales of carbon turnover in soils with mixed crystalline mineralogies. *The Soil*, 3(1), 17–30. <https://doi.org/10.5194/soil-3-17-2017>
- King, A. E., Amsili, J. P., Córdova, S. C., Culman, S., Fonte, S. J., Kotcon, J., Liebigh, M., Masters, M. D., McVay, K., Olk, D. C., Schipanski, M., Schneider, S. K., Stewart, C. E., & Cotrufo, M. F. (2023). A soil matrix capacity index to predict mineral-associated but not particulate organic carbon across a range of climate and soil pH. *Biogeochemistry*, 165(1), 1–14. <https://doi.org/10.1007/s10533-023-01066-3>
- Köhl, A., von Lützow, M., & Kögel-Knabner, I. (2006). Decomposition and distribution of ¹⁵N labelled mustard litter (*Sinapis alba*) in physical soil fractions of a cropland with high- and low-yield field areas. *Soil Biology and Biochemistry*, 38(11), 3292–3302. <https://doi.org/10.1016/j.soilbio.2006.04.010>
- Kong, A. Y. Y., Fonte, S. J., van Kessel, C., & Six, J. (2009). Transitioning from standard to minimum tillage: Trade-offs between soil organic matter stabilization, nitrous oxide emissions, and N availability in irrigated cropping systems. *Soil and Tillage Research*, 104(2), 256–262. <https://doi.org/10.1016/j.still.2009.03.004>
- Larré-Larrouy, M. C., Albrecht, A., Blanchart, E., Chevallier, T., & Feller, C. (2003). Carbon and monosaccharides of a tropical vertisol under pasture and market-gardening: Distribution in primary organomineral separates. *Geoderma*, 117(1–2), 63–79. [https://doi.org/10.1016/S0016-7061\(03\)00135-6](https://doi.org/10.1016/S0016-7061(03)00135-6)
- Lawrence, C. R., Beem-Miller, J., Hoyt, A. M., Monroe, G., Sierra, C. A., Stoner, S., Heckman, K., Blankinship, J. C., Crow, S. E., McNicol, G., Trumbore, S., Levine, P. A., Vinduškova, O., Todd-Brown, K., Rasmussen, C., Hicks Pries, C. E., Schädel, C., McFarlane, K., Doetterl, S., ... Wagai, R. (2020). An open-source database for the synthesis of soil radiocarbon data: International soil radiocarbon database (ISRad) version 1.0. *Earth System Science Data*, 12(1), 61–76. <https://doi.org/10.5194/essd-12-61-2020>
- Li, Y., Xu, M., Zou, X., Shi, P., & Zhang, Y. (2005). Comparing soil organic carbon dynamics in plantation and secondary forest in wet tropics in Puerto Rico. *Global Change Biology*, 11(2), 239–248. <https://doi.org/10.1111/j.1365-2486.2005.00896.x>
- Liang, A., Yang, X., Zhang, X., McLaughlin, N., Shen, Y., & Li, W. (2009). Soil organic carbon changes in particle-size fractions following cultivation of Black soils in China. *Soil and Tillage Research*, 105(1), 21–26. <https://doi.org/10.1016/j.still.2009.05.002>
- Lugato, E., Lavallee, J. M., Haddix, M. L., Panagos, P., & Cotrufo, M. F. (2021). Different climate sensitivity of particulate and mineral-associated soil organic matter. *Nature Geoscience*, 14(5), 295–300. <https://doi.org/10.1038/s41561-021-00744-x>
- Lybrand, R. A., Heckman, K., & Rasmussen, C. (2017). Soil organic carbon partitioning and $\delta^{14}\text{C}$ variation in desert and conifer ecosystems of southern Arizona. *Biogeochemistry*, 134(3), 261–277. <https://doi.org/10.1007/s10533-017-0360-7>
- Marín-Spiotta, E., Swanston, C. W., Torn, M. S., Silver, W. L., & Burton, S. D. (2008). Chemical and mineral control of soil carbon turnover in abandoned tropical pastures. *Geoderma*, 143(1–2), 49–62. <https://doi.org/10.1016/j.geoderma.2007.10.001>
- Mirabito, A. J., & Chambers, L. G. (2023). Quantifying mineral-associated organic matter in wetlands as an indicator of the degree of soil carbon protection. *Geoderma*, 430, 116327. <https://doi.org/10.1016/j.geoderma.2023.116327>
- Mosier, S., Apfelbaum, S., Byck, P., Calderon, F., Teague, R., Thompson, R., & Cotrufo, M. F. (2021). Adaptive multi-paddock grazing enhances soil carbon and nitrogen stocks and stabilization through mineral association in south-eastern U.S. grazing lands. *Journal of Environmental Management*, 288, 112409. <https://doi.org/10.1016/j.jenvman.2021.112409>
- Mosier, S., Paustian, K., Davies, C., Kane, M., & Cotrufo, M. F. (2019). Soil organic matter pools under management intensification of loblolly pine plantations. *Forest Ecology and Management*, 447, 60–66. <https://doi.org/10.1016/j.foreco.2019.05.056>
- Mujuru, L., Mureva, A., Velthorst, E. J., & Hoosbeek, M. R. (2013). Land use and management effects on soil organic matter fractions in Rhodic Ferralsols and Haplic Arenosols in Bindura and Shamva districts of Zimbabwe. *Geoderma*, 209–210, 262–272. <https://doi.org/10.1016/j.geoderma.2013.06.025>
- Nacro, H. B., Benest, D., & Abbadié, L. (1996). Distribution of microbial activities and organic matter according to particle size in a humid savanna soil (Lamto, Côte D'Ivoire). *Soil Biology and Biochemistry*, 28(12), 1687–1697. [https://doi.org/10.1016/S0038-0717\(96\)00246-5](https://doi.org/10.1016/S0038-0717(96)00246-5)
- Piñeiro, G., Paruelo, J. M., Jobbágy, E. G., Jackson, R. B., & Oesterheld, M. (2009). Grazing effects on belowground C and N stocks along a network of cattle enclosures in temperate and subtropical grasslands of South America. *Global Biogeochemical Cycles*, 23(2), GB2003. <https://doi.org/10.1029/2007G003168>
- Pold, G., Grandy, A. S., Melillo, J. M., & DeAngelis, K. M. (2017). Changes in substrate availability drive carbon cycle response to chronic warming. *Soil Biology and Biochemistry*, 110, 68–78. <https://doi.org/10.1016/j.soilbio.2017.03.002>
- Rasmussen, C., Throckmorton, H., Liles, G., Heckman, K., Meding, S., & Horwath, W. (2018). Controls on soil organic carbon partitioning and stabilization in the California Sierra Nevada. *Soil Systems*, 2(3), 41. <https://doi.org/10.3390/soilsystems2030041>
- Rasmussen, C., & White, D. A. (2010). Vegetation effects on soil organic carbon quality in an arid hyperthermic ecosystem. *Soil Science*, 175(9), 438–446. <https://doi.org/10.1097/SS.0b013e3181f38400>
- Riggs, C. E., Hobbie, S. E., Bach, E. M., Hofmockel, K. S., & Kazanski, C. E. (2015). Nitrogen addition changes grassland soil organic matter decomposition. *Biogeochemistry*, 125(2), 203–219. <https://doi.org/10.1007/s10533-015-0123-2>
- Santos, R. S., Oliveira, F. C. C., Ferreira, G. W. D., Ferreira, M. A., Araújo, E. F., & Silva, I. R. (2020). Carbon and nitrogen dynamics in soil organic matter fractions following eucalypt afforestation in southern Brazilian grasslands (Pampas). *Agriculture, Ecosystems & Environment*, 301, 106979. <https://doi.org/10.1016/j.agee.2020.106979>
- Santos, R. S., Wiesmeier, M., Oliveira, D. M. S., Locatelli, J. L., Barreto, M. S. C., Demattê, J. A. M., & Cerri, C. E. P. (2022). Conversion of Brazilian savannah to agricultural land affects quantity and quality of labile soil organic matter. *Geoderma*, 406, 115509. <https://doi.org/10.1016/j.geoderma.2021.115509>
- Schmidt, M. W. I., & Kögel-Knabner, I. (2002). Organic matter in particle-size fractions from A and B horizons of a Haplic Alisol: Organic matter in particle-size fractions. *European Journal of Soil Science*, 53(3), 383–391. <https://doi.org/10.1046/j.1365-2389.2002.00460.x>
- Schrumpf, M., Kaiser, K., Guggenberger, G., Persson, T., Kögel-Knabner, I., & Schulze, E.-D. (2013). Storage and stability of organic carbon in soils as related to depth, occlusion within aggregates, and attachment to minerals. *Biogeosciences*, 10(3), 1675–1691. <https://doi.org/10.5194/bg-10-1675-2013>
- Schulten, H.-R., Leinweber, P., & Sorge, C. (1993). Composition of organic matter in particle-size fractions of an agricultural soil. *Journal of Soil Science*, 44(4), 677–691. <https://doi.org/10.1111/j.1365-2389.1993.tb02332.x>
- Schuur, E. A. G., Chadwick, O. A., & Matson, P. A. (2001). Carbon cycling and soil carbon storage in Mesic to wet Hawaiian montane forests. *Ecology*, 82(11), 3182–3196. [https://doi.org/10.1890/0012-9658\(2001\)082\[3182:CCASCS\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2001)082[3182:CCASCS]2.0.CO;2)
- Sleutel, S., De Neve, S., Singier, B., & Hofman, G. (2006). Organic C levels in intensively managed arable soils—Long-term regional trends and characterization of fractions. *Soil Use and Management*, 22(2), 188–196. <https://doi.org/10.1111/j.1475-2743.2006.00019.x>
- Sollins, P., Kramer, M. G., Swanston, C., Lajtha, K., Filley, T., Aufdenkampe, A. K., Wagai, R., & Bowden, R. D. (2009). Sequential density fractionation across soils of contrasting mineralogy: Evidence for both microbial- and mineral-controlled soil organic matter stabilization. *Biogeochemistry*, 96(1–3), 209–231. <https://doi.org/10.1007/s10533-009-9359-z>
- Sollins, P., Swanston, C., Kleber, M., Filley, T., Kramer, M., Crow, S., Caldwell, B. A., Lajtha, K., & Bowden, R. (2006). Organic C and N stabilization in a forest soil: Evidence from sequential density fractionation. *Soil Biology and Biochemistry*, 38(11), 3313–3324. <https://doi.org/10.1016/j.soilbio.2006.04.014>

- Soong, J. L., Janssens, I. A., Grau, O., Margalef, O., Stahl, C., Van Langenhove, L., Urbina, I., Chave, J., Dourdain, A., Ferry, B., Freycon, V., Herault, B., Sardans, J., Peñuelas, J., & Verbruggen, E. (2020). Soil properties explain tree growth and mortality, but not biomass, across phosphorus-depleted tropical forests. *Scientific Reports*, 10(1), 2302. <https://doi.org/10.1038/s41598-020-58913-8>
- Sousa, R. F. D., Brasil, E. P. F., Figueiredo, C. C. D., & Leandro, W. M. (2015). Soil organic matter fractions in preserved and disturbed wetlands of the cerrado biome. *Revista Brasileira de Ciência do Solo*, 39(2), 222–231. <https://doi.org/10.1590/01000683rbc20150048>
- Steffens, M., Kölbl, A., & Kögel-Knabner, I. (2009). Alteration of soil organic matter pools and aggregation in semi-arid steppe topsoils as driven by organic matter input. *European Journal of Soil Science*, 60(2), 198–212. <https://doi.org/10.1111/j.1365-2389.2008.01104.x>
- Sun, R., Sun, B., Li, X., Wei, S., Zhao, M., Chu, X., Song, W., Yuan, J., & Han, G. (2023). Seven-year experimental warming decreases labile but not recalcitrant soil organic carbon fractions in a coastal wetland. *Journal of Soils and Sediments*, 23(8), 3071–3081. <https://doi.org/10.1007/s11368-023-03536-5>
- Tian, J., McCormack, L., Wang, J., Guo, D., Wang, Q., Zhang, X., Yu, G., Blagodatskaya, E., & Kuzyakov, Y. (2015). Linkages between the soil organic matter fractions and the microbial metabolic functional diversity within a broad-leaved Korean pine forest. *European Journal of Soil Biology*, 66, 57–64. <https://doi.org/10.1016/j.ejsobi.2014.12.001>
- Tian, Q., Wang, D., Li, D., Huang, L., Wang, M., Liao, C., & Liu, F. (2020). Variation of soil carbon accumulation across a topographic gradient in a humid subtropical mountain forest. *Biogeochemistry*, 149(3), 337–354. <https://doi.org/10.1007/s10533-020-00679-2>
- Trumbore, S. E. (1993). Comparison of carbon dynamics in tropical and temperate soils using radiocarbon measurements. *Global Biogeochemical Cycles*, 7(2), 275–290. <https://doi.org/10.1029/93GB00468>
- van der Pol, L. K., Robertson, A., Schipanski, M., Calderon, F. J., Wallenstein, M. D., & Cotrufo, M. F. (2022). Addressing the soil carbon dilemma: Legumes in intensified rotations regenerate soil carbon while maintaining yields in semi-arid dryland wheat farms. *Agriculture, Ecosystems & Environment*, 330, 107906. <https://doi.org/10.1016/j.agee.2022.107906>
- Verret, M., Wang, Y., Bjornson, J., & Lacelle, D. (2019). Hummocks in alpine tundra, northern British Columbia, Canada: Distribution, morphology and organic carbon composition. *Arctic Science*, 5(3), 127–147. <https://doi.org/10.1139/as-2018-0021>
- Wagai, R., Kajiura, M., Asano, M., & Hiradate, S. (2015). Nature of soil organo-mineral assemblage examined by sequential density fractionation with and without sonication: Is allophanic soil different? *Geoderma*, 241–242, 295–305. <https://doi.org/10.1016/j.geoderma.2014.11.028>
- Wang, Q., Zhang, P.-J., Liu, M., & Deng, Z.-W. (2014). Mineral-associated organic carbon and black carbon in restored wetlands. *Soil Biology and Biochemistry*, 75, 300–309. <https://doi.org/10.1016/j.soilbio.2014.04.025>
- Wood, S. A., Sokol, N., Bell, C. W., Bradford, M. A., Naeem, S., Wallenstein, M. D., & Palm, C. A. (2016). Opposing effects of different soil organic matter fractions on crop yields. *Ecological Applications*, 26(7), 2072–2085. <https://doi.org/10.1890/16-0024.1>
- Xu, C., Guo, L., Dou, F., & Ping, C.-L. (2009). Potential DOC production from size-fractionated Arctic tundra soils. *Cold Regions Science and Technology*, 55(1), 141–150. <https://doi.org/10.1016/j.coldregions.2008.08.001>
- Xu, C., Guo, L., Ping, C.-L., & White, D. M. (2009). Chemical and isotopic characterization of size-fractionated organic matter from cryoturbated tundra soils, northern Alaska. *Journal of Geophysical Research: Biogeosciences*, 114(G3), G03002. <https://doi.org/10.1029/2008JG000846>
- Yang, W., Zhao, H., Leng, X., Cheng, X., & An, S. (2017). Soil organic carbon and nitrogen dynamics following *Spartina alterniflora* invasion in a coastal wetland of eastern China. *Catena*, 156, 281–289. <https://doi.org/10.1016/j.catena.2017.03.021>

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Hansen, P. M., Even, R., King, A. E., Lavalley, J., Schipanski, M., & Cotrufo, M. F. (2023). Distinct, direct and climate-mediated environmental controls on global particulate and mineral-associated organic carbon storage. *Global Change Biology*, 30, e17080. <https://doi.org/10.1111/gcb.17080>

Distinct, direct and climate-mediated environmental controls on global particulate and mineral-associated organic carbon storage

Hansen, Paige M.; Even, Rebecca; King, Alison E.; Lavallee, Jocelyn;
Schipanski, Meagan; Cotrufo, M. Francesca

01 mingxi zhang Page 9
15/1/2024 2:02

02 mingxi zhang Page 9
15/1/2024 2:02