

# A conceptual model of organo-mineral interactions in soils: self-assembly of organic molecular fragments into zonal structures on mineral surfaces

M. Kleber · P. Sollins · R. Sutton

Received: 11 March 2006 / Accepted: 27 November 2006 / Published online: 15 June 2007  
© Springer Science+Business Media B.V. 2007

**Abstract** In this paper, we propose a structure for organo-mineral associations in soils based on recent insights concerning the molecular structure of soil organic matter (SOM), and on extensive published evidence from empirical studies of organo-mineral interfaces. Our conceptual model assumes that SOM consists of a heterogeneous mixture of compounds that display a range of amphiphilic or surfactant-like properties, and are capable of self-organization in aqueous solution. An extension of this self-organizational behavior in solution, we suggest that SOM sorbs to mineral surfaces in a discrete zonal sequence. In the *contact zone*, the formation of particularly strong organo-mineral associations appears to be favored by situations where either (i) polar organic functional groups of amphiphiles interact via ligand exchange with singly coordinated mineral

hydroxyls, forming stable inner-sphere complexes, or (ii) proteinaceous materials unfold upon adsorption, thus increasing adhesive strength by adding hydrophobic interactions to electrostatic binding. Entropic considerations dictate that exposed hydrophobic portions of amphiphilic molecules adsorbed directly to mineral surfaces be shielded from the polar aqueous phase through association with hydrophobic moieties of other amphiphilic molecules. This process can create a membrane-like bilayer containing a *hydrophobic zone*, whose components may exchange more easily with the surrounding soil solution than those in the contact zone, but which are still retained with considerable force. Sorbed to the hydrophilic exterior of hemimicellar coatings, or to adsorbed proteins, are organic molecules forming an outer region, or *kinetic zone*, that is loosely retained by cation bridging, hydrogen bonding, and other interactions. Organic material in the kinetic zone may experience high exchange rates with the surrounding soil solution, leading to short residence times for individual molecular fragments. The thickness of this outer region would depend more on input than on the availability of binding sites, and would largely be controlled by exchange kinetics. Movement of organics into and out of this outer region can thus be viewed as similar to a phase-partitioning process. The zonal concept of organo-mineral interactions presented here

M. Kleber (✉)  
Earth Sciences Division, Lawrence Berkeley National  
Laboratory, One Cyclotron Road, Berkeley, CA,  
94720, USA  
e-mail: mkleber@lbl.gov

P. Sollins  
Department of Forest Science, Oregon State  
University, Corvallis, OR, 97331, USA

R. Sutton  
Department of Environmental Science,  
Policy and Management, University of California,  
#3114 Mulford Hall, Berkeley, CA, 94720-3114, USA

offers a new basis for understanding and predicting the retention of organic compounds, including contaminants, in soils and sediments.

**Keywords** Multilayer · Adsorption · Soil organic matter · Nitrogen · Protein · Amphiphilic · Organo-mineral · Micelle · Humic

## Introduction


Decades ago, earth scientists acknowledged the ability of mineral particles to protect soil organic matter (SOM) from biological attack (Jung 1943; Allison et al. 1949). In temperate, cultivated soils, 50–75% of SOM exists within clay-sized organo-mineral particles (Christensen 2001), and numerous researchers have reported positive correlations between the contents of fine mineral particles and carbon in soils (Körschens 1980; Nichols 1984; Burke et al. 1989; Mayer and Xing 2001). As a consequence, it is widely assumed that the inhibiting effect of soil minerals on decomposition can influence strongly the turnover of SOM (Martin and Haider 1986; Theng et al. 1992; Hsieh 1996; Parfitt et al. 1997). It follows that understanding SOM stabilization requires knowledge of the processes operating at the organo-mineral interface. Of particular importance but great complexity are the questions of how strongly SOM may bind to mineral surfaces, how such bonds form and eventually break, and how the mechanisms of attachment of SOM to mineral surfaces affect its residence time in soils.

While the characteristics of mineral surfaces are reasonably well known, the chemical properties of SOM continue to be the subject of much debate. After all, while minerals crystallize towards or dissolve from a defined and comparatively simple crystal structure, SOM encompasses organic molecules representing both compounds released from living plant and microbial cells (e.g., extracellular enzymes, surface-active proteins, chelating compounds, and many others) and complex plant, microbial and animal residues in various stages of alteration due to both biotic and abiotic processes (Baldock and Skjemstad 2000). Considerable spectroscopic evidence (Chien et al. 1997; Engebretson and von

Wandruszka 1997, 1998; von Wandruszka 1998; von Wandruszka et al. 1999; Ferreria et al. 2001; Martin-Neto et al. 2001; von Wandruszka and Engebretson 2001; Nanny and Kontas 2002) indicates that soluble mixtures of organic molecules representing a significant fraction of SOM can form organized structures called micelles within aqueous solution, structures that consist of hydrophilic exterior regions that shield hydrophobic interiors from contact with water molecules (von Wandruszka 1998). Because amphiphilic molecules are requisite to the formation of micelles, the ability of SOM components to form these structures suggests that many of these molecules are amphiphilic. Significant to this insight, Wershaw (1993) previously developed a bilayer model of organo-mineral interactions (Fig. 1) that sharply contrasted with the traditional view of organo-mineral interactions (Stevenson 1985), which were visualized as associations of large, multifunctional polymers with mineral surfaces via a broad range of bonding mechanisms (Stevenson 1985; Leinweber and Schulten 1998). Further, Wershaw and Pinckney (1980) postulated that decayed organic materials are often bound to clay surfaces by amino acids or proteins, based on the observation that deamination of organo-mineral complexes with nitrous acid released organic materials from the clay.

It is our intention to develop this earlier work into a generally applicable concept of the zonal structure of organo-mineral associations (Fig. 2), based on the amphiphilicity of SOM fragments, and the intimate involvement of proteinaceous compounds in stable organo-mineral associations. As defined here, a *zonal structure* is formed when the organic matter attached to a mineral surface is segregated into more than one layer or zone of molecules, such that not all adsorbed molecules are in contact with the mineral surface. Assuming such a zonal structure, we are able to account simultaneously for a number of phenomena observed in soils and sediments, including:

- a) The decrease of C/N ratio with decreasing particle size (Hedges and Keil 1995; Aufdenkampe et al. 2001);
- b) The decrease of C/N ratio with increasing particle density (Sollins et al. 2006);



**Fig. 1** The “Wershaw bilayer model” of organo-mineral interactions (Wershaw et al. 1996a). Figure adapted from Essington (2003). Accumulation of amphiphilic fragments on soil surfaces is initiated by electrostatic interactions between


hydrophilic organic moieties and surface oxygens or hydroxylated surface-metal cations. The hydrophobic portions of these surface molecules are shielded from the polar aqueous phase by a second layer of amphiphiles, forming a bilayer

- c) The decrease of  $^{14}\text{C}$  content with decreasing C content in heavy fractions (Sollins et al. 2006);
- d) The preferential sorption of large, hydrophobic organic compounds in sorption experiments with soils (Jardine et al. 1989; Guo and Chorover 2003) and in marine/estuarine sediments (Zhou et al. 1994);
- e) The existence of at least two different C and N pools in heavy fractions: an older, more stable pool of C and N, and a more recent, fast cycling pool of C and N (Trumbore et al. 1989; Strickland et al. 1992; Swanston et al. 2005);
- f) The observation that mineral-associated organic matter accumulates in A horizons beyond levels consistent with monolayer coverage (Eusterhues et al. 2005);
- g) The correlation of concentrations of poorly crystalline mineral phases with amounts of oxidation-resistant SOM (Kleber et al. 2005);
- h) Seasonal changes in water repellency in soils (Horne and McIntosh 2000).

To support the concept of zonal structure of organo-mineral complexes, we first discuss the idea of SOM as consisting of associations of organic molecules capable of micellar self-organization in aqueous solution. Given this important concept, it is appropriate to view a significant portion of SOM as consisting of a mixture of diverse but typically amphiphilic molecules. We will present empirical evidence that supports the idea of a zonal structure of organic accretions on mineral surfaces, demonstrate the special importance of proteinaceous compounds for the architecture of this structure, and show that the assumption of different zones of sorption implies differential exchange kinetics for organic fragments moving into and out of individual zones.

### Amphiphilic molecules and the micellar association model of soil organic matter

Critical insights regarding organo-mineral complexes arose from studies probing the behavior of



**Fig. 2** The zonal model of organo-mineral interactions. The variety of surface types in soils is represented by a low-charge smectite, a hydroxylated Fe-oxide coating, and a hydrophobic kaolinite siloxane surface. This selection is meant to illustrate major bonding mechanisms, and is not intended to represent the full range of mineral surface types that may be encountered in soils. In the *contact zone*, amphiphilic fragments accumulate on charged surfaces through electrostatic interactions, directing hydrophobic portions outwards toward the polar aqueous solution. Proteins serve as a surface conditioner that adds polar functionality to low-charge siloxane surfaces. Some hydrophobic organic compounds may also associate with noncharged mineral surfaces. The *zone of hydrophobic interactions* is equivalent to the hydrophobic region formed as part of the Wershaw bilayer model, and results from the entropically driven shielding of the hydrophobic portions

of adsorbed organic molecules from the polar aqueous phase by a second layer of amphiphilic molecules. The hydrophobic zone is thought to be discontinuous, as some proteins directly adsorbed to mineral surfaces may well expose ionized or polar functional groups towards the soil solution. In the outer region or *kinetic zone*, further accumulation of organic molecular fragments is possible, probably largely mediated by the presence of multivalent cations. Depending on stereochemical orientation, degree of amphiphilicity, cation content of the soil solution, pH, temperature, and perhaps several other controlling factors, a variety of modes of attachment of organic molecular fragments to the intermediate region appears to be possible. The thickness of the outer zone is likely to depend on input of organic materials and thus to be largely controlled by exchange kinetics

natural organic fractions in aqueous solution. To examine and manipulate the behavior of these complex organic materials within a controlled laboratory environment, alkaline extraction is often used to solubilize a large portion of SOM, producing an operationally defined humic fraction. That alkaline extraction effectively removes between 50% (Rice 2001) and 80% (Scheffer and Schachtschabel 2002) of the organic material in soil samples indicates that while the humic fraction represents only a portion of SOM, it is a

significant portion of this organic material, and it contains many reactive polar or charged moieties that respond to pH. In addition, behavior consistent with micellar organization, observed using a diverse array of spectroscopic techniques including fluorescence (Engbreton and von Wandruszka 1997, 1998; von Wandruszka 1998; von Wandruszka et al. 1999; von Wandruszka and Engbreton 2001; Nanny and Kontas 2002), electron spin resonance (Ferreria et al. 2001; Martin-Neto et al. 2001), and nuclear magnetic resonance

(Chien et al. 1997), suggests that nonpolar moieties are also present within humic fractions. After all, it is the presence of hydrophobic functional groups within amphiphilic molecules that triggers entropically-driven self-organization to shield these nonpolar moieties from the polar aqueous medium through formation of a micelle. Thus, the observation of micellar structure within humic extracts representing a major portion of SOM indicates that a significant fraction of the molecules making up this complex organic material are amphiphilic in nature.

The components of decomposing SOM display a range of functional qualities based on chemical and structural composition (Sutton and Sposito 2005). We suggest that these chemical properties represent a continuum of amphiphilicity that ranges from molecules that are solely nonpolar and hydrophobic, to those that are predominantly amphiphilic because they also contain hydrophilic, highly polar or charged functional groups. Nonpolar, hydrophobic compounds are composed predominantly of alkyl and aromatic functional groups, and may derive from sources such as plant waxes and cutins. Occupying the mid-range of this chemical continuum are mildly polar materials, such as many carbohydrates and their derivatives, which feature functional groups like alcohols or ethers that do not ionize under typical soil and water pH conditions. In addition to hydrophobic functional groups, the amphiphilic compounds that populate the other extreme of this chemical spectrum contain more reactive functional groups like carboxyls and amines, which can develop charge under typical soil and water pH conditions. The dominant role of amphiphilic molecules in the behavior of complex organic mixtures within aqueous environments cannot be ignored when considering the mechanisms by which SOM adsorbs to mineral surfaces.

### Experimental evidence in support of a zonal structure of organic accretions on mineral surfaces

While mounting experimental evidence (Chien et al. 1997; Engebretson and von Wandruszka

1997, 1998; von Wandruszka 1998; von Wandruszka et al. 1999; Ferreria et al. 2001; Martin-Neto et al. 2001; von Wandruszka and Engebretson 2001; Nanny and Kontas 2002) supports the ability of complex natural organic substances to self-assemble into associations featuring regions with different chemical properties, similar behavior is not typically ascribed to SOM adsorbed to mineral surfaces. However, starting with the previously mentioned work by Wershaw and colleagues (Wershaw and Pinckney 1980; Wershaw 1986; Wershaw 1993), information from a broad range of scientific fields has provided numerous clues suggesting that SOM sorbs to minerals in layers or zones that may have different chemical or functional properties. In the following sections we use (i) the fate of organic pollutants in soils, sediments, and aquifers, (ii) the distribution of organic matter over mineral surfaces in natural systems, (iii) information from density fractionation experiments, (iv) evidence from the study of water repellency in soils, and finally, (v) data obtained through laboratory synthesis of layered organo-mineral structures, to draw inferences suggesting that organic accretions on minerals consist of multiple zones of self-assembling materials.

Sorption of organic compounds to organic coatings on mineral surfaces as a model of zonal organo-mineral architecture

The idea that organic matter covers mineral surfaces in a layered or zonal fashion (Fig. 2) is not new (Chassin 1979), with most early concepts of organics sorbing to mineral surfaces in two or more layers resulting from a wealth of studies on the fate of organic pollutants in soils, sediments, and aquifers. A review of this early work led Voice and Weber (1983) to conclude that, although hydrophobic bonding was a major sorption mechanism for hydrophobic compounds, no single sorption mechanism adequately explained all experimental observations. Instead, Weber et al. (1983) state that “*the organic carbon content of a solid appears to be the major factor in determining its sorptive capacity.*”

A step forward in advancing mechanistic knowledge about zonal arrangements of organic

molecules on mineral surfaces was taken by Murphy et al. (1990), who, following up on the work of Hunter and Liss (1982) among others, found that hydrophilic mineral particles exposed to humic fractions developed hydrophobic surfaces, rendering them more capable of sorbing hydrophobic organic compounds. Hydrophobic interactions bind the outer layer hydrophobic compounds to the inner layer organics because the total hydrocarbon surface that is exposed to water is reduced, and entropy maximized, when hydrophobic entities organize to reduce the non-polar surface area exposed to water molecules. Jardine et al. (1989) provided additional support for the importance of hydrophobic interactions to adsorption of organic materials by comparing the sorption of dissolved organic carbon solutions that had been fractionated into operationally defined hydrophobic and hydrophilic organic portions. Hydrophobic organic solutes were preferentially adsorbed by soils relative to hydrophilic organic solutes, leading the authors to conclude that hydrophobic interactions driven by favorable entropy changes represent the primary mechanism of adsorption for these organic solutes. Murphy et al. (1990) further showed that the nature of the mineral surface, particularly the concentration and spatial distribution of hydroxyl sites, influenced the amount of hydrophobic organic C adsorbed to organic coatings. The chemical properties of the humic fraction used to coat the mineral particles influenced the amount of hydrophobic compounds adsorbed, with the most aromatic humic material being the strongest sorbent for hydrophobic material.

Patchy organic matter distributions on mineral surfaces indicate zonal architecture of sorbed materials

Further evidence for the zonal or layered sorption of organic matter to mineral surfaces comes from the examination of natural organo-mineral complexes taken from soil and marine systems. Both specific surface area (SSA) determinations and microscopic analyses indicate that the distribution of organic materials on the mineral surfaces of soils (Mayer and Xing 2001; Kahle

et al. 2002) and marine sediments (Ransom et al. 1998; Mayer 1999) is typically discontinuous. While organic matter loadings on sedimentary mineral surfaces are often on the order of the so-called monolayer equivalent of 0.5–1.0 mg organic C m<sup>-2</sup> (Keil et al. 1994; Mayer 1994), the fact that organic material is not distributed evenly over the available mineral surface indicates that the material must instead be clustered in small patches with some vertical extension. Aluminosilicate sediments with low to moderate loadings of organic matter (<3 mg organic C m<sup>-2</sup>) generally have less than 15% of their surfaces coated, with organic materials existing in discrete spots on the mineral surfaces (Arnarson and Keil 2001).

Speculations about a layered architecture of these patches have been voiced by Kaiser and Guggenberger (2003), who found that the reduction of SSA following sorption of increasing amounts of DOC was not uniform: the decrease in SSA was larger at smaller organic matter loadings than at larger ones. Among other potential explanations, they hypothesized that “*with increasing surface loading an increasing portion of the sorbing molecules did not attach to mineral surfaces but formed organic multilayers as a result of hydrophobic interactions or bridging by polyvalent cations between organic ligands of sorbed and soluted molecules.*” This interpretation was subsequently supported by the work of Wang and Xing (2005), who found that, even under different organic matter loading levels, mineral surface coverage remained nearly constant, indicating that “*the coating must have increased in thickness while retaining practically the same surface coverage at a higher loading. In such a multilayer arrangement, the first several molecular layers close to the mineral surface may take a more compacted form due to the attractive forces of the mineral surface.*” In addition, ocean sediment studies show a direct relationship between organic matter content and mineral surface area at organic loadings 2–5 times that of a monolayer equivalent (Hedges and Keil 1995), further supporting the existence of minerals coated with multiple layers of organic molecules.



Organo-mineral associations isolated through density fractionation may also suggest a zonal architecture of sorbed materials

In addition to the ocean sediments described above (Hedges and Keil 1995), soil particles isolated through density fractionation have also provided evidence indicating that organic material is sorbed to minerals through both direct and indirect interactions with mineral surfaces. Eusterhues et al. (2005) found that organic matter loadings in heavy fractions ( $>2 \text{ g cm}^{-3}$ ) from a series of topsoils were greater than could be explained solely by direct attachment to the available mineral surfaces area in a monolayer fashion. They hypothesized that interactions between organic molecules leading to greater than monolayer coverage would be necessary to explain the large amounts of sorbed organic material (Eusterhues et al. 2005).

Organic C in the heavy fraction of soils is commonly seen as being closely associated with minerals and typically more depleted in  $^{14}\text{C}$  than light-fraction material, which is not intimately bound to minerals (Trumbore and Zheng 1996). However, when Swanston et al. (2005) traced an atmospheric  $^{14}\text{C}$  pulse into a forest ecosystem, they noticed that the elevated  $^{14}\text{C}$  signal appeared not only in the light fraction, but also in the strongly mineral-associated dense fraction. They interpreted this observation as indicating the existence of at least two different C pools in the organo-mineral fraction: an older, more stable pool of C, and a younger, fast cycling C pool. Strickland et al. (1992) found similarly that  $^{15}\text{N}$ -labeled inorganic N was incorporated rapidly into dense (organo-mineral) fractions of five soils of greatly differing mineralogy, then released as both inorganic and organic N during a subsequent incubation. The existence of rapidly cycling C and N in the dense or mineral-associated fraction can satisfactorily be explained by a zonal structure of organic accretions on mineral surfaces. While the organic materials directly adsorbed to the mineral surface are likely to be tightly bound and well protected from microbial degradation, the materials sorbed in outer regions of the organo-mineral complex, and lacking direct interaction with the mineral surface, are presumably more loosely

bound and more readily desorbed or decomposed. That only part of the mineral-bound carbon is protected from oxidation is corroborated by experiments that subject subsoils to chemical oxidation. The resulting C solubilization and mineral dissolution show that most of the organic matter in subsoils is associated with mineral surfaces (Eusterhues et al. 2003; Mikutta et al. 2006). Thus mild chemical oxidation leaves an organic residue with a mass that correlates significantly with mineral surface  $\text{F}^-$  reactivity, and a  $\Delta^{14}\text{C}$  value lower than that of the original organic matter pool (Kleber et al. 2005).

Seasonal water repellency explained through presence of organic material sorbed to mineral surfaces in a zonal fashion

Seasonal changes in soil water repellency may also be explained by alterations to the zonal architecture of sorbed organic matter induced by seasonal changes in the soil environment. Non-seasonal water repellency in soils has often been attributed to the presence of hydrophobic moieties such as those found in long-chain fatty acids (Ma'shum et al. 1988). The presence of hydrophobic compounds alone, however, does not explain *seasonal* changes in water repellency. Using the 'molarity of ethanol droplet' method to measure repellency, combined with a set of organic extractions, Horne and McIntosh (2000) attributed seasonal changes in soil wettability to the existence of multiple, functionally different layers of organic matter on sandy grains. In their model, an inner layer of more hydrophobic compounds is covered by a layer of amphiphilic compounds. When the soil is wettable, the hydrophobic material is effectively screened, especially if the outer surface is well hydrated. Repellence occurs when the hydrophobic compounds making up the inner layer are more exposed. This can be linked to seasonal moisture differences through a combination of factors: during the wet season, the outer layer will be well hydrated, amphiphilic compounds will be arranged in such a fashion as to have their polar portions oriented outwards, the carboxylate anion will predominate, and hydrophobic groups will be effectively screened by more hydrophilic groups. With the onset of the dry season, a more

hydrophobic outer surface develops as the amphiphilic compounds change in orientation, the carboxylate groups are protonated, and contraction of the amphiphilic ‘surface screen’ exposes the interior hydrophobic layer.

#### Laboratory and industrial synthesis of multilayer organic assemblages

Multilayered organo-mineral structures are routinely manufactured and studied in the laboratory, mainly for biomedical (Galeska et al. 2001) and biotechnical purposes (Halthur et al. 2004). Because these artificial organo-mineral structures likely form as a result of some of the same adsorption mechanisms active in soils, they may be viewed as potential analogs of natural organo-mineral complexes. In one such study of artificial organo-mineral assemblages, Spaeth et al. (1997) sorbed protein multilayers to hydrophobic silanized silicon wafers using alternating incubations with poly-streptavidin and bovine serum albumin. They found that the thickness of the organic layers increased 18.75 nm per incubation, yielding a final protein multilayer thickness of 350 nm.

Galeska et al. (2001) sorbed layers of the Aldrich commercial humic fraction onto silicon wafers, noting that the expansion of the organic film could be significantly altered by the incorporation of salt, with the charge screening effect contributing to thicker film deposition irrespective of pH. Further experimentation indicated that the permeability of simple carbohydrates through the humic coating was a function of the number of self-assembled layers of organic material. That the probe glucose penetrated thick films composed of many layers of organic matter more readily than thin films made up of few layers suggests that organic molecules, including both SOM and xenobiotic substances, may not be tightly bound to outer regions of organo-mineral complexes.

Alkylated silicon oxide surfaces were used by Schmidt et al. (1990) in studies of protein adsorption on hydrophobic surfaces. They found that a model with at least three classes of adsorbed molecules was necessary to adequately describe their experimental results. A first layer was formed by the molecules that adsorbed within a short time

after the beginning of the incubation. These molecules made up approximately 65% of the final coverage. They were irreversibly adsorbed and did not measurably exchange with bulk solution molecules. A second layer reached equilibrium after several hours of incubation and showed a pronounced and temperature-dependent exchange with bulk molecules. A third layer consisted of molecules that exchanged rapidly with the bulk solution, but comprised only about 10% of the total coverage. An important implication of this work is that a protein-based multilayer structure allows occurrence of both hydrophobic adsorption on a stable inner layer, as well as phase-partitioning into a dynamic outer layer, thus explaining experimental results suggesting the coexistence of both kinds of sorption mechanisms in the same system.

#### Reconciling the evidence for zonal structures on mineral surfaces

The evidence compiled above suggests that SOM may attach to mineral surfaces in a zonal or layered fashion. In conjunction with the work of Wershaw et al. (1995; 1996a; b) on adsorption of compost leachates to aluminum oxide, studies of the amounts or spatial distributions of organic materials bound to mineral particles, and of variations in C turnover time and seasonal water repellency, have provided a wealth of direct and indirect information supporting the concept of an organized, zonal architecture for SOM sorbed to mineral surfaces. Experiments and observations in the laboratory prove that organic multilayers on mineral surfaces can even be manufactured to serve specific purposes. Ecologically speaking, the work of Schmidt et al. (1990) is especially significant, as it assigns different exchange kinetics to molecules making up different parts of the organo-mineral complex, with molecules of the inner zone being the most stable, and molecules of the outermost zone having the fastest exchange with the soil solution. Such differential, zone-dependent kinetics could explain why organo-mineral isolates from soils can contain C and N pools with different turnover times (Trumbore et al. 1989; Strickland et al. 1992; Swanston et al. 2005).



**Table 1** Relative composition of vascular plants, algae, bacteria, and fungi, compiled using data from Knicker (2004) and White (1997)

	Vascular plants	Algae	Bacteria	Fungi
	% Dry matter			
Lignin	5–30			
Cellulose	15–60			
Hemicellulose	10–30			
N-containing compounds	2–15	24–50	50–60	14–52
Lipids		2–10	10–35	1–42
Carbohydrates		40	4–32	8–60
	Ratio			
C:N ratio	20–50 (tree leaves) 25–80 (herbaceous plants)	6	5–8	≈10

### The role of proteins in organo-mineral associations

Mineral-associated SOM exhibits C/N ratios of 7–14, whereas water-extracted SOM has C/N ratios of 26–55 (Aufdenkampe et al. 2001), indicating N enrichment of adsorbed organic materials. Numerous observations, made in both soils and marine/freshwater environments (Aufdenkampe et al. 2001), similarly suggest that the adsorption of nitrogenous compounds to mineral surfaces may be a broadly relevant ecological process. Thus, a model of organo-mineral associations should account for the body of evidence suggesting preferential adsorption of N-containing moieties over other organic compounds to mineral surfaces (Omoike and Chorover 2006; Hedges and Oades 1997).

Certainly, the relatively high strength of protein adsorption to mineral surfaces (Theng 1979; Chevallier et al. 2003; Wershaw 2004) is a long-established (Ensminger and Gieseking 1942)—and often technologically exploited (Gougeon et al. 2003)—chemical phenomenon, and one that has proved relevant to the laboratory synthesis of multilayer organo-mineral complexes, as discussed previously (Schmidt et al. 1990). We suggest that proteinaceous materials play a prominent role in the formation and structure of organo-mineral complexes due to both the relatively high abundance of such compounds in soils, and the ability of these materials to adsorb irreversibly to mineral surfaces.

*Protein availability at soil mineral surfaces*— $^{15}\text{N}$ -NMR analysis has demonstrated that a significant amount of soil nitrogen is present in amide form

(Almendros et al. 1991; Knicker et al. 1995; Knicker and Lüdemann 1995; Knicker 2000, 2002; Leinweber and Schulten 2000; Mahieu et al. 2000, 2002; Mathers et al. 2000; Zang et al. 2000), and thus most likely in the form of proteins or protein residues. Tremblay and Brenner (2006) showed in a 4-year decomposition experiment that on average 60–75% of the N in highly decomposed detritus was derived from heterotrophic bacteria, rather than plant tissues. As the smallest organisms in the soil, bacteria and fungi are able to enter small, confined pore spaces and die and decompose there, and also contain far more N than vascular plant materials or surface litter (Table 1). In fact, bacteria are known to actively use proteins to attach to surfaces (Bashan and Levanony 1988; Dufrene et al. 1999). Mycelial fungi also secrete a broad class of surface-active glycoproteins (Wright and Upadhyaya 1996). Abundant evidence indicates that microbial products accumulate in fine clay fractions of soils (Leinweber and Schulten 1995; Kleber et al. 2004; Mertz et al. 2005), further corroborating the supposition that the more proteinaceous microbial residues are likely to make initial contact with mineral surfaces, rather than the less proteinaceous plant residues.

*Protein polyfunctionality and adsorption to mineral surfaces*—Proteins are a special class of amphiphilic molecules with a strong tendency to bind to surfaces and to resist desorption (Hlady and Buijs 1996). They consist of chains of amino acids with both apolar and polar (either neutral or charged) side chains, folded into a conformation in which as many apolar residues as possible are packed into the interior, leaving mainly polar and charged residues at the surface.

Given the diversity of functional groups making up proteins, no single process dominates protein adsorption to mineral surfaces. Instead, three subprocesses drive adsorption behavior: (i) structural rearrangement of the protein molecule; (ii) dehydration of parts of the sorbent and protein surfaces; and (iii) redistribution of charged groups at the organo-mineral interface (Haynes and Norde 1994). Numerous electrostatic bonds can form between reactive mineral sites and organic amide and other polar or charged functional groups during the adsorption process (Brash and Horbett 1995). Due to their large size and coiled structure, proteins gain considerable conformational energy by unfolding at the solid/liquid interface, an entropically driven rearrangement that produces numerous organo-mineral interactions and distinguishes protein adsorption from that of other available biomolecules (Haynes and Norde 1994). With adsorption to a mineral, proteins may increase the number of reactive sites on the newly coated surface by exposing their abundant polar functional groups to the soil solution. This effect is known as support preconditioning (Dufrene et al. 1996), and is considered a routine step preceding the adhesion of microorganisms to mineral surfaces (Bos et al. 1999; Bakker et al. 2004). The surface charge densities of both protein and surface can vary with pH, and adsorption is typically greatest at the isoelectric point of the protein (Ramsden 1995).

Proteinaceous compounds can bind especially stably to mineral surfaces over long time periods, as illustrated by the fruitless attempts of archaeologists to remove protein from archaeological artifacts for further study (Craig and Collins 2002). Amelung et al. (2006) used changes in chirality of aspartic acid and lysine to show that some soil proteins achieve residence times of centuries. Sorption experiments (Hedges and Hare 1987; Wang and Lee 1993) as well as observational data from marine sediments and riverine particles (Hedges and Hare 1987; Wang and Lee 1993; Mayer et al. 1995; Arnarson and Keil 2001; Aufdenkampe et al. 2001) also indicate strong and stable binding of amino acids to clay minerals. Glycoproteins secreted by arbuscular mycorrhizal fungi (Wright and Upadhyaya 1996) stabilize soil aggregates (Baveye 2006), likely due to their

ability to bind to both hydrophilic and hydrophobic mineral surfaces. Strong adsorption of defined proteins on clean clays is well documented in the laboratory, with the amount of adsorbed material varying over nearly five orders of magnitude with the nature of the sorbate, sorbent, and experimental conditions (Theng 1979; Burchill et al. 1981; Wang and Lee 1993; De Cristofaro and Violante 2001; Ding and Henrichs 2002).

#### N-containing biomolecules as contact materials within a zonal organo-mineral complex

The proposed special role of proteins in the organo-mineral contact zone has implications for the attachment of additional organic compounds, or the *zonal architecture* of organic materials adsorbed to mineral surfaces. Organo-mineral complexes featuring multiple layers of proteins have been synthesized using a variety of experimental methods (Schmidt et al. 1990; Spaeth et al. 1997; Halthur et al. 2004). The amounts of proteins that may eventually sorb to mineral surfaces can be large, approaching 1 g protein g<sup>-1</sup> clay for clean clays (e.g., Burchill et al. 1981; De Cristofaro and Violante 2001) and buried ceramics (Craig and Collins 2002), with record values of 2.4 g g<sup>-1</sup> (Greenland 1965) on montmorillonite, implying a soil C concentration greater than that allowed for a mineral soil horizon according to Soil Taxonomy (Soil Survey Staff 1999).

However, observations from natural systems provide more compelling evidence regarding the critical role proteins play within soil organo-mineral complexes. Physical fractionation data led Sollins et al. (2006) to postulate that sorption of proteins might create a basal layer to which other amphiphilic molecular fragments would adsorb more strongly than they would to clean clay surfaces, creating a zonal effect, with N-rich compounds more abundant near mineral surfaces, and C-rich compounds more abundant farther from the surface. Physical fractionation reveals the tendency for the C/N ratio to decrease with increasing particle density (Turchenek and Oades 1979; Young and Spycher 1979). Lighter fractions (<1.8 g cm<sup>-3</sup>) have high C/N ratios because they are rich in plant structures high in lignin and cellulose. The continued downward trend in C/N

ratios at densities beyond 2.2 and even  $2.4 \text{ g cm}^{-3}$  has previously been attributed to the accumulation of proteinaceous material (Oades 1988), and is consistent with our suggestion that N-rich compounds have a high probability of remaining sorbed to mineral surfaces.

Sollins et al. (2006) further suggested that sequential density fractionation combined with  $^{14}\text{C}$  analysis might offer a way to assess the role of N-rich compounds in the formation of organo-mineral complexes. They pointed out that variation in particle density correlated well with the percentage of C in the particles, and thus that C loadings decreased with increasing particle density, suggesting thinner coatings of organics, and in turn a more protein-rich inner layer. Assuming these thin inner coatings were more stable,  $^{14}\text{C}$  age should increase with particle density. This proved to be true for an Andic forest soil they examined. However, large differences in mineralogy of the density separates from this soil precluded ascribing the increase in  $^{14}\text{C}$  age solely to coating thinness.

### The zonal structure of organo-mineral complexes: a hypothesis

Recent spectroscopic evidence suggests that the decomposing organic residues in soils can be viewed as a mixture of largely amphiphilic fragments (Chien et al. 1997; Engebretson and von Wandruszka 1997, 1998; von Wandruszka 1998; von Wandruszka et al. 1999; Ferreria et al. 2001; Martin-Neto et al. 2001; von Wandruszka and Engebretson 2001; Nanny and Kontas 2002). The amphiphilicity of many of these molecular fragments, and the polarity of the aqueous soil solution in which they are contained, cause them to self-assemble into micellar associations when suspended at high concentrations, and into layered or zonal structures when in contact with mineral surfaces. Based on the evidence presented above, we propose that the self-assembly of SOM on mineral surfaces creates a layered or zonal structure with regular features (Fig. 2).

*The contact zone*—When brought into contact with a mineral surface, amphiphilic SOM compounds form a contact zone whose structural

properties depend mainly on the type, density, and chemical reactivity of the mineral functional groups, and on the pH of the soil solution. Organic fragments directly attached to mineral surfaces are protected against decomposition and do not participate in significant exchange with organic compounds in the soil solution. This can be explained in part by the strong, ligand-exchange bonding of charged or polar heads of amphiphilic molecules to hydroxylated mineral surfaces (Tiberg et al. 1999). Proteins, a special class of amphiphilic molecules, can bind particularly strongly to mineral surfaces because they reinforce the initial organo-mineral bonds that form with entropically favorable conformational changes that produce further opportunities for more numerous adsorption interactions (Haynes and Norde 1994).

*The hydrophobic zone*—Amphiphilic molecules adsorbed directly to mineral surfaces tend to mask the physicochemical properties of these surfaces (Bos et al. 1999), to the extent that an outwardly oriented region of high hydrophobicity may be created, especially when the underlying mineral surface has high densities of reactive, singly coordinated hydroxyls (Murphy et al. 1990). The hydrophobic zone thus formed provides a suitable region for sorption of hydrophobic moieties, including the hydrophobic portions of amphiphilic molecules in the soil solution, leading to formation of a bilayer structure on the surface of the mineral. In contrast, the direct adsorption of proteins to a mineral surface may lead to an increase in the number of reactive surface sites, rather than hydrophobic sites. Direct adsorption of proteins may not result in formation of a hydrophobic zone or a bilayer structure, but instead enable the attachment of SOM components through electrostatic interactions. Thus, the hydrophobic zone must not be considered an essential and continuous layer present within all organo-mineral associations, but rather a zone likely to be found to varying degrees in many of these associations.

*The kinetic zone*—Depending on stereochemical orientation, degree of amphiphilicity, cation content of the soil solution, pH, temperature, and perhaps several other controlling factors, a variety of modes of attachment of further organic

molecules appears to be possible. No common principle governing these associations is apparent. However, there is reason to assume that the organic compounds trapped in this outer region are partitioned rather than sorbed, and that their residence times within the outer region are shorter than the residence times of compounds in the intermediate or inner regions. Because “*Crowded conditions promote selfassembly*” (Goodsell 2004), the thickness of the outer layer is likely to depend more on input of organic materials than on the availability of binding sites, and to be largely controlled by exchange kinetics. This suggests that more material will be sorbed to kinetic zones in topsoil A horizons than in subsurface B and C horizons. Movement of organics into and out of this outer region may thus resemble a phase-partitioning process, rather than classical adsorption.

## Conclusions

With the proposed zonal model, we attempt to reconcile information from different natural and artificial systems and from various scientific fields into an integrated view of the organo-mineral complex and its most important properties. Our model encompasses interdependent hypotheses regarding (i) the formation of organo-mineral associations, (ii) the specific architecture of the zones of adsorbed organic matter, and (iii) the exchange kinetics of these organic materials with the surrounding soil solution.

We suggest a zonal structure for organic matter accretions on mineral surfaces by recognizing that SOM consists largely of amphiphilic molecular fragments with the thermodynamic impetus to self-assemble into micellar structures on mineral particle surfaces suspended in a polar liquid (Fig. 2). We further propose that microbial proteins are better able to bind to mineral surfaces, as well more likely to arrive at these surfaces, than are residues of vascular plants, thus explaining the proposed existence of a particularly stably bound, N-rich inner layer of organic material. Hydrophobic portions of these surface-bound molecules can be protected from contact with aqueous solution by the additional accumulation

of amphiphiles arranged such that the polar portions point towards the aqueous phase, creating a bilayer structure with a discrete hydrophobic zone. Empirical evidence for organic surface loadings of 2–5 times the monolayer equivalent (Hedges and Keil 1995) suggests that further attachment of organics in a third region or kinetic zone is possible, although for this outer zone or region, exchange rates of organic compounds with the soil solution are likely high, and residence times brief.

The arguments in support of the zonal hypothesis of organo-mineral complexes are indirect in nature. We invite the scientific community to join us in future attempts to challenge the zonal concept and to help us further elucidate the structure and dynamics of organo-mineral associations. If experimentally and empirically validated, the zonal model for organo-mineral associations in natural systems should not only open up a new road towards improved understanding of organic C and N dynamics in soils and sediments, but also facilitate the modeling of the fate and transport of contaminants.

**Acknowledgements** We thank M.S. Torn, T. Filley, C. Swanston, M. Kramer, G. Sposito and especially B. Caldwell for discussions and helpful comments on earlier drafts of the manuscript. The comments of J. Chorover were particularly valuable and are gratefully appreciated. The contribution of MK results from previous support through the German Science Foundation (DFG) priority program SPP 1090 and from current financial support by the Climate Change Research Division, Office of Science, U.S. Department of Energy under Contract No. DE-AC03-76SF00098. Partial funding was provided by NSF SGER and USDA CSRI grants to PS and MK and by the Andrews LTER program.

## References

- Allison FE, Sherman MS, Pinck LA (1949) Maintenance of soil organic matter: I. Inorganic soil colloid as a factor in retention of carbon during formation of humus. *Soil Sci* 68:463–478
- Almendros G, Fründ R, Gonzalez-Vila FJ, Haider KM, Knicker H, Lüdemann HD (1991) Analysis of C-13 and N-15 CPMAS NMR-spectra of soil organic-matter and composts. *FEBS Lett* 282:119–121
- Amelung W, Zhang X, Flach KW (2006) Amino acids in grassland soils: Climatic effects on concentrations and chirality. *Geoderma* 130:207–217
- Arnarson TS, Keil RG (2001) Organic-mineral interactions in marine sediments studied using density fraction-

- ation and X-ray photoelectron spectroscopy. *Org Geochem* 32:1401–1415
- Aufdenkampe A, Hedges J, Richey J, Krusche A, Llerena C (2001) Sorptive fractionation of dissolved organic nitrogen and amino acids onto fine sediments within the Amazon Basin. *Limn Ocean* 46:1921–1935
- Bakker DP, Busscher HJ, van Zanten J, de Vries J, Klijstra JW, van der Mei HC (2004) Multiple linear regression analysis of bacterial deposition to polyurethane coatings after conditioning film formation in the marine environment. *Microbiology* 150:1779–1784
- Baldock JA, Skjemstad JO (2000) Role of the soil matrix and minerals in protecting natural organic materials against biological attack. *Org Geochem* 31:697–710
- Bashan Y, Levanyon H (1988) Active attachment of *Azospirillum brasilense* Cd to quartz sand and to a light-textured soil by protein bridging. *J Gen Microbiol* 134:2269–2279
- Baveye P (2006) Comment on “Soil structure and management: a review” by Bronick CJ. *Lal R Geoderma* 13:231–232
- Bos R, van der Mei HC, Busscher HJ (1999) Physico-chemistry of initial microbial adhesive interactions – its mechanisms and methods for study. *FEMS Microbiol Rev* 23:179–229
- Brash JL, Horbett TA (1995) Proteins at interfaces. An overview. In: Horbett TA, Brash JL (eds) *Proteins at interfaces II fundamentals and applications*. American Chemical Society, Washington, DC, pp 1–25
- Burchill S, Hayes MHB, Greenland DJ (1981) Adsorption. In: Greenland DJ, Hayes MHB (eds) *The chemistry of soil surfaces*. Wiley, Chichester, pp 221–400
- Burke IC, Yonker CM, Parton WJ, Cole CV, Flach K, Schimel DS (1989) Texture, climate and cultivation effects on soil organic matter content in U.S. grassland soils. *Soil Sci Soc Am J* 53:800–805
- Chassin P (1979) Hypothèse sur la structure du complexe argilo-humique. Conséquences sur les processus d’humification. *Science du Sol* 4:323–332
- Chevallier T, Muchaonyerwa P, Chenu C (2003) Microbial utilisation of two proteins adsorbed to a vertisol clay fraction: toxin from *Bacillus thuringiensis* subsp. *tenebrionis* and bovine serum albumin. *Soil Biol Biochem* 35:1211–1218
- Chien Y-Y, Kim E-G, Bleam WF (1997) Paramagnetic relaxation of atrazine solubilized by humic micellar solutions. *Environ Sci Technol* 31:3204–3208
- Christensen BT (2001) Physical fractionation of soil and structural and functional complexity in organic matter turnover. *Eur J Soil Sci* 52:345–353
- Craig OE, Collins MJ (2002) The removal of protein from mineral surfaces: Implications for residue analysis of archaeological materials. *J Arch Sci* 29:1077–1082
- De Cristofaro A, Violante A (2001) Effect of hydroxy-aluminum species on the sorption and interlayering of albumin onto montmorillonite. *Appl Clay Sci* 19:59–67
- Ding X, Henrichs S (2002) Adsorption and desorption of proteins and polyamino acids by clay minerals and marine sediments. *Mar Chem* 77:225–237
- Dufrêne YF, Boonaert CJP, Rouxhet PG (1996) Adhesion of *Azospirillum brasilense*: role of proteins at the cell-support interface. *Coll Surf B Biointerfaces* 7:113–128
- Dufrêne YF, Boonaert CJP, Rouxhet PG (1999) Surface analysis by X-ray photoelectron spectroscopy in study of bioadhesion and biofilms. *Biofilms Methods Enzymol* 310:375–389
- Engelbreton RR, von Wandruszka R (1997) The effects of molecular size on humic acid associations. *Org Geochem* 26:759–767
- Engelbreton RR, von Wandruszka R (1998) Kinetic aspects of cation-enhanced aggregation in aqueous humic acids. *Environ Sci Technol* 32:488–493
- Ensminger LE, Gieseking JE (1942) Resistance of clay-adsorbed proteins to proteolytic hydrolysis. *Soil Sci* 53:205–209
- Essington ME (2003) *Soil and water chemistry*. CRC Press, Boca Raton
- Eusterhues K, Rumpel C, Kögel-Knabner I (2005) Organomineral associations in sandy acid forest soils: importance of specific surface area, iron oxides and micropores. *Eur J Soil Sci* 56:753–763
- Eusterhues K, Rumpel C, Kleber M, Kögel-Knabner I (2003) Stabilisation of soil organic matter by interactions with minerals as revealed by mineral dissolution and oxidative degradation. *Org Geochem* 34:1591–1600
- Ferreira J, Nascimiento OR, Martin-Neto L (2001) Hydrophobic interactions between spin-label 5-SASL and humic acid as revealed by ESR spectroscopy. *Environ Sci Tech* 35:761–765
- Galeska I, Hickey T, Moussy F, Kreutzer D, Papadimitrakopoulos F (2001) Characterization and biocompatibility studies of novel humic acids based films as membrane material for an implantable glucose sensor. *Biomacromolecules* 2:1249–1255
- Goodsell DS (2004) *Bionanotechnology: Lessons from nature*. Wiley-Liss, Inc., Hoboken, New Jersey
- Gougeon WD, Souillard M, Reinholdt M, Miehe-Brendle J, Chezeau JM, Le Dred R, Marchal R, Jeandet P (2003) Polypeptide adsorption on a synthetic montmorillonite: a combined solid-state NMR spectroscopy, X-ray diffraction, thermal analysis and N<sub>2</sub> adsorption study. *Eur J Inorganic Chem* 7:1366–1372
- Greenland DJ (1965) Interaction between clays and organic compounds in soils. Part I: Mechanisms of interaction between clays and defined organic compounds. *Soils Fertilizers* 28:415–420
- Guo M, Chorover J (2003) Transport and fractionation of dissolved organic matter in soil columns. *Soil Sci* 168:108–118
- Halthur TJ, Claesson PM, Elofsson UM (2004) Stability of polypeptide multilayers as studied by in situ ellipsometry: effects of drying and post-buildup changes in temperature and pH. *J Am Chem Soc* 126:17009–17015
- Haynes CA, Norde W (1994) Globular proteins at solid/liquid interfaces. *Coll Surf B: Biointerfaces* 2:517–566
- Hedges JI, Hare P (1987) Amino acid adsorption by clay minerals in distilled water. *Geochim Cosmochim Acta* 51:255–259

- Hedges JJ, Keil RG (1995) Sedimentary organic matter preservation: an assessment and speculative synthesis. *Mar Chem* 49:81–115
- Hedges JJ, Oades JM (1997) Comparative organic geochemistries of soils and marine sediments. *Org Geochem* 27:319–361
- Hlady V, Buijs J (1996) Protein adsorption on solid surfaces. *Curr Opin Biotechnol* 7:72–77
- Horne DJ, McIntosh JC (2000) Hydrophobic compounds in sands in New Zealand – extraction, characterisation and proposed mechanisms for repellency expression. *J Hydr* 231–232:35–46
- Hsieh YP (1996) Soil organic carbon pools of two tropical soils inferred by carbon signatures. *Soil Sci Soc Am J* 60:1117–1121
- Hunter KA, Liss PS (1982) Organic matter and the surface charge of suspended particles in estuarine waters. *Limnol Oceanogr* 27:322–335
- Jardine PM, Weber NL, McCarthy JF (1989) Mechanisms of dissolved organic carbon adsorption on soil. *Soil Sci Soc Am J* 53:1378–1385
- Jung E (1943) Zur Kenntnis der Ton-Humusbindung. *Z Pflanzenern Boden* 32:325–336
- Kahle M, Kleber M, Jahn R (2002) Carbon storage in loess derived surface soils from Central Germany: influence of mineral phase variables. *J Plant Nutr Soil Sci* 165:141–149
- Kaiser K, Guggenberger G (2003) Mineral surfaces and soil organic matter. *Eur J Soil Sci* 54:1–18
- Keil RG, Tsamakis E, Fuh CB, Giddings JC, Hedges JJ (1994) Mineralogical and textural controls on the organic composition of coastal marine sediments: hydrodynamic separation using SPLITT-fractionation. *Geochim Cosmochim Acta* 58:879–893
- Kleber M, Mertz C, Zikeli S, Knicker H, Jahn R (2004) Changes in surface reactivity and organic matter composition of clay subfractions with duration of fertilizer deprivation. *Eur J Soil Sci* 55:381–391
- Kleber M, Mikutta R, Torn MS, Jahn R (2005) Poorly crystalline mineral phases protect organic matter in acid subsoil horizons. *Eur J Soil Sci* 56:717–725
- Knicker H (2004) Stabilization of N-compounds in soil and organic-matter-rich sediments – what is the difference? *Mar Chem* 92:167–195
- Knicker H (2000) Biogenic nitrogen in soils as revealed by solid-state carbon-13 and nitrogen-15 nuclear magnetic resonance spectroscopy. *J Environ Qual* 29:715–723
- Knicker H (2002) The feasibility of using DCPMAS  $^{15}\text{N}$   $^{13}\text{C}$  NMR spectroscopy for a better characterization of immobilized  $^{15}\text{N}$  during incubation of  $^{13}\text{C}$  and  $^{15}\text{N}$ -enriched plant material. *Org Geochem* 33:237–246
- Knicker H, Lüdemann H-D (1995) N-15 and C-13 CPMAS and solution NMR studies of N-15 enriched plant material during 600 days of microbial degradation. *Org Geochem* 23:329–341
- Knicker H, Almendros G, González-Vila FJ, Lüdemann H-D, Martin F (1995)  $^{13}\text{C}$  and  $^{15}\text{N}$  NMR analysis of some fungal melanins in comparison with soil organic matter. *Org Geochem* 23:1023–1028
- Körschens M (1980) Beziehungen zwischen Feinanteil, Ct- und Nt-Gehalt des Bodens. *Arch Acker-Pfl Boden* 24:585–592
- Leinweber P, Schulten H-R (1995) Composition, stability and turnover of soil organic matter: investigations by off-line pyrolysis and direct pyrolysis-mass spectrometry. *J Anal Appl Pyrolysis* 32:91–110
- Leinweber P, Schulten H-R (1998) Advances in analytical pyrolysis of soil organic matter. *J Anal Appl Pyrolysis* 47:165–189
- Leinweber P, Schulten H-R (2000) Nonhydrolyzable forms of soil organic nitrogen: extractability and composition. *J Plant Nutr Soil Sci* 163:433–439
- Mahieu N, Olk DC, Randall EW (2000) Accumulation of heterocyclic nitrogen in humified organic matter: a  $^{15}\text{N}$ -NMR study of lowland rice soils. *Eur J Soil Sci* 51:379–389
- Mahieu N, Olk DC, Randall EW (2002) Multinuclear magnetic resonance analysis of two humic acid fractions from lowland rice soils. *J Environ Qual* 31:421–430
- Martin JP, Haider K (1986) Influence of mineral colloids on turnover rates of soil organic carbon. In: Huang PM, Schnitzer M (eds) Interactions of soil minerals with natural organics and microbes. SSSA, pp 283–304
- Martin-Neto L, Traghetta DG, Vaz CMP, Crestana S, Sposito G (2001) On the interaction of mechanisms of atrazine and hydroxyatrazine with humic substances. *J Environ Qual* 30:520–525
- Ma'shum M, Tate ME, Jones GP, Oades JM (1988) Extraction and characterization of water-repellent materials from Australian soils. *J Soil Sci* 39:99–100
- Mathers NJ, Mao XA, Xu ZH, Saffigna PG, Berners-Price SJ, Perera MCS (2000) Recent advances in the application of  $^{13}\text{C}$  and  $^{15}\text{N}$  NMR spectroscopy to soil organic matter studies. *Aust J Soil Res* 37:769–787
- Mayer LM (1999) Extent of coverage of mineral surfaces by organic matter in marine sediments. *Geochim Cosmochim Acta* 63:207–215
- Mayer LM, Xing B (2001) Organic matter – surface area relationships in acid soils. *Soil Sci Soc Am J* 65:250–258
- Mayer LM, Shick L, Sawyer T, Plante C, Jumars P, Self R (1995) Bioavailable amino acids in sediments: a biomimetic, kinetics-based approach. *Limnol Oceanogr* 40:511–520
- Mayer LM (1994) Relationships between mineral surfaces and organic carbon concentrations in soils and sediments. *Chem Geol* 114:347–363
- Mertz C, Kleber M, Jahn R (2005) Soil organic matter stabilization pathways in clay sub-fractions from a time series of fertilizer deprivation. *Org Geochem* 36:1311–1322
- Mikutta R, Kleber M, Torn M, Jahn R (2006) Stabilization of soil organic matter: association with minerals or chemical recalcitrance? *Biogeochemistry* 77:25–56
- Murphy EM, Zachara JM, Smith SC (1990) Influence of mineral-bound humic substances on the sorption of hydrophobic organic compounds. *Environ Sci Tech* 24:1507–1516
- Nanny MA, Kontas C (2002) Fluorescent spectroscopic analysis of non-covalent interactions between dissolved humic and fulvic acids and 6-propionyl-2-dimethylaminonaphthalene. In Preprints of extended abstracts presented at the 223rd American Chemical



- Society National Meeting. American Chemical Society, Orlando, FL, 42:528–531
- Nichols JD (1984) Relation of organic carbon to soil properties and climate in the southern great plains. *Soil Sci Soc Am J* 48:1382–1384
- Oades JM (1988) The retention of organic matter in soils. *Biogeochemistry* 5:35–70
- Omoike A, Chorover J (2006) Adsorption to goethite of extracellular polymeric substances from *Bacillus subtilis*. *Geochimica et Cosmochimica Acta* 70:827–838
- Parfitt RL, Theng BKG, Whitton JS, Shepherd TG (1997) Effects of clay minerals and land use on organic matter pools. *Geoderma* 75:1–12
- Ramsden JJ (1995) Puzzles and Paradoxes in Protein Adsorption. *Chem Soc Rev* 24:73–78
- Ransom B, Dongsom K, Kastner M, Wainwright S (1998) Organic matter preservation on continental slopes: importance of mineralogy and surface area. *Geochim Cosmochim Acta* 62:1329–1345
- Rice JA (2001) Humins. *Soil Sci* 166:848–857
- Scheffer F, Schachtschabel P (2002) *Lehrbuch der Bodenkunde*, 15. Auflage. Spektrum Akademischer Verlag, Heidelberg, Berlin
- Schmidt CF, Zimmermann RM, Gaub HE (1990) Multilayer adsorption of lysozyme on a hydrophobic substrate. *Biophys J* 57:577–588
- Soil Survey Staff (1999) *Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys*, 2nd edn. U. S. Government Printing Office, Washington, DC
- Sollins P, Swanston C, Kleber M, Filley T, Kramer M, Crow S, Caldwell BA, Lajtha K, Bowden R (2006) Organic C and N stabilization in a forest soil: evidence from sequential density fractionation. *Soil Biol Biochem* 38:3313–3324
- Spaeth K, Brecht A, Gauglitz G (1997) Studies on the biotin-avidin multilayer adsorption by spectroscopic ellipsometry. *J Coll Interface Sci* 196:128–135
- Stevenson FJ (1985) Geochemistry of soil humic substances. In: McKnight DM (ed) *Humic substances in soil, sediment and water: geochemistry, isolation and characterization*. John Wiley and Sons, New York
- Strickland TC, Sollins P, Rudd N, Schimel DS (1992) Rapid stabilization and mobilization of <sup>15</sup>N in forest and range soils. *Soil Biol Biochem* 24:849–855
- Sutton R, Sposito G (2005) Molecular structure in soil humic substances: the new view. *Environ Sci Technol* 39:9009–9015
- Swanston CW, Torn MS, Hanson PJ, Southon JR, Garten CT, Hanlon EM, Ganio L (2005) Initial characterization of processes of soil carbon stabilization using forest stand-level radiocarbon enrichment. *Geoderma* 128:52–62
- Theng BKG (1979) *Formation and properties of clay-polymer complexes*. Elsevier Science Publishing Co., Amsterdam
- Theng BKG, Tate KR, Becker-Heidmann P (1992) Towards establishing the age, location, and identity of the inert soil organic matter of a spodosol. *Z Pflanzenern Bodenk* 155:181–184
- Tiberg F, Brinck J, Grant L (1999) Adsorption and surface-induced self-assembly of surfactants at the solid-aqueous interface. *Curr Opin Colloid Interface Sci* 4:411–419
- Tremblay L, Brenner R (2006) Microbial contributions to N-immobilization and organic matter preservation in decaying plant detritus. *Geochim Cosmochim Acta* 70:133–146
- Trumbore SA, Vogel JS, Southon JR (1989) AMS <sup>14</sup>C measurements of fractionated soil organic matter: an approach to deciphering the soil carbon cycle. *Radiocarbon* 31:644–654
- Trumbore SE, Zheng SH (1996) Comparison of fractionation methods for soil organic matter C-14 analysis. *Radiocarbon* 38:219–229
- Turchenek LW, Oades JM (1979) Fractionation of organo-mineral complexes by sedimentation and density techniques. *Geoderma* 21:311–343
- Voice TC, Weber JWJ (1983) Sorption of hydrophobic compounds by sediments, soils and suspended solids – I. Theory and background. *Water Res* 17:1433–1441
- von Wandruszka R (1998) The micellar model of humic acid: evidence from pyrene fluorescence measurements. *Soil Sci* 163:921–930
- von Wandruszka R, Engebretson R (2001) Kinetics of humic acid associations. In: Clapp CE, Hayes MHB, Senesi N, Bloom PR, Jardine PM (eds) *Humic substances and chemical contaminants*. Soil Science Society of America, Madison, WI, pp 119–126
- von Wandruszka R, Engebretson RR, Yates LM (1999) Humic acid pseudomicelles in dilute aqueous solution: Fluorescence and surface tension measurements. In: Ghabbour EA, Davies G (eds) *Understanding humic substances: Advanced methods, properties and applications*. Royal Society of Chemistry, Cambridge, UK, pp 79–85
- Wang K, Xing B (2005) Structural and sorption characteristics of adsorbed humic acid on clay minerals. *J Environ Qual* 34:342–349
- Wang X-C, Lee C (1993) Adsorption and desorption of aliphatic amines, amino acids and acetate by clay minerals and marine sediments. *Mar Chem* 44:1–23
- Weber JWJ, Voice TC, Pirbazari M, Hunt GE, Ulanoff DM (1983) Sorption of hydrophobic compounds by sediments, soils and suspended solids – II. Sorbent evaluation studies. *Water Res* 17:1443–1452
- Wershaw RL (1986) A new model for humic materials and their interactions with hydrophobic organic chemicals in soil-water or sediment-water systems. *J Cont Hydr* 1:29–45
- Wershaw RL (1993) Model for humus in soils and sediments. *Environ Sci Technol* 27:814–816
- Wershaw RL (2004) Evaluation of conceptual models of natural organic matter (humus) from a consideration of the chemical and biochemical processes of humification. U.S. Geological Survey, Reston, VA, p 44
- Wershaw RL, Pinckney DJ (1980) Isolation and characterization of clay-humic complexes In: Baker RA (ed) *Contaminants and sediments*. Ann Arbor Science Publishers, Ann Arbor, Michigan, pp 207–219

- Wershaw RL, Leenheer JA, Sperline RP, Song Y, Noll LA, Melvin RL, Rigatti GP (1995) Mechanism of formation of humus coatings on mineral surfaces 1. Evidence for multidentate binding of organic acids from compost leachate on alumina. *Coll Surf A: Physicochem Eng Aspects* 96:93–104
- Wershaw RL, Llaguno EC, Leenheer JA (1996a) Mechanism of formation of humus coatings on mineral surfaces 3. Composition of adsorbed organic acids from compost leachate on alumina by solid-state  $^{13}\text{C}$  NMR. *Coll Surf A: Physicochem Eng Aspects* 108:213–223
- Wershaw RL, Llaguno EC, Leenheer JA, Sperline RP, Song Y (1996b) Mechanism of formation of humus coatings on mineral surfaces 2. Attenuated total reflectance spectra of hydrophobic and hydrophilic fractions of organic acids from compost leachate on alumina. *Coll Surf A: Physicochem Eng Aspects* 108:199–211
- White RE (1997) *Principles and practice of soil science: the soil as a natural resource*. Blackwell Publishing, Oxford
- Wright SF, Upadhyaya A (1996) Extraction of an abundant and unusual protein from soil and comparison with hyphal protein of arbuscular mycorrhizal fungi. *Soil Sci* 161:575–586
- Young JL, Spycher G (1979) Water dispersible soil organic-mineral particles. *Soil Sci Soc Am J* 43:324–328
- Zang X, van Heemst JDH, Dria KJ Hatcher PG (2000) Encapsulation of protein in humic acid from a histosol as an explanation for the occurrence of organic nitrogen in soil and sediment. *Org Geochem* 31:679–695
- Zhou JL, Rowland S, Mantoura RFC, Braven J (1994) The formation of humic coatings on mineral particles under simulated estuarine conditions – a mechanistic study. *Water Res* 28:571–579