



Reviews and syntheses: Systematic Earth observations for use in terrestrial carbon cycle data assimilation systems

Marko Scholze¹, Michael Buchwitz², Wouter Dorigo³, Luis Guanter⁴, and Shaun Quegan⁵

¹Department of Physical Geography and Ecosystem Science, Lund University, Lund, Sweden

²Institute of Environmental Physics (IUP), University of Bremen, Bremen, Germany

³Department of Geodesy and Geoinformation, Vienna University of Technology (TU Wien), Vienna, Austria

⁴Remote Sensing Section, German Research Center for Geosciences (GFZ), 14473 Potsdam, Germany

⁵Centre for Terrestrial Carbon Dynamics, The University of Sheffield, Sheffield S3 7RH, UK

Correspondence to: Marko Scholze (marko.scholze@nateko.lu.se)

Received: 22 December 2016 – Discussion started: 12 January 2017

Revised: 19 June 2017 – Accepted: 20 June 2017 – Published: 19 July 2017

Abstract. The global carbon cycle is an important component of the Earth system and it interacts with the hydrology, energy and nutrient cycles as well as ecosystem dynamics. A better understanding of the global carbon cycle is required for improved projections of climate change including corresponding changes in water and food resources and for the verification of measures to reduce anthropogenic greenhouse gas emissions. An improved understanding of the carbon cycle can be achieved by data assimilation systems, which integrate observations relevant to the carbon cycle into coupled carbon, water, energy and nutrient models. Hence, the ingredients for such systems are a carbon cycle model, an algorithm for the assimilation and systematic and well error-characterised observations relevant to the carbon cycle. Relevant observations for assimilation include various in situ measurements in the atmosphere (e.g. concentrations of CO₂ and other gases) and on land (e.g. fluxes of carbon water and energy, carbon stocks) as well as remote sensing observations (e.g. atmospheric composition, vegetation and surface properties).

We briefly review the different existing data assimilation techniques and contrast them to model benchmarking and evaluation efforts (which also rely on observations). A common requirement for all assimilation techniques is a full description of the observational data properties. Uncertainty estimates of the observations are as important as the observa-

tions themselves because they similarly determine the outcome of such assimilation systems. Hence, this article reviews the requirements of data assimilation systems on observations and provides a non-exhaustive overview of current observations and their uncertainties for use in terrestrial carbon cycle data assimilation. We report on progress since the review of model-data synthesis in terrestrial carbon observations by Raupach et al. (2005), emphasising the rapid advance in relevant space-based observations.

1 Introduction

The anthropogenic perturbation of the global carbon cycle has led to a global mean increase of 43 % in atmospheric CO₂ (from 280 to 398 ppm) in 2014 compared to pre-industrial (before 1750) levels (WMO, 2015) and is the main driver of climate change. The main causes for the increase in CO₂ are burning of fossil fuels and land use change, which amount to emissions of 9.8 ± 0.5 GtC in 2014. However, only about 44 % of these emissions stay in the atmosphere; the remainder is currently taken up by the land biosphere ($\approx 30\%$) and the surface ocean ($\approx 26\%$; Le Quéré et al., 2015). Positive climate-carbon cycle feedbacks, predominantly acting on land processes, may reduce this sink capacity and thus

accelerate global warming (Matthews et al., 2007). Also, the sink strength of the terrestrial biosphere is more variable than that of the ocean (Ciais et al., 2013) and its quantification by process-based terrestrial carbon cycle models exhibit large uncertainties (Le Quéré et al., 2015).

A common way to reduce uncertainties from process-based modelling is by confronting these models with observational data. Raupach et al. (2005) pointed out that the systematic combination of observational data with process modelling, which is commonly referred to as “model-data fusion”, is an effective strategy for observing the Earth system. The term model-data fusion is sometimes understood in a more general way, which is that observational data is blended with (pre-computed) model output, whereas the term “data assimilation” refers to a robust mathematical framework for improving model predictions with observational data. Data assimilation is motivated by several benefits to make the best use of observations and models (Mathieu and O’Neill, 2008). These benefits include, among others, (1) forecasting and initialisation (forward predictions in time based on past observations), (2) model and data quality control (regular and systematic confrontation of model output with observations within their uncertainty statistics), (3) a combination of various data streams (combined constraints of independent observations can be stronger than the sum of the individual constraints), (4) filling in regions with sparse observations (consistent propagation of information from data-rich regions to data-poor regions), (5) estimating unobservable quantities (through process-based relations in the model observations constrain modelled quantities which are not directly measured) and (6) observing system design (what is the delta of a new type of observation).

Systematic observations are a key ingredient for data assimilation studies. Here, we focus on the carbon cycle and the land–atmosphere system. The land–atmosphere components of the carbon cycle are an important part of an integrated Earth observation system because of the close interactions on land between the carbon cycle and the water and energy cycles and hence its importance for climate projections and climate change mitigation strategies through the monitoring and management of terrestrial greenhouse gas sources and sinks.

Raupach et al. (2005) provide an analysis of the various elements of a Terrestrial Carbon Observation System (TCOS). The need for, design and steps to be taken towards a TCOS were already outlined by others before (Cihlar et al., 2002; Global Carbon Project, 2003) but Raupach et al. (2005) systematically reviewed two major components of a TCOS: the data assimilation methods and the observational data and data uncertainty characteristics for some selected, main kinds of relevant data. The requirements for a policy-relevant carbon observing system have been outlined by Ciais et al. (2014). They review the current systematic carbon cycle observations and illustrate the implementation of such a policy-relevant carbon observing system.

In this paper we provide an update of the observational data and data uncertainty characteristics as assessed by Raupach et al. (2005) with a focus on existing but also new and upcoming, relevant space-based observations, in the following referred to as Earth observation (EO) data. In contrast to (Ciais et al., 2014), who focus on carbon cycle observations, we focus here on relevant observational data to be (potentially) assimilated in a terrestrial carbon cycle data assimilation system (CCDAS).

In a CCDAS non-carbon observations can be exploited to constrain the simulated carbon cycle indirectly through the relations implemented in the process model. Such observational constraints act by ruling out combinations of the unknowns in a CCDAS (typically a combination of process parameters, initial- or boundary conditions), which are inconsistent with the observations and thereby reduce uncertainties in the simulated output. In that sense we are somewhat broader in terms of observed variables because the “non-carbon” observations (such as soil moisture or land surface temperature) are also able to constrain the carbon cycle indirectly through process information embedded in the underlying models. At the same time, the focus of our review is narrower than that of Ciais et al. (2014), who also addressed ocean and anthropogenic components.

The paper is organised as follows: in the next section we contrast data assimilation with recently established benchmarking activities and give a brief overview of commonly used data assimilation approaches and their applications in terrestrial carbon cycling. We continue with a short overview on data characteristics including an update on progress for some of the observations discussed in Raupach et al. (2005). Since there has been much developments in the provision of remotely sensed observations, we focus here on the characteristics of the most relevant EO data streams.

2 Data assimilation

2.1 Data assimilation versus benchmarking

In the recent past the international land surface and terrestrial ecosystem modelling communities have recognised the importance of model benchmarking and evaluation (e.g. Luo et al., 2012; Foley et al., 2013). One of the reasons for this development is the huge range of model results from different models in key diagnostics of the land–atmosphere interface such as gross primary productivity (GPP) and latent heat flux (Prentice et al., 2015).

In general “benchmarking” is understood as the quantification of performance against a reference using some pre-defined metrics. The reference can either be output from some previous model simulations, other (ensembles of) models or reference data sets based on observations if the model simulates the analogue quantity. Luo et al. (2012) suggest a theoretical framework for benchmarking land models based

on standardised references and metrics to measure model performance skills. A large variety of such metrics and their characteristics is introduced by Foley et al. (2013). Some examples of benchmarking terrestrial carbon cycle models (either stand-alone or coupled to climate models) are given, for example, by Randerson et al. (2009), Cadule et al. (2010) and Kelley et al. (2013).

The commonality between benchmarking/evaluation and data assimilation lies in the quantitative assessment of model output. In benchmarking the quantitative assessment is performed by calculating some metrics against either observations or other references, while in data assimilation this is achieved by defining a cost function, which quantifies the mismatch of some simulated model quantity against observations weighted by the inverse of their uncertainties (including a model uncertainty). However, data assimilation goes beyond benchmarking as it minimises the quantified mismatch to improve model performance directly by adjusting initial and boundary conditions, state variables and/or model process parameters.

As pointed out by Prentice et al. (2015) there is a need for both model benchmarking and data assimilation: benchmarking may be used as a routine application to improve confidence and evaluate the performance (over time) in terrestrial carbon cycle modelling. However, if a benchmark test for a given model fails, this could simply imply that the model parameter values have not been specified correctly and optimised against observations. In contrast, data assimilation, in particular when used for parameter optimisation, potentially identifies structural model and/or data deficiencies if the model-data mismatch (or the benchmark test) is still inadequate after optimisation (see also Fig. 1). On the other hand, a better fit between the posterior maximum likelihood simulation (i.e. using the optimised parameters) and the observations is not necessarily an indication for correct parameters and/or model structure as has been pointed out by MacBean et al. (2016).

2.2 Data assimilation methods

The general problem of data assimilation can be formulated (following the notation of Rayner et al., 2016) as follows: given a model M , a set of observations \mathbf{y} of some observables $\mathbf{o} = H(\mathbf{z})$, with \mathbf{z} being the state variables of the model, H the observation operator and prior information on some target variables \mathbf{x} , produce an updated description of \mathbf{x} . \mathbf{x} may include elements of \mathbf{z} and \mathbf{p} (parameters; quantities not changed by the model, i.e. process parameters, boundary and initial conditions). The observation operator maps the model state onto observables. In the case of a CCDAS-assimilating atmospheric CO₂ the observation operator is the atmospheric transport model that maps the net CO₂ surface exchange fluxes as calculated by the terrestrial carbon cycle onto simulated atmospheric CO₂ concentrations.




Figure 1. Schematic of a data assimilation system with \mathbf{x} being the control vector containing the quantities to be updated by the assimilation. The loop between the “evaluation of J ” box to the “model and observation operator” box indicates the assimilation process (assimilation loop). Often, the analysis of residuals in model-data comparison leads to either model improvements or adjustment of the measurement strategies (“model improvement” and “adjusting measurement strategy” arrows).

A data assimilation system consists of three main ingredients: a set of observations, a dynamical model including the observation operator and an assimilation method. When assimilating multiple data streams, each data stream usually requires its own observation operator (see e.g. Kaminski and Mathieu, 2017). In the Bayesian formulation of the assimilation problem uncertainties (i.e. the description of quantities by probability density functions, PDFs) are central to the concept of data assimilation. Both observations as well as models have errors arising for various reasons. We will detail the observational errors in the next section. Dynamical models as well as observation operators have errors arising from the parameterisations and the discretisation of analytical dynamics into a numerical model; for a more complete description of uncertainty in Earth system models or components of such we refer to Scholze et al. (2012).

We distinguish two basic approaches in data assimilation: sequential assimilation, which assimilates observations subsequently at discrete model time steps, and variational assimilation, which assimilates all observations at once at their respective measurement times over a given period, the so-called assimilation window. They differ in their numerical efficiency and adequacy for their specific use. A general data-assimilation scheme is shown in Fig. 1. In the sequential approach the assimilation loop is evaluated sequentially over time following the dynamics of the model. In the case of

variational assimilation the assimilation loop is evaluated iteratively (assuming a non-linear model). Both cases evaluate a cost function J , formulated in the Bayesian framework as follows:

$$J = \frac{1}{2} \left[(\mathbf{x} - \mathbf{x}^b)^T \mathbf{B}^{-1} (\mathbf{x} - \mathbf{x}^b) + (\mathbf{H}(\mathbf{x}) - \mathbf{y})^T \mathbf{R}^{-1} (\mathbf{H}(\mathbf{x}) - \mathbf{y}) \right], \quad (1)$$

where \mathbf{x}^b is the prior information, \mathbf{B} the prior uncertainty covariance and \mathbf{R} the observational uncertainty covariance. When multiple data streams with different observation operators are assimilated, there will be several summands of the form of the second term on the right hand side of Eq. (1), one for each data stream.

Rayner et al. (2016) introduce the theory fundamental to data assimilation and illustrate how the different implementations of data assimilation relate to this theory in a more narrative style. A more complete and mathematically precise introduction to the concepts of data assimilation is given in the textbooks by e.g. Daley (1991) and Tarantola (2005).

2.3 Examples of terrestrial carbon cycle data assimilation

A variety of the methods as described by Rayner et al. (2016) have been applied by the carbon cycle community. One example that is making use of formal assimilation methodologies for inferring surface-atmosphere CO₂ exchange fluxes is based on atmospheric transport inversions. As mentioned before, in atmospheric inversions the observation model is an atmospheric tracer transport model. In atmospheric inversions both sequential and variational methods have been used together with observations of atmospheric trace gas concentrations such as from the flask sampling network, continuous in situ and aircraft measurements and more recently also remotely sensed total column measurements. The techniques for atmospheric transport inversions have been detailed in Enting (2002) and a recent comparison of results from different transport inversion is given by Peylin et al. (2013).

A more recent development is the assimilation of observations into terrestrial biosphere models. Here, various methods and observations have been used to optimise model process parameters at different scales. A comparison of a whole suite of these assimilation methods applied to a test case using a simplified model at local scale is given by Trudinger et al. (2007) and Fox et al. (2009).

Kaminski et al. (2002) were among the first who applied a formal algorithm together with observations of atmospheric CO₂ concentrations to constrain the simple diagnostic biosphere model at global scale. This work was continued by the development of the first carbon cycle data assimilation system (CCDAS) with a process-based model (BETHY) at its core (Rayner et al., 2005). The advantage of using a process-based model at the core of a CCDAS is that once the process parameters have been optimised the constrained model can also be used for predictions as demonstrated by Scholze et al.

(2007). Also, such systems are capable of ingesting multiple independent data streams besides atmospheric CO₂ concentrations. Kaminski et al. (2013) provide an overview on the developments of the CCDAS-BETHY since its first application while Scholze et al. (2016) demonstrate the latest application of CCDAS-BETHY, assimilating atmospheric CO₂ and remotely sensed surface soil moisture simultaneously. Since then several global terrestrial ecosystem models have been included in CCDAS employing a variational approach (e.g. Schüermann et al., 2016; Peylin et al., 2016).

Concurrently, there have been several studies at the local/regional scale assimilating various types of observations. For instance, Barrett (2002) used a genetic algorithm to infer soil carbon turnover times in a terrestrial carbon cycle model over Australia from in situ observations of plant production, biomass, litter and soil carbon. Local eddy covariance flux tower measurements of net exchange of CO₂ and latent and sensible heat fluxes have been assimilated to optimise parameters related to photosynthesis, respiration and energy fluxes of terrestrial ecosystem models using Monte-Carlo-type methods (e.g. Braswell et al., 2005; Knorr and Kattge, 2005; Moore et al., 2008; Ricciuto et al., 2008; Post et al., 2017), sequential methods (Williams et al., 2005) as well as variational approaches (e.g. Wang et al., 2001; Kuppel et al., 2012; Raoult et al., 2016)

Recent advances focus on multiple independent data stream assimilation to provide a more rigorous constraint on the multiple components of terrestrial ecosystem models and avoid equifinality, i.e. different parameter solutions providing the same cost function value at the minimum. Examples for such studies on local/regional scale are the assimilation of eddy covariance CO₂ fluxes together with observations of vegetation structural information or carbon stocks (e.g. Richardson et al., 2010; Keenan et al., 2012; Thum et al., 2017). The assimilation of multiple data streams can be performed either in a stepwise (e.g. Peylin et al., 2016) or simultaneous approach (e.g. Kaminski et al., 2012); in the case of non-linear models or non-linear observation operators only the simultaneous assimilation makes optimal use of the observations (MacBean et al., 2016). In Sect. 3.2 we provide more terrestrial carbon cycle data assimilation examples using some of the remotely sensed products discussed in the following.

3 Data characteristics and provision

Observations are our measurable representation of the “truth”. They come with different characteristics in terms of spatial and temporal resolution, coverage of the observed system and errors. In analogy, models are also some representation of the truth, but via knowledge embodied in some form of functional relationships (with their own errors as mentioned before). The paper by Raupach et al. (2005) has been instrumental in highlighting the challenges in combin-




Figure 2. Space–time diagram for a range of observations relevant for a Terrestrial Carbon Observation System.

ing models and observational data for building a TCOS focusing on the observational requirements. Ciais et al. (2014) argue for a globally integrated carbon observation system to improve our understanding of the carbon cycle for predicting future changes and to be able to independently verify the impact of emission reduction measures. Such a system relies on atmospheric carbon observations as a backbone but also concerns observations of the terrestrial and ocean carbon cycle. They focus on a strategy towards a global carbon cycle monitoring system for achieving the above mentioned objectives.

Figure 2 depicts exemplarily the main observations of a TCOS and their space–time characteristics. In the following we briefly summarise the aspects of uncertainty in the observations and highlight progress on the specification of uncertainty for some of the observations in Fig. 2 as well as on their monitoring since Raupach et al. (2005).

3.1 Observational uncertainty

As mentioned before an important ingredient to any data assimilation system is not only the observations themselves but also the uncertainties associated to them. We distinguish three main types of observation errors:

- Random errors are always present in measurements and are caused by unpredictable changes in the measurement system (e.g. electronic noise in electrical instrument). They show up as different readings of the same repeated measurement and thus can be reduced by taking the average of multiple measurements. Random errors are usually assumed to be normal (Gaussian) distributed, however, in some cases the random error distribution is log-normal (e.g. precipitation) or skewed by outliers due to unpredictable corruptions of the measurement system. Random errors are therefore related to the precision of a measurement system.

– Systematic (bias) errors in observations are usually due to some recurring problems in the overall measurement system. They are caused by instrument miscalibrations or interferences with the measurement system. They can vary in space and time but they affect the measurement system in a predictable way. Biases can be both additive (absolute mean bias) and multiplicative (biases in the dynamic range affecting the amplitude of a signal). If the source for systematic errors is known they can usually be fixed and should be removed. Systematic errors are therefore related to the accuracy of a measurement system.

– Representativeness error occurs when information is represented at a scale different from the source of the information. For instance a quantity simulated by a model is representative for a given spatial and temporal resolution of the model grid. In fact, the scale at which we trust the model may be larger than a grid cell. An individual measurement, however, represents information influenced by the local environment that is not resolved by the model grid (e.g. representation of atmospheric flask data in an atmospheric transport model grid cell).

For both random and systematic errors not only the magnitude of the error for a single observation is important, i.e. the diagonal elements in the observational uncertainty covariance matrix \mathbf{B} , but also the correlations between errors for different observations. Hence there is a need to specify the off-diagonal elements in the error covariance matrix \mathbf{B} . These off-diagonal elements are usually hard to specify, but it is important to quantify them in a data assimilation system. They have considerable impact on the solution because of their influence on the weight of the respective observations in the cost function.

In addition to the observational errors, models have errors, which, in a data assimilation system, are usually included in the observation errors. These errors in dynamical models are mainly caused by process parameterisations (instead of resolving the process) and by the discretisation of analytical dynamics into a numerical model. A more detailed description of the different model error sources is given in Scholze et al. (2012).

As mentioned before, Raupach et al. (2005) have already reflected on the main properties of the data and their error covariances for observations of remotely sensed land surface properties (mainly the normalised differential vegetation index, NDVI), atmospheric CO₂ concentrations, land–atmosphere net CO₂ exchange fluxes and terrestrial carbon stores. The in situ measurements of CO₂ concentrations are either based on flask samples or on continuous monitoring stations. The flask sampling network was established in 1961 by Keeling (1961) and has been extended since then to more than 200 sites globally. The continuous in situ network provide measurements at higher precision and temporal resolution than the flask networks. For both the flask and the contin-

uous stations, improvements in precision and accuracy have been achieved through propagation of frequent comparisons and international standards (Francey et al., 2001).

The global FluxNet network consists of more than 200 sites globally measuring land–atmosphere fluxes of CO₂, latent and sensible heat and others using the eddy covariance technique at a half-hourly temporal resolution (Balocchi et al., 2001). Many other (mostly meteorological) variables are measured at these sites as well. In the past years, there has been substantial progress in the homogenisation and availability of these direct CO₂ flux measurements. The publicly available FLUXNET2015 data set includes more than 1500 site-years of data covering all major biome types from about 165 sites worldwide, spanning a period from 1991 (for some sites) up to 2014 (Pastorello et al., 2017). There has also been substantial progress in the specification of uncertainties in eddy covariance measurements of the land–atmosphere net CO₂ exchange flux (net ecosystem productivity, NEP) and its component fluxes (GPP and ecosystem respiration, R_{eco}). For instance, Lasslop et al. (2008) analysed the error distribution and found that the eddy flux data can almost entirely be represented by a superposition of Gaussian distributions with inhomogeneous variance. Richardson et al. (2008) showed that the measurement errors in NEP are heteroscedastic; i.e. the error variance varies with the magnitude of the flux. In a more recent study Raj et al. (2016) investigated the uncertainty of GPP derived from partitioning the eddy covariance NEP measurements. They used a light-use efficiency model embedded in a Bayesian framework to estimate the uncertainty in the separated GPP from the posterior distribution at half-hourly time steps.

3.2 Examples of systematic observations from satellite EO data

There has been a vast extension of EO capabilities during the past 10 years or so both in terms of product quality (including, for instance, improved accuracy) and quantity (new products).

In any data assimilation system using satellite EO data one needs to decide in the design phase of the system whether to assimilate observations at the sensor level (i.e. the spectral radiances for optical sensors or brightness temperatures for microwave sensor, referred to as level 1 data) or to assimilate the biogeophysical variable derived from the radiances through a retrieval algorithm (level 2 data product). When assimilating level 1 data the retrieval algorithm is part of the observation operator linking the model state to the observations in the data assimilation system. A more detailed description of the two alternatives in assimilating EO satellite observations into models of the Earth system is given by Kaminski and Mathieu (2017). In carbon cycle data assimilation systems level 2 data products (or even level 3 data, which are provided on a regular space–time grid) are most commonly used. However, there is a risk that when using products at

level 2 or higher, the parameters/processes implemented in the retrieval algorithm may not be consistent with the corresponding equivalent parameters/processes in the underlying model, thus causing additional errors in the assimilation.

In the next subsections we present some selected remotely sensed Earth observation products, which are relevant for terrestrial carbon cycle data assimilation, in more detail:

- atmospheric CO₂,
- vegetation activity (FAPAR and SIF),
- soil moisture,
- terrestrial biomass.

These EO products either have already been used, are in the process of being used or would potentially be a useful data constraint in a CCDAS. For vegetation activity we distinguish two major types of products: more “traditional” reflectance- or radiative-based products, such as the fraction of absorbed photosynthetically active radiation (FAPAR), and recently developed products based on biogeochemical processes, such as sun-induced fluorescence (SIF). Leaf area index (LAI, e.g. Liu et al., 2014), which is in effect closely related to FAPAR, is another geophysical parameter that represents vegetation activity. There is also a range of remotely sensed vegetation indices, of which NDVI is an example. Both LAI and NDVI have been used in data assimilation studies: an example for NDVI is given by MacBean et al. (2015) and for LAI by Luke (2011) and Barbu et al. (2014). In Sect. 3.2.2 we detail the difference between NDVI and FAPAR and explain that FAPAR is based on physical principles. FAPAR has already been demonstrated to provide a strong constraint on terrestrial carbon and water fluxes through its impact on the phenology components of the carbon cycle model either by assimilating only FAPAR data (e.g. Knorr et al., 2010) or in combination with other data streams (e.g. Kaminski et al., 2012; Kato et al., 2013; Forkel et al., 2014). SIF is a promising observation for constraining the gross uptake of CO₂ by plant photosynthesis. First assimilation results using SIF observations in a CCDAS show that the uncertainty in global annual GPP is largely reduced by constraining parameters that describe leaf phenology (Norton et al., 2016). Remotely sensed atmospheric CO₂ concentration (XCO₂; see Sect. 3.2.1) has also been assimilated into a diagnostic terrestrial carbon cycle model to derive net CO₂ fluxes consistent with independent in situ measurements of atmospheric CO₂ and to reduce posterior uncertainties in the inferred net and gross CO₂ fluxes (Kaminski et al., 2016). Barbu et al. (2014) and Albergel et al. (2017) assimilated both soil moisture and LAI data into a land surface model, but their focus was on improving the hydrological and land surface physical quantities and not the carbon cycle. van der Molen et al. (2016) assessed the impact of assimilating various remotely sensed soil moisture products into the SiB-CASA ecosystem model on simulated carbon fluxes in bo-

real Eurasia. Although the impact of assimilating ASCAT surface soil moisture was significant, its skill in this hydrologically complex environment strongly depends on surface water and vegetation dynamics. In contrast, Scholze et al. (2016) showed that when assimilating SMOS soil moisture simultaneously with in situ atmospheric CO₂ concentrations, the reduction of uncertainty in gross and net CO₂ fluxes relative to the prior is considerably higher than when only assimilating CO₂, which quantifies the added value of SMOS observations as a constraint on the terrestrial carbon cycle. So far, remotely sensed biomass data have not been used in carbon cycle data assimilation studies, but several studies (e.g. Richardson et al., 2010; Keenan et al., 2012; Thum et al., 2017) have demonstrated the added value of in situ above-ground biomass observations in constraining the terrestrial carbon cycle.

This list of EO products described in this paper is admittedly subjective and there are of course a whole range of additional remotely sensed products available, which are relevant for carbon cycle studies as well, e.g. burned area (e.g. Giglio et al., 2013), land cover (e.g. Bontemps et al., 2012), land surface temperature (e.g. Li et al., 2013) or vegetation optical depth (VOD; e.g. Konings et al. (2016)). However, these products are rather used as input or boundary conditions for terrestrial carbon cycle models (burned area and land cover) or, in the case of land surface temperature and VOD, they have so far not been used in carbon cycle data assimilation studies.

3.2.1 Atmospheric CO₂ and CH₄

Satellite retrievals of atmospheric carbon dioxide (CO₂) and methane (CH₄) are available from several satellite instruments such as mid-tropospheric CO₂ and CH₄ columns from the Infrared Atmospheric Sounding Interferometer (IASI; e.g. (Crevoisier et al., 2009a, b)) on EUMETSAT's Metop satellite series, vertical profiles with highest sensitivity in the middle/upper troposphere from AIRS on Aqua (e.g. Xiong et al., 2013), stratospheric profiles from MIPAS on ENVISAT limb observations (e.g. Laeng et al., 2015) and from the solar occultation observations of SCIAMACHY on ENVISAT (Noël et al., 2011, 2016) and ACE-FTS (e.g. Boone et al., 2005; Foucher et al., 2009). These observations have, however, only little or no sensitivity to CO₂ and CH₄ concentration changes close to the Earth's surface and therefore contain only limited information on regional or local CO₂ and CH₄ sources and sinks. Satellites with high near-surface sensitivity are nadir (down-looking) satellites which measure radiance spectra of reflected solar radiation in the relevant spectral bands in the near-infrared/shortwave-infrared (NIR/SWIR) spectral region, which are located around 1.6 μm (CO₂ and CH₄) and around 2.0 μm (CO₂). Satellites instruments which perform (or have performed) these observations are SCIAMACHY on board ENVISAT (2002–2012; Burrows et al., 1995; Bovensmann et al., 1999),

TANSO-FTS on board GOSAT (launched in 2009; Kuze et al., 2009, 2014) and NASA's Orbiting Carbon Observatory 2 (OCO-2) mission (launched in 2014; Crisp et al., 2004; Boesch et al., 2011).

The main CO₂ and CH₄ data products of these sensors are near-surface-sensitive column-averaged dry-air mole fractions of CO₂ and CH₄, denoted XCO₂ and XCH₄. The quantities XCO₂ and XCH₄ are both retrieved from SCIAMACHY/ENVISAT (ground pixel size: 30 × 50 km², along track times across track; swath width 960 km with contiguous ground pixels) and TANSO-FTS/GOSAT (10 km pixel size; several, e.g. 3 or 5 non-contiguous pixels across track). OCO-2 delivers XCO₂ (8 ground pixels across track, each ≈ 1.3 km) and other satellites have been or will be launched, such as Europe's Sentinel-5-Precursor satellite (S5P; Veefkind et al., 2012), which will deliver (among several other parameters) XCH₄ (7 km pixel size at nadir, 2600 km swath width with contiguous ground pixels; planned launch: autumn 2017) (Butz et al., 2012) and China's TanSat (launched end of 2016), which will deliver XCO₂ with similar characteristics to NASA's OCO-2. It can be expected that future satellites will provide improved measurements, in particular with respect to more localised emission sources (e.g. Buchwitz et al., 2013; Ciais et al., 2015). In the following we focus the discussion on sensors that have already delivered multiyear XCO₂ and XCH₄ data sets, i.e. on SCIAMACHY and TANSO.

These satellite-derived XCO₂ and XCH₄ data products are sensitive to surface fluxes because CO₂ and CH₄ emission and uptake by surface sources and sinks result in the largest changes of the atmospheric CO₂ and CH₄ mixing ratio close to the Earth's surface and therefore modify the observed vertical columns. This results in local or regional atmospheric enhancements (e.g. Buchwitz et al., 2017, discussing localised methane sources) or large-scale atmospheric gradients (e.g. Reuter et al., 2014, discussing CO₂ uptake by the terrestrial biosphere).

The XCO₂ and XCH₄ data products retrieved from SCIAMACHY and TANSO are generated from the radiance observations using different approaches. Most approaches are based on optimal estimation (OE; e.g. Rogers, 2000; Reuter et al., 2010), also called Bayesian inference. OE permits constraining the retrieval using a priori information on, e.g. atmospheric vertical profiles of trace gases and aerosols. In general, the radiances are simulated using a radiative transfer model (RTM) and RTM and other parameters (state vector elements) are adjusted until an optimal match is achieved between observed and simulated radiances. One algorithm (WFM-DOAS, WFMD Buchwitz et al., 2000; Schneising et al., 2008, 2009) is based on least-squares and does not use a priori information to constrain the fit parameters. As a consequence, the resulting XCO₂ and XCH₄ products are typically somewhat "noisier" compared to the OE products.

The XCO₂ and XCH₄ data products from SCIAMACHY are generated within the GHG-CCI project (Buchwitz et al.,

Table 1. Overview of SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT XCO₂ and XCH₄ level 2 data products (individual ground-pixel retrievals). For some products, level 3 (i.e. gridded data) products are also available (e.g. for CO₂_SCI_WFMD and CH₄_SCI_WFMD from http://www.iup.uni-bremen.de/sciamachy/NIR_NADIR_WFM_DOAS/ and merged SCIAMACHY and TANSO-FTS XCO₂ and XCH₄ products in Obs4MIPs format from <http://www.esa-ghg-cci.org/>).

| Variable | Sensor | Source Product (reference) |
|------------------|--------------------------|---|
| XCO ₂ | SCIAMACHY | http://www.esa-ghg-cci.org/ CO ₂ _SCI_BESD (Reuter et al., 2011) |
| | TANSO | http://www.gosat.nies.go.jp/en/ NIES operational GOSAT (Yoshida et al., 2013) http://www.esa-ghg-cci.org/ CO ₂ _GOS_OCFP (Cogan et al., 2012) CO ₂ _GOS_SRFP/RemoTeC (Butz et al., 2011) http://www.iup.uni-bremen.de/~heymann/besd_gosat.php GOSAT BESD (Heymann et al., 2015) http://disc.sci.gsfc.nasa.gov/acdisc/documentation/ACOS.html NASA ACOS (Crisp et al., 2012) |
| | SCIAMACHY & TANSO merged | http://www.esa-ghg-cci.org/ CO ₂ _EMMA (Reuter et al., 2013) http://disc.sci.gsfc.nasa.gov/OCO-2 NASA OCO-2 (Boesch et al., 2011) |
| XCH ₄ | SCIAMACHY | http://www.esa-ghg-cci.org/ CH ₄ _SCI_WFMD (Schneising et al., 2011) CH ₄ _SCI_IMAP (Frankenberg et al., 2011a) |
| | TANSO | http://www.gosat.nies.go.jp/en/ NIES operational GOSAT (Yoshida et al., 2013) http://www.esa-ghg-cci.org/ CH ₄ _GOS_OCPR (Parker et al., 2011) CH ₄ _GOS_SRPR/RemoTeC (Butz et al., 2010) CH ₄ _GOS_OCFP (Parker et al., 2011) CH ₄ _GOS_SRFP/RemoTeC (Butz et al., 2011) |
| | SCIAMACHY & TANSO merged | http://www.esa-ghg-cci.org/ CH ₄ _EMMA (Reuter et al., 2013) |

2015) of ESA's Climate Change Initiative (CCI, Hollmann et al., 2013) and these products are available from the GHG-CCI website (<http://www.esa-ghg-cci.org/>). XCO₂ and/or XCH₄ products from GOSAT are generated at several institutions in Japan, Europe and the USA and are available from several sources as shown in Table 1. The quality of these GHG-CCI products and the XCO₂ and XCH₄ products generated elsewhere has been significantly improved during recent years (e.g. Schneising et al., 2012; Yoshida et al., 2013; Dils et al., 2014; Buchwitz et al., 2015) and has now reached quite high maturity when compared to user requirements as formulated in, e.g. GCOS (2011). This can be concluded, for example, from the quality of the latest version of the GHG-CCI SCIAMACHY and TANSO XCO₂ and XCH₄ data set (Climate Research Data Package No. 3, CRDP3; Buchwitz et al., 2016). Based on comparisons with ground-based observations of the Total Carbon Column Observing Network (TCCON, Wunch et al., 2010, 2011) it has been found that the GCOS requirements for systematic er-

ror (< 1 ppm for XCO₂, < 10 ppb for XCH₄) and long-term stability (< 0.2 ppm/year for XCO₂, < 2 ppb yr⁻¹ for XCH₄) are met for nearly all products. As also shown in Buchwitz et al. (2016), the single observation (ground pixel) retrieval precision (random error primarily due to instrument noise) is about 2 ppm for XCO₂ from SCIAMACHY and TANSO and ≈ 15 ppb for TANSO XCH₄. For SCIAMACHY XCH₄ the precision depends on the time period and retrieval algorithm and is in the range 35–80 ppb. For some products it has also been investigated to what extent the uncertainty can be reduced upon averaging (Kulawik et al., 2016) and recommendations are given on how to take into account error correlations (Reuter et al., 2016), i.e. which values to use for the non-diagonal elements of the error covariance matrix, as an important contribution to the full characterisation of the data needs for data assimilation studies.

Figure 3 presents an overview of GHG-CCI CRDP3 XCO₂ (left) and XCH₄ (right) data set in terms of time series and maps. These figures have been generated by gridding the un-

derlying individual ground pixel (level 2) products to generate a $5^\circ \times 5^\circ$ monthly level 3 Obs4MIPs product (Buchwitz and Reuter, 2016). Each $5^\circ \times 5^\circ$ monthly grid cell also contains an estimate of the overall uncertainty (also shown in Fig. 3) which has been computed by taking into account random and systematic error components. The grid cell uncertainty is computed from two terms: (i) using the reported uncertainties as given in the level 2 (individual ground pixel) product files for each of the used satellite products (using an ensemble of SCIAMACHY and GOSAT level 2 products) and (ii) using a term accounting for potential regional/temporal biases as obtained from validation using TCCON ground-based data (see above). The first term depends on the number of individual observations added (the error reduces in proportion to the square root of the number of observations added) whereas the latter term is constant and in the range 0.57–0.87 ppm depending on the satellite XCO₂ product or in the range 6–10 ppb for XCH₄. As can be seen from Fig. 3, the uncertainty of the satellite XCO₂ retrievals for monthly $5^\circ \times 5^\circ$ averages is estimated to be typically around 0.5–1 ppm (values larger than 1 ppm are typically associated with regions where only few observations per grid cell exist, e.g. due to clouds or higher latitudes corresponding to low sun elevation). For XCH₄ the uncertainty is of the order of a few ppb (typically 4–8 ppb). In Buchwitz and Reuter (2016), initial TCCON validation results of the Obs4MIPs products are also presented. It is shown that the XCO₂ product agrees with monthly averaged TCCON XCO₂ within 0.29 ± 1.2 ppm (1σ) and the XCH₄ product within 2.0 ± 10.7 ppb. This is hardly worse than the results which have been obtained by careful validation of the individual ground pixel retrievals taking into account the best possible spatio-temporal co-location and considering the averaging kernels, etc. (e.g. Buchwitz et al., 2016). Note that the computed differences of Obs4MIPs monthly $5^\circ \times 5^\circ$ satellite products with monthly averaged TCCON include the errors of the satellite data, errors of the TCCON products, errors due to neglecting altitude sensitivity differences (averaging kernels) and representativity error. This indicates that the representativity error is quite small (at least for monthly $5^\circ \times 5^\circ$ spatio-temporal sampling and resolution), probably of the order of 0.1–0.2 ppm for XCO₂ and a few ppb for XCH₄ (it is planned to quantify this error in the future but currently only these rough estimates are available). Note that detailed information on all GHG-CCI products is available on the GHG-CCI website in terms of technical documents, links to peer-reviewed publications and figures including detailed maps for each month and each individual data product.

The SCIAMACHY and TANSO XCO₂ and XCH₄ retrievals have been used in a number of scientific studies to address important questions related to the sources and sinks of atmospheric CO₂ and CH₄ by atmospheric inversion studies (e.g. Bergamaschi et al., 2013; Houweling et al., 2015) and more recently also in a data assimilation context for optimising model parameters (Chevallier et al., 2017). Obviously,

the longer the time series and the more accurate it is, the larger the information content of a given data set. Therefore, further improvements are desired (Chevallier et al., 2017) and possible (at least in terms of time series extension but likely also in further reduction of remaining biases).

3.2.2 Reflectance-based vegetation dynamics/activity

Since the early beginnings of remote sensing the state and evolution of the vegetation has been monitored by satellites. An early attempt to analyse vegetation dynamics from space is to calculate the normalised difference vegetation index (NDVI), defined as the ratio between the difference of NIR and visible red (RED) spectral bands, and the sum of NIR and RED: $NDVI = (NIR - RED)/(NIR + RED)$ (Deering et al., 1975). The advantage of an index such as NDVI lies in its simplicity and applicability to sensors with few spectral bands such as the Advanced Very High Resolution Radiometer (AVHRR). Therefore this index has been applied for numerous purposes over the last 30 years or so. However NDVI is not a geophysical variable and it is sensitive to various perturbing factors such as atmospheric constituents (aerosols, water vapour), directional effects (geometry of illumination and observation), changes in soil background colour (depending on soil water content; e.g. Pinty et al., 1993; Goel and Qin, 1994; Leprieur et al., 1994; Dorigo et al., 2007). There have been many attempts to modify NDVI and develop additional vegetation indices (VIs) to overcome its limitations, for example the Soil-Adjusted Vegetation Index (Huete, 1988), Atmospherically Resistant Vegetation Index (Kaufman and Tanre, 1992) or Global Environmental Monitoring Index (Pinty and Verstraete, 1992). These indices generally exhibit some improvement in one respect but at the expense of degradation in another respect. Pinty et al. (2009) demonstrate the limitations of such VIs in representing the complex radiative properties of the canopy–soil system over the visible to NIR albedo range. Satellite-derived LAI products (e.g. Liu et al., 2014) seem to be an alternative to VIs. LAI is, however, model dependent; i.e. the correct interpretation of this variable depends on the formulation of the model used in the retrieval scheme and may differ from the interpretation adopted by the land biosphere model used for assimilating the LAI product (Disney et al., 2016).

A rational approach to addressing all these issues together is to design a physically based quantity which is determined by the state of the canopy–soil system. The fraction of absorbed photosynthetically active radiation (FAPAR), which is a normalised fraction with values ranging from 0 to 1, provides information on the photosynthetic activity of the land vegetation. It is recognised as an essential climate variable (ECV; GCOS, 2011) and is based on the land surface radiation budget. It is defined as the fraction of the photosynthetically active radiation (i.e. incoming solar radiation in the spectral region 0.4–0.7 µm) that is absorbed by the vegetation canopy (see also Pickett-Heaps et al., 2014 for

Table 2. Characteristics of a variety of FAPAR products; more details and products are provided by D’Odorico et al. (e.g. 2014); Pickett-Heaps et al. (e.g. 2014).

| Name | Time period | Temporal resolution | Definition | Reference |
|----------------|--------------|---------------------|---------------------------------------|-----------------------|
| MODIS | 2000–present | 8 days | Green canopy, direct radiation | (Myneni et al., 2002) |
| SeaWiFS* | 1997–2006 | 10 days | Green canopy, diffuse radiation | (Gobron et al., 2006) |
| TIP-MODIS | 2000–present | 16 days | FAPAR/green canopy, diffuse radiation | (Pinty et al., 2011b) |
| TIP-GlobAlbedo | 2002–2011 | 8 days | FAPAR/green canopy, diffuse radiation | (Disney et al., 2016) |
| Vegetation | 1999–present | 10 days | FAPAR, direct radiation | (Baret et al., 2007) |

* The same algorithm is used for MERIS (called JRC MGVI, 2002–2011) and SPOT-Vegetation (2012–present) with a 1.2 km, 10-day resolution (Gobron et al., 2008).




Figure 3. Time series of satellite-derived XCO₂ in three latitude bands (see annotation bottom left, e.g. red line: 30 to 60° N) and maps showing the spatial distribution of XCO₂ for April 2014 (top left) and corresponding XCO₂ uncertainty (top right). (b) As (a) but for XCH₄ (maps: September 2014).

a mathematical definition). Several FAPAR products are derived from a variety of optical sensors (e.g. ATSR, MERIS, MISR, MODIS, SEVIRI, SeaWiFS, VEGETATION) at different spatial and temporal resolutions. Although there have been substantial efforts to harmonise products across sensors (Ceccherini et al., 2013) and establish standards and validation practices (e.g. Widlowski, 2010), there are still considerable differences among the products. These differences can mainly be associated with differences in the retrieval methodology as well as to the quality of input variables. A recent overview of various FAPAR products and their specifications, but without an assessment of product uncertainties, is given by Gobron and Verstraete (2009). Table 2 summarises the characteristics of the most common FAPAR products.

Several studies have compared the performance of different satellite-derived FAPAR products: McCallum et al.

(2010) looked at four FAPAR data sets over northern Eurasia for the year 2000, Pickett-Heaps et al. (2014) evaluated six products across Australia, D’Odorico et al. (2014) compared three products over Europe, Tao et al. (2015) assessed five products over different land cover types, and Disney et al. (2016) compared two FAPAR products derived from Glob-Albedo and MODIS data. Pickett-Heaps et al. (2014) concluded that, although all six evaluated products display robust spatial and temporal patterns, there is considerable disagreement in the absolute magnitude among the products and none of the products outperforms the others. This has also been confirmed by the studies of D’Odorico et al. (2014) and Tao et al. (2015). One of the reasons for these differences are different assumptions on the underlying biome types. They also reviewed the consistency of the FAPAR products against in situ field measurements: the mean difference be-

tween the EO products and the in situ field measurements is around 0.1. This estimate is confirmed by the study of Tao et al. (2015), who suggest an average uncertainty of 0.14 from validation against total FAPAR and 0.09 from validation against green FAPAR in situ measurements. In their comparison of the Joint Research Centre – Two-stream Inversion Package (JRC-TIP) MODIS, JRC MGVI (based on MERIS) and Boston University MODIS products (see Table 2), D’Odorico et al. (2014) placed special emphasis on the assessment of the product uncertainties by not only comparing the uncertainties (or quality indicators) as proposed by the product teams but also by calculating an independent theoretical uncertainty based on the triple collocation (TC) method (see Sect. 3.2.4). While the uncertainties specified by the product teams differed by up to 0.1 among the products, the TC method suggested more consistent uncertainties among the three products of around 10–20 % of the signal.

The JRC-TIP (Pinty et al., 2007) is an inverse modelling system that was explicitly designed to retrieve a set of land surface variables, including FAPAR, in a form that is compliant with the requirements for assimilation into terrestrial biosphere models; hence we focus in the following on this product. TIP is based on a one-dimensional two-stream representation of the radiative transfer in the canopy–soil system (Pinty et al., 2006) and applies the same inversion approach as CCDAS, which is briefly sketched in Sect. 2.2 and detailed in Rayner et al. (2016) and Kaminski and Mathieu (2017). In a first step it retrieves a set of model parameters describing the state of the vegetation canopy system including the full uncertainty covariance of the parameters by combining prior information with observed radiant fluxes. Further, the model is used to propagate this PDF forward onto simulated fluxes such as FAPAR. TIP uses observed broadband albedo in the NIR and visible spectral domains as input. The prior information used in the retrieval is constant in space and time; i.e. all variability is determined from space (Kaminski et al., 2017). This is in contrast to other retrieval approaches, which are based on prescribed land cover maps (e.g. Liu et al., 2014). Long-term global records of JRC-TIP products (see Table 2) have been retrieved from broadband albedos provided by MODIS collection 5 (Pinty et al., 2011b, c) and Globalbedo (Disney et al., 2016). Products are provided for each of the respective 16-day (MODIS) and 8-day (Globalbedo) synthesis periods. To reduce disk space, by default, JRC-TIP products are delivered without correlations among the uncertainties between individual variables, even though these correlations are available. An estimate of uncertainty correlation in space and time is not provided. Both JRC-TIP records are provided in the native 1 km resolution of the albedo input products. In order to maintain the above-mentioned compliance with terrestrial models, coarser-resolution products are to be derived by applying JRC-TIP to aggregated albedo inputs (as in Disney et al., 2016). JRC-TIP products are validated at site (Pinty et al., 2007, 2008, 2011a) and regional

scales (Disney et al., 2016); more details on JRC-TIP are given in Kaminski et al. (2017).

3.2.3 Biogeochemical-based vegetation activity

Sun-induced fluorescence (SIF) is an electromagnetic signal emitted as a two-peak spectrum between 650 and 850 nm by the chlorophyll *a* of green plants under solar radiation. SIF can be directly related to photosynthetic electron transport rates and yields a mechanistic link to photosynthesis and the subsequent gross carbon uptake by terrestrial vegetation (GPP; Porcar-Castell et al., 2014). Recent developments in satellite-based spectroscopy have enabled the first retrievals of SIF from space (Frankenberg et al., 2011c; Joiner et al., 2011), which holds the promise of enabling new approaches to globally monitoring terrestrial photosynthesis. For example, a high linear correlation between data-driven GPP estimates and SIF retrievals at global and annual scales was reported by Frankenberg et al. (2011c); Guanter et al. (2012). The skills of SIF as a proxy for photosynthetic activity and GPP were also reported by studies over different ecosystems, like the Amazon rainforest (Lee et al., 2013; Parazoo et al., 2013), large crop belts (Guanter et al., 2014) and the boreal forests in Eurasia and North America (Walther et al., 2015). However, in the context of DA and in order to extract the maximal benefit from SIF data, the complex processes responsible for SIF in the plants’ photochemical systems (as mentioned above) require complex models as observation operators for SIF.

The global retrieval of SIF from space relies on the principle of in-filling of solar Fraunhofer lines by SIF (Frankenberg et al., 2011b). Fraunhofer lines are absorption features in the solar spectrum, caused by elements in the solar atmosphere and sufficiently resolved by atmospheric spectrometers. Because of the additive nature of SIF, the fractional depth of the Fraunhofer lines detected by the satellite instrument decreases with the amount of SIF being emitted at the same wavelength. The retrieval of SIF from space is then based on the evaluation of the depth of the Fraunhofer lines present in red and NIR top-of-atmosphere spectra. The retrieval forward model is thus simple and can be linearised (e.g. Guanter et al., 2012; Köhler et al., 2015b), which simplifies the inversion.

Fraunhofer line-based SIF retrievals tend to be accurate but not precise: uncertainties are dominated by a random component associated to instrumental noise, which is linearly mapped into SIF retrievals. The amplitude of instrumental noise, and hence 1σ single-retrieval errors, scale with at-sensor radiance for the most common case of grating-based spectrometers dominated by multiplicative noise. This implies that retrieval errors are mostly driven by surface brightness and sun zenith angles (Guanter et al., 2015). Because of this high contribution of random errors to the total retrieval uncertainty, single SIF retrievals are commonly linearly aggregated as spatio-temporal composites in which

Table 3. Selected characteristics of operating and planned space-borne instruments able to deliver SIF data. Names of upcoming instruments are highlighted in italics. NIR stands for near-infrared. It must be noted that GOME-2 on MetOp-A has been operating with a reduced pixel size of $40 \times 40 \text{ km}^2$ since July 2013. References are examples: the full list is given in the text.

| Name | Time period | Overpass time | Spectral sampling | Spatial resolution | Temporal resolution | Reference |
|----------------|-------------|---------------|-------------------|-------------------------------|---------------------|--|
| GOSAT | 2009–today | Midday | NIR | 10 km diam. | 3 days | e.g. (Frankenberg et al., 2011c) |
| GOME-2 | 2007–today | Morning | red & NIR | $40 \times 80 \text{ km}^2$ | < 2 days | e.g. (Joiner et al., 2013), (Köhler et al., 2015b) |
| SCIAMACHY | 2003–2012 | Morning | red & NIR | $30 \times 240 \text{ km}^2$ | < 3 days | e.g. (Joiner et al., 2013), (Köhler et al., 2015b) |
| OCO-2 | 2014–today | Midday | NIR | $1.3 \times 2.3 \text{ km}^2$ | 16 days | (Frankenberg et al., 2014) |
| <i>TROPOMI</i> | ~ 2017 | Midday | red & NIR | $7 \times 7 \text{ km}^2$ | < 1 day | (Guanter et al., 2015) |
| <i>FLEX</i> | ~ 2022 | Morning | red & NIR | $0.3 \times 0.3 \text{ km}^2$ | < 27 days | (Drusch et al., 2017) |

random errors are reduced. The number of retrievals to be aggregated into a given grid cell results from a compromise between spatial resolution, temporal resolution and precision of the gridded product, the size of the spatial and temporal bins being exchangeable in terms of their effect on the random uncertainty. The random uncertainty of the resulting spatio-temporal composites is then not only driven by surface albedo and illumination, but also by the number of soundings going into a given grid cell, which is in turn defined by cloudiness and latitude (in the case of overlapping orbits). Detailed analyses of random errors in SIF retrievals for different space-borne instruments can be found in Frankenberg et al. (2011b) and Guanter et al. (2015).

Global SIF data sets have been or are being derived from GOSAT, MetOp's Global Ozone Monitoring Experiment-2 (GOME-2), ENVISAT's SCIAMACHY and the OCO-2 mission (Joiner et al., 2011, 2012, 2013, 2016; Frankenberg et al., 2011c, 2014; Guanter et al., 2012; Köhler et al., 2015a, b; Wolanin et al., 2015). All four missions except for SCIAMACHY are still operating. Sample SIF maps from GOSAT, GOME-2 and SCIAMACHY for July 2010 are displayed in Fig. 4. The spectral, spatial and temporal sampling of single SIF soundings varies for each instrument, as it is summarised in Table 3. For example, GOME-2 and SCIAMACHY provide SIF retrievals in the red and NIR spectral regions with global coverage and a relatively high temporal resolution. However, this comes at the expense of a coarse spatial resolution, which is $40 \times 80 \text{ km}^2$ for GOME-2 ($40 \times 40 \text{ km}^2$ for GOME-2 on MetOp-A since July 2013) and $30 \times 240 \text{ km}^2$ for SCIAMACHY. On the other hand, GOSAT and OCO-2 do not provide spatially continuous measurements (i.e. no global coverage), but single soundings in the NIR have a much higher spatial resolution than those of GOME-2 and SCIAMACHY. In particular, OCO-2 soundings correspond to ground areas of about 4 km^2 , which is substantially finer than that of the other data sets. The number of soundings per day by OCO-2 is also much larger (about $100\times$) than that by the other instruments (Frankenberg et al., 2014), which makes OCO-2 SIF to be the most suited data set for studies over areas not requiring a continuous spatial sampling but

benefiting from a high spatial resolution. This is the case, for example, of tropical and boreal forests: spatial continuity is less critical for those ecosystems because they are relatively homogeneous over large spatial scales, whereas the high spatial resolution is important for maximising the number of clear-sky soundings during the parts of the year with persistent cloudiness.

Concerning near-future perspectives for SIF monitoring, it can be expected that the limitations in spatial resolution and coverage of existing SIF products will be alleviated with the advent of the TROPOspheric Monitoring Instrument (TROPOMI) scheduled for launch on board the Sentinel-5 Precursor satellite mission by mid-2017 (Table 3). TROPOMI will enable SIF retrievals in the red and NIR regions similar to GOME-2 and SCIAMACHY, but with a 7 km pixel, daily global coverage and a number of clear-sky observations per day, ≈ 200 times larger than GOME-2 and ≈ 600 times larger than SCIAMACHY. The SIF product from TROPOMI can therefore be anticipated to have a much higher spatio-temporal resolution and signal-to-noise ratio than those from GOME-2 and SCIAMACHY (Guanter et al., 2015). Complementarily, the FLuorescence EXplorer (FLEX; Drusch et al., 2017) has recently been selected for implementation by ESA, with launch currently expected for 2022. FLEX will provide global measurements of SIF in the red and NIR at a relatively low temporal resolution, but with the finest spatial resolution of all existing and upcoming space-borne instruments.

3.2.4 Soil moisture

Soil moisture is measured in situ through large-scale soil moisture monitoring networks (Dorigo et al., 2011; Ochsner et al., 2013) or at various FLUXNET sites (Baldochini et al., 2001). Yet, these point observations have only limited coverage in space time, have spatially very divergent properties (Dorigo et al., 2013), and often contain large representativeness errors at the scale of global ecosystem models (Gruber et al., 2013). Satellite remote sensing in the microwave domain has the potential to overcome many of these

issues. Microwave remote sensing uses the contrasting dielectric properties of water, air, ice and soil particles to infer the water content in the soil column (Owe et al., 2008). Both passive radiometer systems, measuring the emitted microwave radiance (brightness temperatures), and active radar systems, measuring backscattered microwave radiance, can be used to retrieve soil moisture. Various approaches exist that convert brightness temperatures and backscatter measurements into estimates of soil moisture, including radiative transfer model inversion approaches (e.g. Kerr et al., 2012; Owe et al., 2008), neural networks (e.g. Rodriguez-Fernandez et al., 2015), linear regressions (Al-Yaari et al., 2016, e.g.) and change detection methods (Wagner et al., 1999). The latter is commonly applied to scatterometer measurements and yields, in contrast to the other approaches which provide soil moisture as volumetric water content, soil moisture as a percentage of total saturation. Microwave sensors operate in different frequency (wavelength) domains, of which L-band (with a wavelength of ≈ 23 cm) and C-band (≈ 5 cm) are most commonly used for retrieving soil moisture (Kerr et al., 2012; Owe et al., 2008; Wagner et al., 1999). Smaller wavelengths are more sensitive to the vegetation canopy covering the soil and increasingly lose their sensitivity to water. Still, frequencies up to 19 GHz (≈ 1.5 cm) have proven potential for providing robust soil moisture estimates at the global scale for moderately to sparsely vegetated areas (Owe et al., 2008). Due to the relatively low energy levels and the technical challenges in microwave domain, spatial resolutions of the satellite observations are generally coarse ($\approx 25\text{--}50$ km) but with high revisit frequencies (up to 1 day). Only synthetic aperture radar is able to provide much higher spatial resolutions, up to a few tens of metres, yet at the cost of long revisit times. Also observations made by the Gravity Recovery and Climate Experiment (GRACE; Rodell et al., 2009) are sensitive to soil moisture, but the estimation of soil moisture content from these observations is not straightforward because they are also sensitive to changes in snow, surface water, groundwater and vegetation.

Since the release of the first global soil moisture data sets from microwave sensors in the early 2000s, the number of available soil moisture products and missions has rapidly expanded (De Jeu and Dorigo, 2016). Several (pre-)operational products are now available from a wide variety of data providers and space organisations (Table 4). While initially soil moisture products were based on sensors mainly designed for other purposes (such as ASCAT, AMSR2 and Sentinel-1), ESA and NASA launched their own dedicated soil moisture satellite missions SMOS and SMAP (Kerr et al., 2012; Entekhabi et al., 2010). Differences between the various products exist in their technical design, observation bands and retrieval algorithms, which often result in complementary strengths over different land cover types (Alyaari et al., 2015; Dorigo et al., 2010; Liu et al., 2011). The missions also differ in their degree of operationalisation: while SMOS and SMAP are primarily scientific concept demon-

strators, AMSR2 continues the legacy of C-band radiometer observations started by JAXA and NASA in 2002 with the launch of AMSR-E, while ASCAT is embedded in a fully operational programme of weather observing satellites with guaranteed continuation at least until 2023 and a follow-on mission already under development (Wagner et al., 2013). Apart from the target variable surface soil moisture, some products come with estimates of freeze/thaw state and VOD, which are disentangled from the soil moisture impacts on the measured microwave signal during the retrieval process.

As none of the currently active missions cover a period long enough to study climate change impacts, ESA's Climate Change Initiative (CCI) endorsed the combination of available soil moisture products from active and passive microwave sensors into a consistent multidecadal record. The ESA CCI soil moisture product currently combines soil moisture products from 11 different sensors into a homogenised daily product spanning the period 1978–2015 (Liu et al., 2012, 2011; Dorigo et al., 2016). Several studies have demonstrated the value of ESA CCI soil moisture for assessing long-term interactions between soil moisture and vegetation productivity (Barichivich et al., 2014; Chen et al., 2014; Dorigo et al., 2012; Muñoz et al., 2014).

Key to a proper assimilation of remotely sensed soil moisture into carbon models is a correct characterisation of its errors. Apart from instrument errors which are common to all observations, the quality of microwave-based soil moisture retrievals is particularly impacted by vegetation cover, soil frost, snow cover, open water, topography, surface roughness, urban structures and radio frequency interference (Dorigo et al., 2010; Kerr et al., 2012). Observations in which a strong adverse impact of these factors is detected are usually masked during processing, which may lead to data gaps for certain areas or periods of the year (Dorigo et al., 2015). If cases in which their impact on the soil moisture retrieval are only moderate, the errors that they introduce are either simulated during the retrieval itself using error propagation methods, or assessed *a posteriori* against reference data using various statistical methods (Draper et al., 2013).

While the ASCAT and AMSR2 products come with an estimate of the error variance for each observation by error propagation (Naeimi et al., 2009; Parinussa et al., 2011), this is still not common practice for all soil moisture products. Yet, no error propagation model perfectly represents all error sources and interactions (Draper et al., 2013). On the other hand, the use of *in situ* soil moisture measurements to estimate random errors is hampered by their heterogeneous nature and large spatial representativeness errors (Gruber et al., 2013). As an alternative, in recent years triple collocation analysis (TCA) has firmly established itself as a robust alternative to estimate random errors in soil moisture data sets without the need of an absolute “true” reference (Dorigo et al., 2010; Scipal et al., 2008). TCA estimates the error variances of three spatially and temporally collocated soil moisture data sets with independent error structures, e.g.




Figure 4. Maps of sun-induced fluorescence (SIF) for July 2010 derived from GOSAT, GOME-2 and SCIAMACHY satellite data.

Table 4. Current (pre-)operational global soil moisture missions and products (see the list of acronyms).

| Mission | Organisation | Measurement concept | Band | Mission start | Data access |
|---------------|-----------------|---------------------------------------|--------------------|---------------|---|
| MetOp – ASCAT | EUMETSAT | Real aperture radar (scatterometer) | C-band | Jan. 2007 | http://hsaf.meteoam.it/soil-moisture.php |
| SMOS | ESA | Interferometric radiometer | L-band | Nov. 2009 | http://land.copernicus.eu/global/products/swi |
| GCOM-W1 | JAXA | Radiometer | C-band | May 2012 | http://www.vandersat.com/ |
| AMSR2 | | | | | http://suzaku.eorc.jaxa.jp/GCOM_W/ |
| SMAP | NASA | Radiometer & radar* | L-band | Jan. 2015 | http://smap.jpl.nasa.gov/ |
| Sentinel-1 | ESA/ Copernicus | Synthetic aperture radar | C-band | Apr. 2014 | https://www.eodc.eu/ |
| CCI | ESA | Combined scatterometer and radiometer | L-, C-, X- Ku-band | Nov. 1978 | http://www.esa-soilmoisture-cci.org |

* SMAP's radar failed in July 2015.

a radiometer-based, a scatterometer-based and a land surface model soil moisture data set. Recently, the TCA has been intensively elaborated, e.g. to solve for colinearities between errors (Gruber et al., 2016b) and non-linear dependencies between data sets (Zwieback et al., 2016). The most remarkable advancement has been to express TCA-based error estimates as a signal-to-noise ratio, which facilitates a direct intercomparison of the skill of data sets independent of their dynamic ranges (Gruber et al., 2016a); see Fig. 5. Although the TCA provides an estimate that is entirely independent of any retrieval model assumptions, it only provides a single average error estimate for the entire period under consideration.

Thus, synergistic use of error propagation and triple collocation estimates shall be exploited to better capture the temporal error dynamics needed for an optimal assimilation into carbon models. Due to the recent progress in product quality, error characterisation and operationalisation, satellite-based soil moisture products have reached the level of maturity that allows for a systematic assimilation into land surface models. These products have been used to improve model hydrology by, for example, Martens et al. (2017) who showed that the assimilation of SMOS and ESA CCI soil moisture generally has a small positive impact on soil water storage and evaporative fluxes as simulated by the GLEAM land evapo-

ration model. Surface soil moisture from ASCAT is assimilated operationally in near-real-time into the ECMWF Land Data Assimilation System to obtain root-zone soil moisture (Albergel et al., 2012). Global root-zone soil moisture products based on SMOS and SMAP are derived by a slightly different approach, which assimilate the observed brightness temperatures instead of the retrieved surface soil moisture products (Lannoy and Reichle, 2016). The assimilation of satellite-based soil moisture products in terrestrial carbon cycle models has been described above.

3.2.5 Biomass

Continental-scale biomass maps have been produced from space using both radar and lidar. Biomass here refers to above-ground biomass (AGB), since there are no methods to measure the below-ground component, and this is typically inferred from AGB using allometric equations. Furthermore, the emphasis is on the AGB of forests, although a global data set of AGB in all biomes for the period 1993–2012 has been produced based on VOD data from global passive microwave sensors, hence with spatial resolution of 10 km or coarser (Liu et al., 2015). The AGB product is derived from a regression of VOD against observations of AGB from ground-based inventory data.

Using long time series of C-band radar data provided by the ESA Envisat satellite, the growing stock volume of Northern Hemisphere boreal and temperate forests has been estimated (Santoro et al., 2011). Although available at 0.01° resolution, the accuracy of growing stock volume at this scale is comparatively poor, and spatial averaging provides more reliable results: at 0.5° spacing, estimated growing stock volume has a relative accuracy of 20–30 % when tested against inventory data (Santoro et al., 2013). Thurner et al. (2014) used this product to derive the carbon stock (above- and below-ground) in boreal, temperate mixed and broadleaf and temperate coniferous forests of forests above 30° N (40.7, 24.5 and 14.5 PgC respectively). These values have estimated accuracies of around 33–39 % under a conservative approach to estimate uncertainty. Santoro et al. (2015) provide a high-resolution data set (0.01°) over the Northern Hemisphere with a relative RMSE against National Forest Inventory between 12 and 45 %.

For tropical forests, the key sensor is the Geoscience Laser Altimeter System (GLAS) on board the Ice, Cloud and land Elevation Satellite (ICESat) which failed in 2009 (Lefsky, 2010). Its archive of forest height estimates was the core data set exploited to produce two pantropical biomass maps (Saatchi et al., 2011; Baccini et al., 2012) at grid scales of 1 km and 500 m respectively; Saatchi et al. (2011) also provide a map of the errors associated with the biomass estimates at each pixel. This is produced by combining measurement errors, allometry errors, sampling errors and prediction errors, which are treated as independent and spatially uncorrelated. Further details are given in the supplementary

material to Saatchi et al. (2011). In an attempt to resolve differences between these two maps, Avitabile et al. (2016) used an independent reference data set of field observations to remove the biases in the maps and then combined them to estimate the AGB in the tropical belt (23.4° S to 23.4° N). Testing against a reference data set not used in the fusion process indicated that the fused map had a RMSE 15–21 % lower than that of the input maps and nearly unbiased estimates.

However, there are unresolved questions about large-scale biomass patterns across the Amazon inferred from in situ and satellite data. Biomass maps derived from satellite data in Saatchi et al. (2011) and Baccini et al. (2012) differ significantly from each other and from biomass maps derived from in situ plots distributed across Amazonia using kriging (Mitchard et al., 2014). Neither satellite product exhibits the strong increase in biomass from southwestern to northeastern Amazonia inferred from in situ data. Mitchard et al. (2014) attributed this to failure to account for gradients in wood density and regionally varying tree height–diameter relations when estimating biomass from the satellite data. Saatchi et al. (2015) reject this analysis and claim that the trends and patterns in Mitchard et al. (2014) are erroneous and a consequence of inadequate sampling. Resolving this disagreement is of fundamental importance since it raises basic questions on accuracy, uncertainty and representativeness for both in situ and satellite-derived biomass data.

The next 4–5 years will dramatically improve our global knowledge of biomass, with the launch of three missions aimed at measuring forest structure and biomass. The ESA BIOMASS mission (European Space Agency, 2012), to be launched in 2021, is a P-band radar that will provide near-global measurements of forest biomass and height. Measurements from airborne sensors indicate that even in dense tropical forests affected by topography, the P-band frequency used by BIOMASS will give sensitivity to biomass up to 350–450 t ha⁻¹ (Minh et al., 2014; Villard and Toan, 2015). Around the same time the NASA-ISRO SAR mission (NISAR) based on an L-band sensor will be deployed, providing measurements of biomass in lower biomass forests (up to 100 t ha⁻¹). These highly complementary missions will be further complemented by the NASA Global Ecosystem Dynamics Investigation vegetation lidar to be placed on the International Space Station around 2019; this aims to provide the first global, high-resolution observations of the vertical structure of tropical and temperate forests, from which biomass may be estimated.

As well as limitations caused by mission lifetimes, satellite measurements of biomass are unlikely to be sensitive enough to measure biomass increment except in rapidly growing plantations and tropical forests. Hence an important ancillary data set for studies aiming to relate biomass to climate and environment is tree ring data (<https://www.ncdc.noaa.gov/data-access/paleoclimatology-data/datasets/tree-ring>).




Figure 5. Signal-to-noise ratio (in dB), estimated with the triple collocation analysis for four different satellite-based soil moisture products and a land surface model. (a) MetOp-A ASCAT based on the TU Wien method (Wagner et al., 1999); (b) AMSR2 based on the LPRM model (Owe et al., 2008); (c) SMOS L3 (Kerr et al., 2010); (d) SMAP (Jackson, 1993). An SNR of -3 indicates a signal variance that is half of the noise variance, an SNR of 0 a signal variance equal to the noise variance, an SNR of 3 a signal variance that is twice the noise variance and so on. In areas without data the TC could not be computed, e.g. because of too few observations in one of the data sets. For details see Gruber et al. (2016b).

4 Conclusions

In the context of carbon cycle data assimilation this paper reviews the requirements and summarises the availability and characteristics of some selected observations with a special focus on remotely sensed Earth observation data. Observations are key for understanding the carbon cycle processes and are an important component for any data assimilation system. In this context the provision of systematic and sustained observing systems on an operational basis is becoming more and more important.

An example for such an operational network for in situ data is the Integrated Carbon Observing System (ICOS; see also <https://www.icos-ri.eu>). ICOS is a pan-European infrastructure for carbon observations, which provides high-quality in situ observations (both fluxes as well as atmospheric concentrations) over Europe and over ocean regions adjacent to Europe with a long-term perspective. ICOS consists of central facilities for co-ordination, calibration and data in conjunction with networks of atmospheric, oceanic and ecosystem observations as well as a data distribution centre, the Carbon Portal, providing discovery of and access to ICOS data products such as derived flux information. Other (quasi-)operational networks measuring atmospheric CO₂ concentrations are maintained, for instance, by the National Oceanic and Atmospheric Administration (NOAA) Climate

Monitoring and Diagnostics Laboratory and the Scripps Institution of Oceanography, both USA, as well as the CSIRO Global Atmospheric Sampling Laboratory, Australia.

An example for an operational space-based Earth observing programme in Europe is the fleet of so-called Sentinel satellites of the Copernicus programme. Copernicus aims to provide Europe with continuous and independent access to Earth observation data and associated services (transforming the satellite and additional in situ data into value-added information by processing and analysing the data) in support of Earth system science (Berger et al., 2012). Currently, six different Sentinel missions are planned (and have partly been launched). So far, a dedicated mission for monitoring the carbon cycle, i.e. an instrument measuring the atmospheric CO₂ composition, is not yet included in the Copernicus monitoring programme (see Ciais et al., 2015); however, the series of Sentinel satellites will be extended in the future and will likely include a CO₂ mission. Other operational EO programmes are operated by e.g. NOAA and the Japanese Aerospace Exploration Agency.

The paper also briefly recapitulates the assimilation systems capable of integrating these data: a more comprehensive description of the underlying formalism is given in Rayner et al. (2016) while MacBean et al. (2016) discuss the implementation strategies for a multiple data assimilation system and their impacts on the results. To take maximum advantage

of these data streams in carbon cycle data assimilation studies it is of utmost importance to have the appropriate knowledge of the uncertainty characteristics of the observational data, here with a focus on satellite products. This includes an understanding of the observable and its representativeness in order to develop the appropriate observation operator (see also Kaminski and Mathieu, 2017) but also the structure of any biases, random errors and error covariances (that is both the diagonal and off-diagonal elements quantifying the error correlations between different observations).

The benefit of using multiple data streams in a CCDAS lies in the complementarity of the data and thus in the ability to constrain different components of the underlying process model. In fact, because of the model internal interactions and feedbacks among the components the simultaneous assimilation of complementary observations has synergistic effects such that the constraint is larger than the sum of the individual constraints, as shown for instance by Kato et al. (2013), who assimilated observations of FAPAR and latent heat flux.

As a final remark one important aspect of observational data is their continuity, since much of the important information is contained in response to climate anomalies. Fortunately, the set up of operational observing systems such as ICOS for in situ data or Copernicus for satellite data has created the necessary infrastructure to ensure a long-term perspective in the provision of Earth observations.

Data availability. No data sets were used in this article.

Appendix A: List of acronyms

| | |
|-----------|---|
| ACE-FTS | Atmospheric Chemistry Experiment – Fourier transform spectrometer |
| AGB | Above-ground biomass |
| AIRS | Atmospheric Infrared Sounder |
| AMSR2 | Advanced Microwave Scanning Radiometer 2 |
| AMSR-E | Advanced Microwave Scanning Radiometer – Earth observing system |
| ASCAT | Advanced scatterometer |
| ATSR | Along Track Scanning Radiometers |
| AVHRR | Advanced Very High Resolution Radiometer |
| CCDAS | Carbon cycle data assimilation system |
| CCI | Climate Change Initiative |
| ECMWF | European Centre for Medium-Range Weather Forecasts |
| ECV | Essential climate variable |
| EO | Earth observation (in this form generally understood as from space) |
| ESA | European Space Agency |
| FAPAR | Fraction of absorbed photosynthetically active radiation |
| FLEX | FLuorescence EXplorer |
| GCOM-W1 | Global Change Observation Mission 1st-Water |
| GLAS | Geoscience Laser Altimeter System |
| GLEAM | Global Land Evaporation Amsterdam Model |
| GOME-2 | Global Ozone Monitoring Experiment-2 |
| GOSAT | Greenhouse Gases Observing Satellite |
| GPP | Gross primary productivity |
| IASI | Infrared Atmospheric Sounding Interferometer |
| ICOS | Integrated Carbon Observing System |
| ICESat | Ice, Cloud and land Elevation Satellite |
| ISRO | Indian Space Research Organisation |
| JAXA | Japan Aerospace Exploration Agency |
| JRC-MGVI | Joint Research Centre – MERIS Global Vegetation Index |
| JRC-TIP | Joint Research Centre – Two-stream Inversion Package |
| LAI | Leaf area index |
| MERIS | Medium Resolution Imaging Spectrometer |
| MIPAS | Michelson Interferometer for Passive Atmospheric Sounding |
| MISR | Multi-angle Imaging SpectroRadiometer |
| MODIS | Moderate Resolution Imaging Spectroradiometer |
| NASA | National Aeronautics and Space Administration |
| NDVI | Normalised difference vegetation index |
| NIR | Near infrared |
| NOAA | National Oceanic and Atmospheric Administration |
| Obs4Mips | Observations for Model Intercomparisons Project |
| OCO-2 | Orbiting Carbon Observatory 2 |
| OE | Optimal estimation |
| PDF | Probability density function |
| SAR | Synthetic aperture radar |
| SCIAMACHY | SCanning Imaging Absorption spectrometer for Atmospheric CHartography |
| SeaWiFS | Sea-viewing Wide Field-of-view Sensor |
| SEVIRI | Spinning Enhanced Visible and InfraRed Imager |
| SIF | Sun-induced fluorescence |
| SMAP | Soil Moisture Active Passive |
| SMOS | Soil Moisture Ocean Salinity |
| SWIR | Shortwave infrared |
| TANSO-FTS | Thermal And Near infrared Sensor for carbon Observations – Fourier Transform Spectrometer |
| TCA | Triple collocation analysis |
| TCCON | Total Carbon Column Observing Network |
| TCOS | Terrestrial Carbon Observation System |
| TROPOMI | TROPOspheric Monitoring Instrument |
| VI | Vegetation index |
| VOD | Vegetation optical depth |

Competing interests. The authors declare that they have no conflict of interest.

Special issue statement. This article is part of the special issue “Data assimilation in carbon/biogeochemical cycles: consistent assimilation of multiple data streams” (BG/ACP/GMD interjournal SI). It is not associated with a conference.

Acknowledgements. Michael Buchwitz has received funding from ESA via the GHG-CCI project. Wouter Dorigo is supported by the TU Wien Wissenschaftspris 2015, a personal grant awarded by the Vienna University of Technology. Figure 4 was kindly provided by Philipp Köhler, California Institute of Technology. We acknowledge the support from the International Space Science Institute (ISSI). This publication is an outcome of the ISSI’s working group on “carbon cycle data assimilation: how to consistently assimilate multiple data streams”.

Edited by: Victor Brovkin

Reviewed by: Natasha MacBean and Thomas Kaminski

References

- Al-Yaari, A., Wigneron, J., Kerr, Y., de Jeu, R., Rodriguez-Fernandez, N., van der Schalie, R., Bitar, A. A., Mialon, A., Richaume, P., Dolman, A., and Ducharne, A.: Testing regression equations to derive long-term global soil moisture datasets from passive microwave observations, *Remote Sens. Environ.*, 180, 453–464, <https://doi.org/10.1016/j.rse.2015.11.022>, 2016.
- Albergel, C., de Rosnay, P., Gruhier, C., Muñoz Sabater, J., Hassenauer, S., Isaksen, L., Kerr, Y., and Wagner, W.: Evaluation of remotely sensed and modelled soil moisture products using global ground-based in situ observations, *Remote Sens. Environ.*, 118, 215–226, <https://doi.org/10.1016/j.rse.2011.11.017>, 2012.
- Albergel, C., Munier, S., Leroux, D. J., Dewaele, H., Fairbairn, D., Barbu, A. L., Gelati, E., Dorigo, W., Faroux, S., Meurey, C., Le Moigne, P., Decharme, B., Mahfouf, J.-F., and Calvet, J.-C.: Sequential assimilation of satellite-derived vegetation and soil moisture products using SURFEX_v8.0: LDAS-Monde assessment over the Euro-Mediterranean area, *Geosci. Model Dev. Discuss.*, <https://doi.org/10.5194/gmd-2017-121>, in review, 2017.
- Alyouri, A., Wigneron, J. P., Ducharne, A., Kerr, Y., Wagner, W., De Lannoy, G., Reichle, R., Al Bitar, A., Dorigo, W., Richaume, P., and Mialon, A.: Global-scale comparison of passive (SMOS) and active (ASCAT) satellite-based microwave soil moisture retrievals with soil moisture simulations (MERRA-Land), *Remote Sens. Environ.*, 152, 614–626, <https://doi.org/10.1016/j.rse.2014.07.013>, 2015.
- Avitabile, V., Herold, M., Heuvelink, G. B. M., Lewis, S. L., Phillips, O. L., Asner, G. P., Armston, J., Ashton, P. S., Banin, L., Bayol, N., Berry, N. J., Boeckx, P., de Jong, B. H. J., DeVries, B., Girardin, C. A. J., Kearsley, E., Lindsell, J. A., Lopez-Gonzalez, G., Lucas, R., Malhi, Y., Morel, A., Mitchard, E. T. A., Nagy, L., Qie, L., Quinones, M. J., Ryan, C. M., Ferry, S. J. W., Sunderland, T., Laurin, G. V., Gatti, R. C., Valentini, R., Verbeeck, H., Wijaya, A., and Willcock, S.: An integrated pan-tropical biomass map using multiple reference datasets, *Glob. Change Biol.*, 22, 1406–1420, <https://doi.org/10.1111/gcb.13139>, 2016.
- Baccini, A., Goetz, S. J., Walker, W. S., Laporte, N. T., Sun, M., Sulla-Menashe, D., Hackler, J., Beck, P. S. A., Dubayah, R., Friedl, M. A., Samanta, S., and Houghton, R. A.: Estimated carbon dioxide emissions from tropical deforestation improved by carbon-density maps, *Nature Climate Change*, 2, 182–185, <https://doi.org/10.1038/nclimate1354>, 2012.
- Baldocchi, D., Falge, E., Gu, L., Olson, R., Hollinger, D., Running, S., Anthoni, P., Bernhofer, C., Davis, K., Evans, R., Fuentes, J., Goldstein, A., Katul, G., Law, B., Lee, X., Malhi, Y., Meyers, T., Munger, W., Oechel, W., Paw, K. T., Pilegaard, K., Schmid, H. P., Valentini, R., Verma, S., Vesala, T., Wilson, K., and Wofsy, S.: FLUXNET: A New Tool to Study the Temporal and Spatial Variability of Ecosystem-Scale Carbon Dioxide, Water Vapor, and Energy Flux Densities, *B. Am. Meteorol. Soc.*, 82, 2415–2434, [https://doi.org/10.1175/1520-0477\(2001\)082<2415:FANTTS>2.3.CO;2](https://doi.org/10.1175/1520-0477(2001)082<2415:FANTTS>2.3.CO;2), 2001.
- Barbu, A. L., Calvet, J.-C., Mahfouf, J.-F., and Lafont, S.: Integrating ASCAT surface soil moisture and GEOV1 leaf area index into the SURFEX modelling platform: a land data assimilation application over France, *Hydrol. Earth Syst. Sci.*, 18, 173–192, <https://doi.org/10.5194/hess-18-173-2014>, 2014.
- Baret, F., Hagolle, O., Geiger, B., Bicheron, P., Miras, B., Huc, M., Berthelot, B., Nino, F., Weiss, M., Samain, O., Roujean, J. L., and Leroy, M.: LAI, fAPAR and fCover CYCLOPES global products derived from VEGETATION: Part 1: Principles of the algorithm, *Remote Sens. Environ.*, 110, 275–286, <https://doi.org/10.1016/j.rse.2007.02.018>, 2007.
- Barichivich, J., Briffa, K. R., Myneni, R., Van der Schrier, G., Dorigo, W., Tucker, C. J., Osborn, T., and Melvin, T.: Temperature and Snow-Mediated Moisture Controls of Summer Photosynthetic Activity in Northern Terrestrial Ecosystems between 1982 and 2011, *Remote Sensing*, 6, 1390–1431, <https://doi.org/10.3390/rs6021390>, 2014.
- Barrett, D. J.: Steady state turnover time of carbon in the Australian terrestrial biosphere, *Global Biogeochem. Cy.*, 16, 1108, <https://doi.org/10.1029/2002GB001860>, 2002.
- Bergamaschi, P., Houweling, S., Segers, A., Krol, M., Frankenberg, C., Scheepmaker, R. A., Dlugokencky, E., Wofsy, S. C., Kort, E. A., Sweeney, C., Schuck, T., Brenninkmeijer, C., Chen, H., Beck, V., and Gerbig, C.: Atmospheric CH₄ in the first decade of the 21st century: Inverse modeling analysis using SCIAMACHY satellite retrievals and NOAA surface measurements, *J. Geophys. Res.-Atmos.*, 118, 7350–7369, <https://doi.org/10.1002/jgrd.50480>, 2013.
- Berger, M., Moreno, J., Johannessen, J. A., Levelt, P. F., and Hanssen, R. F.: ESA’s sentinel missions in support of Earth system science, *Remote Sens. Environ.*, 120, 84–90, <https://doi.org/10.1016/j.rse.2011.07.023>, 2012.
- Boesch, H., Baker, D., Connor, B., Crisp, D., and Miller, C.: Global Characterization of CO₂ Column Retrievals from Shortwave-Infrared Satellite Observations of the Orbiting Carbon Observatory-2 Mission, *Remote Sensing*, 3, 270–304, <https://doi.org/10.3390/rs3020270>, 2011.
- Bontemps, S., Herold, M., Kooistra, L., van Groenestijn, A., Hartley, A., Arino, O., Moreau, I., and Defourny, P.: Revisiting land cover observation to address the needs of the

- climate modeling community, *Biogeosciences*, 9, 2145–2157, <https://doi.org/10.5194/bg-9-2145-2012>, 2012.
- Boone, C. D., Nassar, R., Walker, K. A., Rochon, Y., McLeod, S. D., Rinsland, C. P., and Bernath, P. F.: Retrievals for the atmospheric chemistry experiment Fourier-transform spectrometer, *Appl. Opt.*, 44, 7218–7231, <https://doi.org/10.1364/AO.44.007218>, 2005.
- Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noël, S., Rozanov, V. V., Chance, K. V., and Goede, A. P. H.: SCIAMACHY: Mission Objectives and Measurement Modes, *J. Atmos. Sci.*, 56, 127–150, [https://doi.org/10.1175/1520-0469\(1999\)056<0127:SMOAMM>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<0127:SMOAMM>2.0.CO;2), 1999.
- Braswell, B. H., Sacks, W. J., Linder, E., and Schimel, D. S.: Estimating diurnal to annual ecosystem parameters by synthesis of a carbon flux model with eddy covariance net ecosystem exchange observations, *Glob. Change Biol.*, 11, 335–355, <https://doi.org/10.1111/j.1365-2486.2005.00897.x>, 2005.
- Buchwitz, M. and Reuter, M.: Merged SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT atmospheric column-average dry-air mole fraction of CO₂ (XCO₂), Technical Note, Version 1, available at: http://www.esa-ghg-cci.org/?q=webfm_send/319 (last access: 14 July 2017), 2016.
- Buchwitz, M., Rozanov, V. V., and Burrows, J. P.: A near-infrared optimized DOAS method for the fast global retrieval of atmospheric CH₄, CO, CO₂, H₂O, and N₂O total column amounts from SCIAMACHY Envisat-1 nadir radiances, *J. Geophys. Res.-Atmos.*, 105, 15231–15245, <https://doi.org/10.1029/2000JD900191>, 2000.
- Buchwitz, M., Reuter, M., Bovensmann, H., Pillai, D., Heymann, J., Schneising, O., Rozanov, V., Krings, T., Burrows, J. P., Boesch, H., Gerbig, C., Meijer, Y., and Lüscher, A.: Carbon Monitoring Satellite (CarbonSat): assessment of atmospheric CO₂ and CH₄ retrieval errors by error parameterization, *Atmos. Meas. Tech.*, 6, 3477–3500, <https://doi.org/10.5194/amt-6-3477-2013>, 2013.
- Buchwitz, M., Reuter, M., Schneising, O., Boesch, H., Guerlet, S., Dils, B., Aben, I., Armante, R., Bergamaschi, P., Blumenthal, T., Bovensmann, H., Brunner, D., Buchmann, B., Burrows, J., Butz, A., Chédin, A., Chevallier, F., Crevoisier, C., Deutscher, N., Frankenberg, C., Hase, F., Hasekamp, O., Heymann, J., Kaminski, T., Laeng, A., Lichtenberg, G., Mazière, M. D., Noël, S., Notholt, J., Orphal, J., Popp, C., Parker, R., Scholze, M., Sussmann, R., Stiller, G., Warneke, T., Zehner, C., Bril, A., Crisp, D., Griffith, D., Kuze, A., O'Dell, C., Oshchepkov, S., Sherlock, V., Suto, H., Wennberg, P., Wunch, D., Yokota, T., and Yoshida, Y.: The Greenhouse Gas Climate Change Initiative (GHG-CCI): Comparison and quality assessment of near-surface-sensitive satellite-derived CO₂ and CH₄ global data sets, *Remote Sens. Environ.*, 162, 344–362, <https://doi.org/10.1016/j.rse.2013.04.024>, 2015.
- Buchwitz, M., Dils, B., Boesch, H., Crevoisier, C., Detmers, D., Frankenberg, C., Hasekamp, O., Hewson, W., Laeng, A., Noël, S., Notholt, J., Parker, R., Reuter, M., and Schneising, O.: ESA Climate Change Initiative (CCI) Product Validation and Intercomparison Report (P VIR) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG) for data set Climate Research Data Package No. 3 (CRDP No. 3), Version 4.0, available at: http://www.esa-ghg-cci.org/?q=webfm_send/300 (last access: 14 July 2017), 2016.
- Buchwitz, M., Schneising, O., Reuter, M., Heymann, J., Krautwurst, S., Bovensmann, H., Burrows, J. P., Boesch, H., Parker, R. J., Somkuti, P., Detmers, R. G., Hasekamp, O. P., Aben, I., Butz, A., Frankenberg, C., and Turner, A. J.: Satellite-derived methane hotspot emission estimates using a fast data-driven method, *Atmos. Chem. Phys.*, 17, 5751–5774, <https://doi.org/10.5194/acp-17-5751-2017>, 2017.
- Burrows, J. P., Hödlzle, E., Goede, A. P. H., Visser, H., and Fricke, W.: SCIAMACHY – scanning imaging absorption spectrometer for atmospheric chartography, *Acta Astronautica*, 35, 445–451, [https://doi.org/10.1016/0094-5765\(94\)00278-T](https://doi.org/10.1016/0094-5765(94)00278-T), 1995.
- Butz, A., Hasekamp, O. P., Frankenberg, C., Vidot, J., and Aben, I.: CH₄ retrievals from space-based solar backscatter measurements: Performance evaluation against simulated aerosol and cirrus loaded scenes, *J. Geophys. Res.-Atmos.*, 115, D24302, <https://doi.org/10.1029/2010JD014514>, 2010.
- Butz, A., Guerlet, S., Hasekamp, O., Schepers, D., Galli, A., Aben, I., Frankenberg, C., Hartmann, J.-M., Tran, H., Kuze, A., Keppel-Aleks, G., Toon, G., Wunch, D., Wennberg, P., Deutscher, N., Griffith, D., Macatangay, R., Messerschmidt, J., Notholt, J., and Warneke, T.: Toward accurate CO₂ and CH₄ observations from GOSAT, *Geophys. Res. Lett.*, 38, L14812, <https://doi.org/10.1029/2011GL047888>, 2011.
- Butz, A., Galli, A., Hasekamp, O., Landgraf, J., Tol, P., and Aben, I.: TROPOMI aboard Sentinel-5 Precursor: Prospective performance of CH₄ retrievals for aerosol and cirrus loaded atmospheres, *Remote Sens. Environ.*, 120, 267–276, <https://doi.org/10.1016/j.rse.2011.05.030>, 2012.
- Cadule, P., Friedlingstein, P., Bopp, L., Sitch, S., Jones, C. D., Ciais, P., Piao, S. L., and Peylin, P.: Benchmarking coupled climate-carbon models against long-term atmospheric CO₂ measurements, *Global Biogeochem. Cy.*, 24, GB2016, <https://doi.org/10.1029/2009GB003556>, 2010.
- Ceccherini, G., Gobron, N., and Robustelli, M.: Harmonization of Fraction of Absorbed Photosynthetically Active Radiation (FAPAR) from Sea-ViewingWide Field-of-View Sensor (SeaWiFS) and Medium Resolution Imaging Spectrometer Instrument (MERIS), *Remote Sensing*, 5, 3357–3376, <https://doi.org/10.3390/rs5073357>, 2013.
- Chen, T., de Jeu, R. A. M., Liu, Y. Y., van der Werf, G. R., and Dolman, A. J.: Using satellite based soil moisture to quantify the water driven variability in NDVI: A case study over mainland Australia, *Remote Sens. Environ.*, 140, 330–338, 2014.
- Chevallier, F., Alexe, M., Bergamaschi, P., Brunner, D., Feng, L., Houweling, S., Kaminski, T., Knorr, W., van Leeuwen, T. T., Marshall, J., Palmer, P. I., Scholze, M., Sundström, A.-M., and Vossbeck, M.: ESA Climate Change Initiative (CCI) Climate Assessment Report (CAR) for Climate Research Data Package No. 4 (CRDP No. 4) of the Essential Climate Variable (ECV) Greenhouse Gases (GHG), Version 4, available at: http://www.esa-ghg-cci.org/?q=webfm_send/385, last access: 14 July 2017.
- Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R., Piao, S., and Thornton, P.: Carbon and Other Biogeochemical Cycles, book section 6, 465–570, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/CBO9781107415324.015>, 2013.

- Ciais, P., Dolman, A. J., Bombelli, A., Duren, R., Peregon, A., Rayner, P. J., Miller, C., Gobron, N., Kinderman, G., Marland, G., Gruber, N., Chevallier, F., Andres, R. J., Balsamo, G., Bopp, L., Bréon, F.-M., Broquet, G., Dargaville, R., Battin, T. J., Borges, A., Bovensmann, H., Buchwitz, M., Butler, J., Canadell, J. G., Cook, R. B., DeFries, R., Engelen, R., Gurney, K. R., Heinze, C., Heimann, M., Held, A., Henry, M., Law, B., Luysaert, S., Miller, J., Moriyama, T., Moulin, C., Myneni, R. B., Nussli, C., Obersteiner, M., Ojima, D., Pan, Y., Paris, J.-D., Piao, S. L., Poulter, B., Plummer, S., Quegan, S., Raymond, P., Reichstein, M., Rivier, L., Sabine, C., Schimel, D., Tarasova, O., Valentini, R., Wang, R., van der Werf, G., Wickland, D., Williams, M., and Zehner, C.: Current systematic carbon-cycle observations and the need for implementing a policy-relevant carbon observing system, *Biogeosciences*, 11, 3547–3602, <https://doi.org/10.5194/bg-11-3547-2014>, 2014.
- Ciais, P., Crisp, D., Denier van der Gon, H., Engelen, R., Heimann, M., Janssens-Maenhout, G., Rayner, P., and Scholze, M.: Towards a European Operational Observing System to Monitor Fossil CO₂ emissions, Final Report from the expert group, European Commission, B-1049 Brussels, Belgium, available at: http://www.copernicus.eu/sites/default/files/library/CO2_Report_22Oct2015.pdf (last access: 14 July 2017), 2015.
- Cihlar, J., Denning, S., Ahem, F., Arino, O., Belward, A., Bretherton, F., Cramer, W., Dedieu, G., Field, C., Francey, R., Gommes, R., Gosz, J., Hibbard, K., Igarashi, T., Kabat, P., Olson, D., Plummer, S., Rasool, I., Raupach, M., Scholes, R., Townshend, J., Valentini, R., and Wickland, D.: Initiative to quantify terrestrial carbon sources and sinks, *Eos, Transactions American Geophysical Union*, 83, 1–7, <https://doi.org/10.1029/2002EO000002>, 2002.
- Cogan, A. J., Boesch, H., Parker, R. J., Feng, L., Palmer, P. I., Blavier, J.-F. L., Deutscher, N. M., Macatangay, R., Notholt, J., Roehl, C., Warneke, T., and Wunch, D.: Atmospheric carbon dioxide retrieved from the Greenhouse gases Observing SATellite (GOSAT): Comparison with ground-based TCCON observations and GEOS-Chem model calculations, *J. Geophys. Res.-Atmos.*, 117, D21301, <https://doi.org/10.1029/2012JD018087>, d21301, 2012.
- Crevoisier, C., Chédin, A., Matsueda, H., Machida, T., Armante, R., and Scott, N. A.: First year of upper tropospheric integrated content of CO₂ from IASI hyperspectral infrared observations, *Atmos. Chem. Phys.*, 9, 4797–4810, <https://doi.org/10.5194/acp-9-4797-2009>, 2009a.
- Crevoisier, C., Nobileau, D., Fiore, A. M., Armante, R., Chédin, A., and Scott, N. A.: Tropospheric methane in the tropics – first year from IASI hyperspectral infrared observations, *Atmos. Chem. Phys.*, 9, 6337–6350, <https://doi.org/10.5194/acp-9-6337-2009>, 2009b.
- Crisp, D., Atlas, R., Breon, F.-M., Brown, L., Burrows, J., Ciais, P., Connor, B., Doney, S., Fung, I., Jacob, D., Miller, C., O'Brien, D., Pawson, S., Randerson, J., Rayner, P., Salawitch, R., Sander, S., Sen, B., Stephens, G., Tans, P., Toon, G., Wennberg, P., Wofsy, S., Yung, Y., Kuang, Z., Chudasama, B., Sprague, G., Weiss, B., Pollock, R., Kenyon, D., and Schroll, S.: The Orbiting Carbon Observatory (OCO) mission, *Adv. Space Res.*, 34, 700–709, <https://doi.org/10.1016/j.asr.2003.08.062>, 2004.
- Crisp, D., Fisher, B. M., O'Dell, C., Frankenberg, C., Basilio, R., Bösch, H., Brown, L. R., Castano, R., Connor, B., Deutscher, N. M., Eldering, A., Griffith, D., Gunson, M., Kuze, A., Mandrake, L., McDuffie, J., Messerschmidt, J., Miller, C. E., Morino, I., Natraj, V., Notholt, J., O'Brien, D. M., Oyafuso, F., Polonsky, I., Robinson, J., Salawitch, R., Sherlock, V., Smyth, M., Suto, H., Taylor, T. E., Thompson, D. R., Wennberg, P. O., Wunch, D., and Yung, Y. L.: The ACOS CO₂ retrieval algorithm – Part II: Global XCO₂ data characterization, *Atmos. Meas. Tech.*, 5, 687–707, <https://doi.org/10.5194/amt-5-687-2012>, 2012.
- Daley, R.: *Atmospheric data analysis*, Cambridge University Press, Cambridge, UK, 1991.
- De Jeu, R. and Dorigo, W.: On the importance of satellite observed soil moisture, *International Journal of Applied Earth Observation and Geoinformation*, 45, Part B, 107–109, <https://doi.org/10.1016/j.jag.2015.10.007>, 2016.
- Deering, D., Rouse, J., Haas, R., and Schell, J.: Measuring forage production of grazing units from Landsat MSS data, *Proc. 10th Int. Symp. Remote Sensing Environ.*, University of Michigan, Ann Arbor, USA, 1975.
- Dils, B., Buchwitz, M., Reuter, M., Schneing, O., Boesch, H., Parker, R., Guerlet, S., Aben, I., Blumenstock, T., Burrows, J. P., Butz, A., Deutscher, N. M., Frankenberg, C., Hase, F., Hasekamp, O. P., Heymann, J., De Mazière, M., Notholt, J., Sussmann, R., Warneke, T., Griffith, D., Sherlock, V., and Wunch, D.: The Greenhouse Gas Climate Change Initiative (GHG-CCI): comparative validation of GHG-CCI SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT CO₂ and CH₄ retrieval algorithm products with measurements from the TCCON, *Atmos. Meas. Tech.*, 7, 1723–1744, <https://doi.org/10.5194/amt-7-1723-2014>, 2014.
- Disney, M., Muller, J.-P., Kharbouche, S., Kaminski, T., Vossbeck, M., Lewis, P., and Pinty, B.: A New Global fAPAR and LAI Dataset Derived from Optimal Albedo Estimates: Comparison with MODIS Products, *Remote Sensing*, 8, 27, <https://doi.org/10.3390/rs8040275>, 2016.
- D'Odorico, P., Gonsamo, A., Pinty, B., Gobron, N., Coops, N., Mendez, E., and Schaepman, M. E.: Intercomparison of fraction of absorbed photosynthetically active radiation products derived from satellite data over Europe, *Remote Sens. Environ.*, 142, 141–154, <https://doi.org/10.1016/j.rse.2013.12.005>, 2014.
- Dorigo, W., Zurita-Milla, R., de Wit, A., Brazile, J., Singh, R., and Schaepman, M.: A review on reflective remote sensing and data assimilation techniques for enhanced agroecosystem modeling, *International Journal of Applied Earth Observation and Geoinformation*, 9, 165–193, <https://doi.org/10.1016/j.jag.2006.05.003>, 2007.
- Dorigo, W., De Jeu, R., Chung, D., Parinussa, R., Liu, Y., Wagner, W., and Fernandez-Prieto, D.: Evaluating global trends (1988–2010) in homogenized remotely sensed surface soil moisture, *Geophys. Res. Lett.*, 39, L18405, <https://doi.org/10.1029/2012gl052988>, 2012.
- Dorigo, W., Xaver, A., Vreugdenhil, M., Gruber, A., Hegyiová, A., Sanchis-Dufau, A., Wagner, W., and Drusch, M.: Global automated quality control of in-situ soil moisture data from the International Soil Moisture Network, *Vadose Zone J.*, 12, <https://doi.org/10.2136/vzj2012.0097>, 2013.
- Dorigo, W., Wagner, W., Albergel, C., Albrecht, F., Balsamo, G., Brocca, L., Chung, D., Ertl, M., Forkel, M., Gruber, A., Haas, E., Hamer, P., Hirschi, M., Ikonen, J., Jeu, R., Kidd, R., Lahoz, W., Liu, Y., Miralles, D., Mistelbauer, T., Nicolai-Shaw, N., Par-

- inussa, R., Pratola, C., Reimer, C., Schalie, R., Seneviratne, S., Smolander, T., and Lecomte, P.: ESA CCI Soil Moisture for improved Earth system understanding: state-of-the art and future directions, *Remote Sens. Environ.*, under review, 2016.
- Dorigo, W. A., Scipal, K., Parinussa, R. M., Liu, Y. Y., Wagner, W., de Jeu, R. A. M., and Naeimi, V.: Error characterisation of global active and passive microwave soil moisture datasets, *Hydrol. Earth Syst. Sci.*, 14, 2605–2616, <https://doi.org/10.5194/hess-14-2605-2010>, 2010.
- Dorigo, W. A., Wagner, W., Hohensinn, R., Hahn, S., Paulik, C., Xaver, A., Gruber, A., Drusch, M., Mecklenburg, S., van Oevelen, P., Robock, A., and Jackson, T.: The International Soil Moisture Network: a data hosting facility for global in situ soil moisture measurements, *Hydrol. Earth Syst. Sci.*, 15, 1675–1698, <https://doi.org/10.5194/hess-15-1675-2011>, 2011.
- Dorigo, W. A., Gruber, A., De Jeu, R. A. M., Wagner, W., Stacke, T., Loew, A., Albergel, C., Brocca, L., Chung, D., Parinussa, R. M., and Kidd, R.: Evaluation of the ESA CCI soil moisture product using ground-based observations, *Remote Sens. Environ.*, 162, 380–395, <https://doi.org/10.1016/j.rse.2014.07.023>, 2015.
- Draper, C., Reichle, R., de Jeu, R., Naeimi, V., Parinussa, R., and Wagner, W.: Estimating root mean square errors in remotely sensed soil moisture over continental scale domains, *Remote Sens. Environ.*, 137, 288–298, <https://doi.org/10.1016/j.rse.2013.06.013>, 2013.
- Drusch, M., Moreno, J., Bello, U. D., Franco, R., Goulas, Y., Huth, A., Kraft, S., Middleton, E. M., Miglietta, F., Mohammed, G., Nedbal, L., Rascher, U., Schüttmeyer, D., and Verhoef, W.: The FLuorescence EXplorer Mission Concept – ESA's Earth Explorer 8, *IEEE T. Geosci. Remote*, 55, 1273–1284, <https://doi.org/10.1109/TGRS.2016.2621820>, 2017.
- Entekhabi, D., Njoku, E. G., O'Neill, P. E., Kellogg, K. H., Crow, W. T., Edelstein, W. N., Entin, J. K., Goodman, S. D., Jackson, T. J., Johnson, J., Kimball, J., Piepmeyer, J. R., Koster, R. D., Martin, N., McDonald, K. C., Moghaddam, M., Moran, S., Reichle, R., Shi, J. C., Spencer, M. W., Thurman, S. W., Tsang, L., and Van Zyl, J.: The soil moisture active passive (SMAP) mission, *Proceedings of the IEEE*, 98, 704–716, 2010.
- Enting, I. G.: Inverse Problems in Atmospheric Constituent Transport, Cambridge University Press, Cambridge, 2002.
- European Space Agency: Report for Mission Selection: Biomass, Science authors: Quegan, S., Le Toan T., Chave, J., Dall, J., Perrera, A., Papathanassiou, K., Rocca, F., Saatchi, S., Scipal, K., Shugart, H., Ulander, L., and Williams, M., ESA SP 1324/1, European Space Agency, Noordwijk, the Netherlands, available at: http://esamultimedia.esa.int/docs/EarthObservation/SP1324-1_BIOMASSr.pdf (last access: 14 July 2017), 2012.
- Foley, A. M., Dalmonech, D., Friend, A. D., Aires, F., Archibald, A. T., Bartlein, P., Bopp, L., Chappellaz, J., Cox, P., Edwards, N. R., Feulner, G., Friedlingstein, P., Harrison, S. P., Hopcroft, P. O., Jones, C. D., Kolassa, J., Levine, J. G., Prentice, I. C., Pyle, J., Vázquez Riveiros, N., Wolff, E. W., and Zaehle, S.: Evaluation of biospheric components in Earth system models using modern and palaeo-observations: the state-of-the-art, *Biogeosciences*, 10, 8305–8328, <https://doi.org/10.5194/bg-10-8305-2013>, 2013.
- Forkel, M., Carvalhais, N., Schaphoff, S., v. Bloh, W., Migliavacca, M., Thurner, M., and Thonicke, K.: Identifying environmental controls on vegetation greenness phenology through model-data integration, *Biogeosciences*, 11, 7025–7050, <https://doi.org/10.5194/bg-11-7025-2014>, 2014.
- Foucher, P. Y., Chédin, A., Dufour, G., Capelle, V., Boone, C. D., and Bernath, P.: Technical Note: Feasibility of CO₂ profile retrieval from limb viewing solar occultation made by the ACE-FTS instrument, *Atmos. Chem. Phys.*, 9, 2873–2890, <https://doi.org/10.5194/acp-9-2873-2009>, 2009.
- Fox, A., Williams, M., Richardson, A. D., Cameron, D., Gove, J. H., Quaife, T., Ricciuto, D., Reichstein, M., Tomelleri, E., Trudinger, C. M., and Wijk, M. T. V.: The {REFLEX} project: Comparing different algorithms and implementations for the inversion of a terrestrial ecosystem model against eddy covariance data, *Agr. Forest Meteorol.*, 149, 1597–1615, <https://doi.org/10.1016/j.agrformet.2009.05.002>, 2009.
- Francey, R. J., Rayner, P. J., and Allison, C. E.: Global Biogeochemical Cycles in the Climate System, chap. Constraining the global carbon budget from global to regional scales – the measurement challenge, Academic Press, San Diego, USA, 245–252, 2001.
- Frankenberg, C., Aben, I., Bergamaschi, P., Dlugokencky, E. J., van Hees, R., Houweling, S., van der Meer, P., Snel, R., and Tol, P.: Global column-averaged methane mixing ratios from 2003 to 2009 as derived from SCIAMACHY: Trends and variability, *J. Geophys. Res.-Atmos.*, 116, D04302, <https://doi.org/10.1029/2010JD014849>, 2011a.
- Frankenberg, C., Butz, A., and Toon, G. C.: Disentangling chlorophyll fluorescence from atmospheric scattering effects in O2A-band spectra of reflected sun-light, *Geophys. Res. Lett.*, 38, L03801, <https://doi.org/10.1029/2010GL045896>, 2011b.
- Frankenberg, C., Fisher, J. B., Worden, J., Badgley, G., Saatchi, S. S., Lee, J.-E., Toon, G. C., Butz, A., Jung, M., Kuze, A., and Yokota, T.: New global observations of the terrestrial carbon cycle from GOSAT: Patterns of plant fluorescence with gross primary productivity, *Geophys. Res. Lett.*, 38, L17706, <https://doi.org/10.1029/2011GL048738>, 2011c.
- Frankenberg, C., O'Dell, C., Berry, J., Guanter, L., Joiner, J., Köhler, P., Pollock, R., and Taylor, T. E.: Prospects for chlorophyll fluorescence remote sensing from the Orbiting Carbon Observatory-2, *Remote Sens. Environ.*, 147, 1–12, 2014.
- GCOS: Global Climate Observing System: Systematic Observation Requirements for Satellite-based Products for Climate, GCOS – 154, available at: <https://www.wmo.int/pages/prog/geos/Publications/gcos-154.pdf> (last access: 14 July 2017), 2011.
- Giglio, L., Randerson, J. T., and van der Werf, G. R.: Analysis of daily, monthly, and annual burned area using the fourth-generation global fire emissions database (GFED4), *J. Geophys. Res.-Biogeo.*, 118, 317–328, <https://doi.org/10.1002/jgrg.20042>, 2013.
- Global Carbon Project: Science Framework and Implementation. Earth System Science Partnership (IGBP, IHDP, WCRP, DIVERSITAS), Report No. 1; Global Carbon Project Report No. 1, 69 pp., Canberra, 2003.
- Gobron, N. and Verstraete, M. M.: FAPAR, fraction of absorbed photosynthetically active radiation – Assessment of the status of the development of the standards for the terrestrial essential climate variables, Version 8, GTOS Secretariat, FAO, Italy, available at: <http://www.fao.org/gtos/doc/ECVs/T10/T10.pdf> (last access: 14 July 2017), 2009.

- Gobron, N., Pinty, B., Aussedat, O., Chen, J. M., Cohen, W. B., Fensholt, R., Gond, V., Huemmrich, K. F., Lavergne, T., Mélin, F., Privette, J. L., Sandholt, I., Taberner, M., Turner, D. P., Verstraete, M. M., and Widlowski, J.-L.: Evaluation of fraction of absorbed photosynthetically active radiation products for different canopy radiation transfer regimes: Methodology and results using Joint Research Center products derived from SeaWiFS against ground-based estimations, *J. Geophys. Res.-Atmos.*, 111, D13110, <https://doi.org/10.1029/2005JD006511>, 2006.
- Gobron, N., Pinty, B., Aussedat, O., Taberner, M., Faber, O., Melin, F., Lavergne, T., Robustelli, M., and Snoejj, P.: Uncertainty estimates for the FAPAR operational products derived from MERIS: Impact of top-of-atmosphere radiance uncertainties and validation with field data, *Remote Sens. Environ.*, 112, 1871–1883, <https://doi.org/10.1016/j.rse.2007.09.011>, remote Sensing Data Assimilation Special Issue, 2008.
- Goel, N. S. and Qin, W.: Influences of canopy architecture on relationships between various vegetation indices and LAI and Fpar: A computer simulation, *Remote Sensing Reviews*, 10, 309–347, <https://doi.org/10.1080/02757259409532252>, 1994.
- Gruber, A., Dorigo, W., Zwieback, S., Xaver, A., and Wagner, W.: Characterizing coarse-scale representativeness of in-situ soil moisture measurements from the International Soil Moisture Network, *Vadose Zone J.*, 12, 16 pp., <https://doi.org/10.2136/vzj2012.0170>, 2013.
- Gruber, A., Su, C., Zwieback, S., Crow, W. T., Wagner, W., and Dorigo, W.: Recent advances in (soil moisture) triple collocation analysis, *International Journal of Applied Earth Observation and Geoinformation Part B*, 45, 200–211, 2016a.
- Gruber, A., Su, C. H., Crow, W. T., Zwieback, S., Dorigo, W. A., and Wagner, W.: Estimating error cross-correlations in soil moisture data sets using extended collocation analysis, *J. Geophys. Res.-Atmos.*, 121, 1208–1219, <https://doi.org/10.1002/2015JD024027>, 2016b.
- Guanter, L., Frankenberg, C., Dudhia, A., Lewis, P. E., Gómez-Dans, J., Kuze, A., Suto, H., and Grainger, R. G.: Retrieval and global assessment of terrestrial chlorophyll fluorescence from GOSAT space measurements, *Remote Sens. Environ.*, 121, 236–251, 2012.
- Guanter, L., Zhang, Y., Jung, M., Joiner, J., Voigt, M., Berry, J. A., Frankenberg, C., Huete, A. R., Zarco-Tejada, P., Lee, J.-E., Moran, M. S., Ponce-Campos, G., Beer, C., Camps-Valls, G., Buchmann, N., Gianelle, D., Klumpp, K., Cescatti, A., Baker, J. M., and Griffis, T. J.: Global and time-resolved monitoring of crop photosynthesis with chlorophyll fluorescence, *P. Natl. Acad. Sci. USA*, 111, E1327–E1333, 2014.
- Guanter, L., Aben, I., Tol, P., Krijger, J. M., Hollstein, A., Köhler, P., Damm, A., Joiner, J., Frankenberg, C., and Landgraf, J.: Potential of the TROPOspheric Monitoring Instrument (TROPOMI) onboard the Sentinel-5 Precursor for the monitoring of terrestrial chlorophyll fluorescence, *Atmos. Meas. Tech.*, 8, 1337–1352, <https://doi.org/10.5194/amt-8-1337-2015>, 2015.
- Heymann, J., Reuter, M., Hilker, M., Buchwitz, M., Schneising, O., Bovensmann, H., Burrows, J. P., Kuze, A., Suto, H., Deutscher, N. M., Dubey, M. K., Griffith, D. W. T., Hase, F., Kawakami, S., Kivi, R., Morino, I., Petri, C., Roehl, C., Schneider, M., Sherlock, V., Sussmann, R., Velazco, V. A., Warneke, T., and Wunch, D.: Consistent satellite XCO₂ retrievals from SCIAMACHY and GOSAT using the BESD algorithm, *Atmos. Meas. Tech.*, 8, 2961–2980, <https://doi.org/10.5194/amt-8-2961-2015>, 2015.
- Hollmann, R., Merchant, C. J., Saunders, R., Downy, C., Buchwitz, M., Cazenave, A., Chuvieco, E., Defourny, P., de Leeuw, G., Forsberg, R., Holzer-Popp, T., Paul, F., Sandven, S., Sathyendranath, S., van Roozendael, M., and Wagner, W.: The ESA Climate Change Initiative: Satellite Data Records for Essential Climate Variables, *B. Am. Meteorol. Soc.*, 94, 1541–1552, <https://doi.org/10.1175/BAMS-D-11-00254.1>, 2013.
- Houweling, S., Baker, D., Basu, S., Boesch, H., Butz, A., Chevallier, F., Deng, F., Dlugokencky, E. J., Feng, L., Ganeshin, A., Hasekamp, O., Jones, D., Maksyutov, S., Marshall, J., Oda, T., O'Dell, C. W., Oshchepkov, S., Palmer, P. I., Peylin, P., Poussi, Z., Reum, F., Takagi, H., Yoshida, Y., and Zhuravlev, R.: An intercomparison of inverse models for estimating sources and sinks of CO₂ using GOSAT measurements, *J. Geophys. Res.-Atmos.*, 120, 5253–5266, <https://doi.org/10.1002/2014JD022962>, 2014JD022962, 2015.
- Huete, A.: A soil-adjusted vegetation index (SAVI), *Remote Sens. Environ.*, 25, 295–309, [https://doi.org/10.1016/0034-4257\(88\)90106-X](https://doi.org/10.1016/0034-4257(88)90106-X), 1988.
- Jackson, T.: Measuring surface soil moisture using passive microwave remote sensing, *Hydrol. Process.*, 7, 139–152, 1993.
- Joiner, J., Yoshida, Y., Vasilkov, A. P., Yoshida, Y., Corp, L. A., and Middleton, E. M.: First observations of global and seasonal terrestrial chlorophyll fluorescence from space, *Biogeosciences*, 8, 637–651, <https://doi.org/10.5194/bg-8-637-2011>, 2011.
- Joiner, J., Yoshida, Y., Vasilkov, A. P., Middleton, E. M., Campbell, P. K. E., Yoshida, Y., Kuze, A., and Corp, L. A.: Filling-in of near-infrared solar lines by terrestrial fluorescence and other geophysical effects: simulations and space-based observations from SCIAMACHY and GOSAT, *Atmos. Meas. Tech.*, 5, 809–829, <https://doi.org/10.5194/amt-5-809-2012>, 2012.
- Joiner, J., Guanter, L., Lindstrot, R., Voigt, M., Vasilkov, A. P., Middleton, E. M., Huemmrich, K. F., Yoshida, Y., and Frankenberg, C.: Global monitoring of terrestrial chlorophyll fluorescence from moderate-spectral-resolution near-infrared satellite measurements: methodology, simulations, and application to GOME-2, *Atmos. Meas. Tech.*, 6, 2803–2823, <https://doi.org/10.5194/amt-6-2803-2013>, 2013.
- Joiner, J., Yoshida, Y., Guanter, L., and Middleton, E. M.: New methods for the retrieval of chlorophyll red fluorescence from hyperspectral satellite instruments: simulations and application to GOME-2 and SCIAMACHY, *Atmos. Meas. Tech.*, 9, 3939–3967, <https://doi.org/10.5194/amt-9-3939-2016>, 2016.
- Kaminski, T. and Mathieu, P.-P.: Reviews and syntheses: Flying the satellite into your model: on the role of observation operators in constraining models of the Earth system and the carbon cycle, *Biogeosciences*, 14, 2343–2357, <https://doi.org/10.5194/bg-14-2343-2017>, 2017.
- Kaminski, T., Knorr, W., Rayner, P., and Heimann, M.: Assimilating Atmospheric data into a Terrestrial Biosphere Model: A case study of the seasonal cycle, *Global Biogeochem. Cy.*, 16, 14–1–14–16, 2002.
- Kaminski, T., Knorr, W., Scholze, M., Gobron, N., Pinty, B., Giering, R., and Mathieu, P.-P.: Consistent assimilation of MERIS FAPAR and atmospheric CO₂ into a terrestrial vegetation model and interactive mission benefit analysis, *Biogeosciences*, 9, 3173–3184, <https://doi.org/10.5194/bg-9-3173-2012>, 2012.

- Kaminski, T., Knorr, W., Schürmann, G., Scholze, M., Rayner, P. J., Zaehle, S., Blessing, S., Dorigo, W., Gayler, V., Giering, R., Gobron, N., Grant, J. P., Heimann, M., Hooker-Stroud, A., Houweling, S., Kato, T., Kattge, J., Kelley, D., Kemp, S., Koffi, E. N., Köstler, C., Mathieu, P.-P., Pinty, B., Reick, C. H., Rödenbeck, C., Schnur, R., Scipal, K., Sebald, C., Stacke, T., van Scheltinga, A. T., Vossbeck, M., Widmann, H., and Ziehn, T.: The BETHY/JSBACH Carbon Cycle Data Assimilation System: experiences and challenges, *J. Geophys. Res.-Biogeosci.*, 118, 1414–1426, <https://doi.org/10.1002/jgrg.20118>, 2013.
- Kaminski, T., Scholze, M., Vossbeck, M., Knorr, W., Buchwitz, M., and Reuter, M.: Constraining a terrestrial biosphere model with remotely sensed atmospheric carbon dioxide, *Remote Sens. Environ.*, submitted, 2016.
- Kaminski, T., Pinty, B., Voßbeck, M., Lopatka, M., Gobron, N., and Robustelli, M.: Consistent retrieval of land surface radiation products from EO, including traceable uncertainty estimates, *Biogeosciences*, 14, 2527–2541, <https://doi.org/10.5194/bg-14-2527-2017>, 2017.
- Kato, T., Knorr, W., Scholze, M., Veenendaal, E., Kaminski, T., Kattge, J., and Gobron, N.: Simultaneous assimilation of satellite and eddy covariance data for improving terrestrial water and carbon simulations at a semi-arid woodland site in Botswana, *Biogeosciences*, 10, 789–802, <https://doi.org/10.5194/bg-10-789-2013>, 2013.
- Kaufman, Y. J. and Tanre, D.: Atmospherically resistant vegetation index (ARVI) for EOS-MODIS, *IEEE T. Geosci. Remote*, 30, 261–270, <https://doi.org/10.1109/36.134076>, 1992.
- Keeling, C. D.: The concentration and isotopic abundance of carbon dioxide in rural and marine air, *Geochim. Cosmochim. Ac.*, 24, 277–298, 1961.
- Keenan, T. F., Davidson, E., Moffat, A. M., Munger, W., and Richardson, A. D.: Using model-data fusion to interpret past trends, and quantify uncertainties in future projections, of terrestrial ecosystem carbon cycling, *Glob. Change Biol.*, 18, 2555–2569, <https://doi.org/10.1111/j.1365-2486.2012.02684.x>, 2012.
- Kelley, D. I., Prentice, I. C., Harrison, S. P., Wang, H., Simard, M., Fisher, J. B., and Willis, K. O.: A comprehensive benchmarking system for evaluating global vegetation models, *Biogeosciences*, 10, 3313–3340, <https://doi.org/10.5194/bg-10-3313-2013>, 2013.
- Kerr, Y. H., Waldteufel, P., Wigneron, J. P., Delwart, S., Cabot, F., Boutin, J., Escorihuela, M. J., Font, J., Reul, N., Gruhier, C., Juglea, S. E., Drinkwater, M. R., Hahne, A., Martin-Neira, M., and Mecklenburg, S.: The SMOS Mission: New Tool for Monitoring Key Elements of the Global Water Cycle, *Proceedings of the IEEE*, 98, 666–687, <https://doi.org/10.1109/JPROC.2010.2043032>, 2010.
- Kerr, Y. H., Waldteufel, P., Richaume, P., Wigneron, J. P., Ferrazzoli, P., Mahmoodi, A., Bitar, A. A., Cabot, F., Gruhier, C., Juglea, S. E., Leroux, D., Mialon, A., and Delwart, S.: The SMOS Soil Moisture Retrieval Algorithm, *IEEE T. Geosci. Remote*, 50, 1384–1403, <https://doi.org/10.1109/TGRS.2012.2184548>, 2012.
- Knorr, W. and Kattge, J.: Inversion of terrestrial biosphere model parameter values against eddy covariance measurements using Monte Carlo sampling, *Glob. Change Biol.*, 11, 1333–1351, 2005.
- Knorr, W., Kaminski, T., Scholze, M., Gobron, N., Pinty, B., Giering, R., and Mathieu, P.-P.: Carbon cycle data assimilation with a generic phenology model, *J. Geophys. Res.-Biogeosci.*, 115, G04017, <https://doi.org/10.1029/2009JG001119>, 2010.
- Köhler, P., Guanter, L., and Frankenberg, C.: Simplified Physically Based Retrieval of Sun-Induced Chlorophyll Fluorescence From GOSAT Data, *IEEE Geosci. Remote S.*, 12, 1446–1450, <https://doi.org/10.1109/LGRS.2015.2407051>, 2015a.
- Köhler, P., Guanter, L., and Joiner, J.: A linear method for the retrieval of sun-induced chlorophyll fluorescence from GOME-2 and SCIAMACHY data, *Atmos. Meas. Tech.*, 8, 2589–2608, <https://doi.org/10.5194/amt-8-2589-2015>, 2015b.
- Konings, A. G., Piles, M., Rötzer, K., McColl, K. A., Chan, S. K., and Entekhabi, D.: Vegetation optical depth and scattering albedo retrieval using time series of dual-polarized L-band radiometer observations, *Remote Sens. Environ.*, 172, 178–189, <https://doi.org/10.1016/j.rse.2015.11.009>, 2016.
- Kulawik, S., Wunch, D., O'Dell, C., Frankenberg, C., Reuter, M., Oda, T., Chevallier, F., Sherlock, V., Buchwitz, M., Osterman, G., Miller, C. E., Wennberg, P. O., Griffith, D., Morino, I., Dubey, M. K., Deutscher, N. M., Notholt, J., Hase, F., Warneke, T., Sussmann, R., Robinson, J., Strong, K., Schneider, M., De Mazière, M., Shiomi, K., Feist, D. G., Iraci, L. T., and Wolf, J.: Consistent evaluation of ACOS-GOSAT, BESD-SCIAMACHY, CarbonTracker, and MACC through comparisons to TCCON, *Atmos. Meas. Tech.*, 9, 683–709, <https://doi.org/10.5194/amt-9-683-2016>, 2016.
- Kuppel, S., Peylin, P., Chevallier, F., Bacour, C., Maignan, F., and Richardson, A. D.: Constraining a global ecosystem model with multi-site eddy-covariance data, *Biogeosciences*, 9, 3757–3776, <https://doi.org/10.5194/bg-9-3757-2012>, 2012.
- Kuze, A., Suto, H., Nakajima, M., and Hamazaki, T.: Thermal and near infrared sensor for carbon observation Fourier-transform spectrometer on the Greenhouse Gases Observing Satellite for greenhouse gases monitoring, *Appl. Opt.*, 48, 6716–6733, <https://doi.org/10.1364/AO.48.006716>, 2009.
- Kuze, A., Taylor, T. E., Kataoka, F., Bruegge, C. J., Crisp, D., Harada, M., Helmlinger, M., Inoue, M., Kawakami, S., Kikuchi, N., Mitomi, Y., Murooka, J., Naitoh, M., O'Brien, D. M., O'Dell, C. W., Ohya, H., Pollock, H., Schwandner, F. M., Shiomi, K., Suto, H., Takeda, T., Tanaka, T., Urabe, T., Yokota, T., and Yoshida, Y.: Long-Term Vicarious Calibration of GOSAT Short-Wave Sensors: Techniques for Error Reduction and New Estimates of Radiometric Degradation Factors, *IEEE T. Geosci. Remote*, 52, 3991–4004, <https://doi.org/10.1109/TGRS.2013.2278696>, 2014.
- Laeng, A., Plieninger, J., von Clarmann, T., Grabowski, U., Stiller, G., Eckert, E., Glatthor, N., Haenel, F., Kellmann, S., Kiefer, M., Linden, A., Lossow, S., Deaver, L., Engel, A., Hervig, M., Levin, I., McHugh, M., Noël, S., Toon, G., and Walker, K.: Validation of MIPAS IMK/IAA methane profiles, *Atmos. Meas. Tech.*, 8, 5251–5261, <https://doi.org/10.5194/amt-8-5251-2015>, 2015.
- Lannoy, G. J. M. D. and Reichle, R. H.: Global Assimilation of Multiangle and Multipolarization SMOS Brightness Temperature Observations into the GEOS-5 Catchment Land Surface Model for Soil Moisture Estimation, *J. Hydrometeorol.*, 17, 669–691, <https://doi.org/10.1175/JHM-D-15-0037.1>, 2016.
- Lasslop, G., Reichstein, M., Kattge, J., and Papale, D.: Influences of observation errors in eddy flux data on inverse model parameter estimation, *Biogeosciences*, 5, 1311–1324, <https://doi.org/10.5194/bg-5-1311-2008>, 2008.

- Le Quéré, C., Moriarty, R., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I., Friedlingstein, P., Peters, G. P., Andres, R. J., Boden, T. A., Houghton, R. A., House, J. I., Keeling, R. F., Tans, P., Arneth, A., Bakker, D. C. E., Barbero, L., Bopp, L., Chang, J., Chevallier, F., Chini, L. P., Ciais, P., Fader, M., Feely, R. A., Gkrizalis, T., Harris, I., Hauck, J., Ilyina, T., Jain, A. K., Kato, E., Kitidis, V., Klein Goldewijk, K., Koven, C., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lima, I. D., Metzl, N., Millero, F., Munro, D. R., Murata, A., Nabel, J. E. M. S., Nakaoaka, S., Nojiri, Y., O'Brien, K., Olsen, A., Ono, T., Pérez, F. F., Pfeil, B., Pierrot, D., Poulter, B., Rehder, G., Rödenbeck, C., Saito, S., Schuster, U., Schwinger, J., Séférian, R., Steinhoff, T., Stocker, B. D., Sutton, A. J., Takahashi, T., Tilbrook, B., van der Laan-Luijkx, I. T., van der Werf, G. R., van Heuven, S., Vandemark, D., Viovy, N., Wiltshire, A., Zaehle, S., and Zeng, N.: Global Carbon Budget 2015, *Earth Syst. Sci. Data*, 7, 349–396, <https://doi.org/10.5194/essd-7-349-2015>, 2015.
- Lee, J.-E., Frankenberg, C., van der Tol, C., Berry, J. A., Gartner, L., Boyce, C. K., Fisher, J. B., Morrow, E., Worden, J. R., Asefi, S., Badgley, G., and Saatchi, S.: Forest productivity and water stress in Amazonia: observations from GOSAT chlorophyll fluorescence, *P. Roy. Soc. B-Biol. Sci.*, 280, 20130171, <https://doi.org/10.1098/rspb.2013.0171>, 2013.
- Lefsky, M. A.: A global forest canopy height map from the Moderate Resolution Imaging Spectroradiometer and the Geoscience Laser Altimeter System, *Geophys. Res. Lett.*, 37, L15401, <https://doi.org/10.1029/2010GL043622>, 2010.
- Leprieur, C., Verstraete, M. M., and Pinty, B.: Evaluation of the performance of various vegetation indices to retrieve vegetation cover from AVHRR data, *Remote Sensing Reviews*, 10, 265–284, <https://doi.org/10.1080/02757259409532250>, 1994.
- Li, Z.-L., Tang, B.-H., Wu, H., Ren, H., Yan, G., Wan, Z., Trigo, I. F., and Sobrino, J. A.: Satellite-derived land surface temperature: Current status and perspectives, *Remote Sens. Environ.*, 131, 14–37, <https://doi.org/10.1016/j.rse.2012.12.008>, 2013.
- Liu, Q., Liang, S., Xiao, Z., and Fang, H.: Retrieval of leaf area index using temporal, spectral, and angular information from multiple satellite data, *Remote Sens. Environ.*, 145, 25–37, <https://doi.org/10.1016/j.rse.2014.01.021>, 2014.
- Liu, Y. Y., Parinussa, R. M., Dorigo, W. A., De Jeu, R. A. M., Wagner, W., van Dijk, A. I. J. M., McCabe, M. F., and Evans, J. P.: Developing an improved soil moisture dataset by blending passive and active microwave satellite-based retrievals, *Hydrol. Earth Syst. Sci.*, 15, 425–436, <https://doi.org/10.5194/hess-15-425-2011>, 2011.
- Liu, Y., Dorigo, W., Parinussa, R., De Jeu, R., Wagner, W., McCabe, M., Evans, J., and Van Dijk, A. I. J. M.: Trend-preserving blending of passive and active microwave soil moisture retrievals, *Remote Sens. Environ.*, 123, 280–297, <https://doi.org/10.1016/j.rse.2012.03.014>, 2012.
- Liu, Y. Y., van Dijk, A. I. J. M., de Jeu, R. A. M., Canadell, J. G., McCabe, M. F., Evans, J. P., and Wang, G.: Recent reversal in loss of global terrestrial biomass, *Nature Climate Change*, 5, 470–474, <https://doi.org/10.1038/nclimate2581>, 2015.
- Luke, C. M.: Modelling aspects of land-atmosphere interaction: Thermal instability in peatland soils and land parameter through data assimilation, PhD thesis, University of Exeter, UK, 2011.
- Luo, Y. Q., Randerson, J. T., Abramowitz, G., Bacour, C., Blyth, E., Carvalhais, N., Ciais, P., Dalmonech, D., Fisher, J. B., Fisher, R., Friedlingstein, P., Hibbard, K., Hoffman, F., Huntzinger, D., Jones, C. D., Koven, C., Lawrence, D., Li, D. J., Mahecha, M., Niu, S. L., Norby, R., Piao, S. L., Qi, X., Peylin, P., Prentice, I. C., Riley, W., Reichstein, M., Schwalm, C., Wang, Y. P., Xia, J. Y., Zaehle, S., and Zhou, X. H.: A framework for benchmarking land models, *Biogeosciences*, 9, 3857–3874, <https://doi.org/10.5194/bg-9-3857-2012>, 2012.
- MacBean, N., Maignan, F., Peylin, P., Bacour, C., Bréon, F.-M., and Ciais, P.: Using satellite data to improve the leaf phenology of a global terrestrial biosphere model, *Biogeosciences*, 12, 7185–7208, <https://doi.org/10.5194/bg-12-7185-2015>, 2015.
- MacBean, N., Peylin, P., Chevallier, F., Scholze, M., and Schürmann, G.: Consistent assimilation of multiple data streams in a carbon cycle data assimilation system, *Geosci. Model Dev.*, 9, 3569–3588, <https://doi.org/10.5194/gmd-9-3569-2016>, 2016.
- Martens, B., Miralles, D. G., Lievens, H., van der Schalie, R., de Jeu, R. A. M., Fernández-Prieto, D., Beck, H. E., Dorigo, W. A., and Verhoest, N. E. C.: GLEAM v3: satellite-based land evaporation and root-zone soil moisture, *Geosci. Model Dev.*, 10, 1903–1925, <https://doi.org/10.5194/gmd-10-1903-2017>, 2017.
- Mathieu, P.-P. and O'Neill, A.: Data assimilation: From photon counts to Earth System forecasts, *Remote Sens. Environ.*, 112, 1258–1267, <https://doi.org/10.1016/j.rse.2007.02.040>, 2008.
- Matthews, H. D., Eby, M., Ewen, T., Friedlingstein, P., and Hawkins, B. J.: What determines the magnitude of carbon cycle-climate feedbacks?, *Global Biogeochem. Cy.*, 21, GB2012, <https://doi.org/10.1029/2006GB002733>, 2007.
- McCallum, I., Wagner, W., Schmullius, C., Shvidenko, A., Obersteiner, M., Fritz, S., and Nilsson, S.: Comparison of four global FAPAR datasets over Northern Eurasia for the year 2000, *Remote Sens. Environ.*, 114, 941–949, <https://doi.org/10.1016/j.rse.2009.12.009>, 2010.
- Minh, D. H. T., Toan, T. L., Rocca, F., Tebaldini, S., d'Alessandro, M. M., and Villard, L.: Relating P-Band Synthetic Aperture Radar Tomography to Tropical Forest Biomass, *IEEE T. Geosci. Remote.*, 52, 967–979, <https://doi.org/10.1109/TGRS.2013.2246170>, 2014.
- Mitchard, E. T. A., Feldpausch, T. R., Brienen, R. J. W., Lopez-Gonzalez, G., Monteagudo, A., Baker, T. R., Lewis, S. L., Lloyd, J., Quesada, C. A., Gloor, M., ter Steege, H., Meir, P., Alvarez, E., Araujo-Murakami, A., Aragao, L. E. O. C., Arroyo, L., Aymard, G., Banki, O., Bonal, D., Brown, S., Brown, F. I., Cerón, C. E., Chama Moscoso, V., Chave, J., Comiskey, J. A., Cornejo, F., Corrales Medina, M., Da Costa, L., Costa, F. R. C., Di Fiore, A., Domingues, T. F., Erwin, T. L., Fredrickson, T., Higuchi, N., Honorio Coronado, E. N., Killeen, T. J., Laurance, W. F., Levis, C., Magnusson, W. E., Marimon, B. S., Marimon Junior, B. H., Mendoza Polo, I., Mishra, P., Nascimento, M. T., Neill, D., Nunez Vargas, M. P., Palacios, W. A., Parada, A., Pardo Molina, G., Peña-Claros, M., Pitman, N., Peres, C. A., Poorter, L., Prieto, A., Ramirez-Angulo, H., Restrepo Correa, Z., Roopsind, A., Roucoux, K. H., Rudas, A., Salomao, R. P., Schietti, J., Silveira, M., de Souza, P. F., Steininger, M. K., Stropp, J., Terborgh, J., Thomas, R., Toledo, M., Torres-Lezama, A., van Andel, T. R., van der Heijden, G. M. F., Vieira, I. C. G., Vieira, S., Vilanova-Torre, E., Vos, V. A., Wang, O., Zartman, C. E., Malhi, Y., and Phillips, O. L.: Markedly divergent estimates of Amazon forest carbon density

- from ground plots and satellites, *Global Ecol. Biogeogr.*, 23, 935–946, <https://doi.org/10.1111/geb.12168>, 2014.
- Moore, D. J., Hu, J., Sacks, W. J., Schimel, D. S., and Monson, R. K.: Estimating transpiration and the sensitivity of carbon uptake to water availability in a subalpine forest using a simple ecosystem process model informed by measured net CO₂ and H₂O fluxes, *Agr. Forest Meteorol.*, 148, 1467–1477, <https://doi.org/10.1016/j.agrformet.2008.04.013>, 2008.
- Muñoz, A. A., Barichivich, J., Christie, D. A., Dorigo, W., Sauchyn, D., González-Reyes, A., Villalba, R., Lara, A., Riquelme, N., and González, M. E.: Patterns and drivers of Araucaria araucana forest growth along a biophysical gradient in the northern Patagonian Andes: Linking tree rings with satellite observations of soil moisture, *Aust. Ecol.*, 39, 158–169, <https://doi.org/10.1111/aec.12054>, 2014.
- Mynden, R., Hoffman, S., Knyazikhin, Y., Privette, J., Glassy, J., Tian, Y., Wang, Y., Song, X., Zhang, Y., Smith, G., Lotsch, A., Friedl, M., Morisette, J., Votava, P., Nemani, R., and Running, S.: Global products of vegetation leaf area and fraction absorbed {PAR} from year one of {MODIS} data, *Remote Sens. Environ.*, 83, 214 – 231, [https://doi.org/10.1016/S0034-4257\(02\)00074-3](https://doi.org/10.1016/S0034-4257(02)00074-3), 2002.
- Naeimi, V., Scipal, K., Bartalis, Z., Hasenauer, S., and Wagner, W.: An Improved Soil Moisture Retrieval Algorithm for ERS and METOP Scatterometer Observations, *IEEE T. Geosci. Remote*, 47, 1999–2013, <https://doi.org/10.1109/Tgrs.2009.2011617>, 2009.
- Noël, S., Bramstedt, K., Rozanov, A., Bovensmann, H., and Burrows, J. P.: Stratospheric methane profiles from SCIAMACHY solar occultation measurements derived with onion peeling DOAS, *Atmos. Meas. Tech.*, 4, 2567–2577, <https://doi.org/10.5194/amt-4-2567-2011>, 2011.
- Noël, S., Bramstedt, K., Hilker, M., Liebing, P., Plieninger, J., Reuter, M., Rozanov, A., Sioris, C. E., Bovensmann, H., and Burrows, J. P.: Stratospheric CH₄ and CO₂ profiles derived from SCIAMACHY solar occultation measurements, *Atmos. Meas. Tech.*, 9, 1485–1503, <https://doi.org/10.5194/amt-9-1485-2016>, 2016.
- Norton, A., Rayner, P. J., Scholze, M., and Koffi, E.: Global Gross Primary Productivity for 2015 inferred from OCO-2 SIF and a Carbon-Cycle Data Assimilation System, Abstract B53L-01 presented at 2016, Fall Meeting, AGU, San Francisco, CA, 12–16 December, 2016.
- Ochsner, T., Cosh, M., Cuenca, R., Dorigo, W., Draper, C., Hagimoto, Y., Kerr, Y., Larson, K., Njoku, E., Small, E., and Zreda, M.: State of the art in large-scale soil moisture monitoring, *Soil Sci. Soc. Am. J.*, 77, 1888–1919, <https://doi.org/10.2136/sssaj2013.03.0093>, 2013.
- Owe, M., de Jeu, R., and Holmes, T.: Multisensor historical climatology of satellite-derived global land surface moisture, *J. Geophys. Res.-Earth*, 113, F01002, <https://doi.org/10.1029/2007jf000769>, 2008.
- Parazoo, N. C., Bowman, K., Frankenberg, C., Lee, J.-E., Fisher, J. B., Worden, J., Jones, D. B. A., Berry, J., Collatz, G. J., Baker, I. T., Jung, M., Liu, J., Osterman, G., O'Dell, C., Sparks, A., Butz, A., Guerlet, S., Yoshida, Y., Chen, H., and Gerbig, C.: Interpreting seasonal changes in the carbon balance of southern Amazonia using measurements of XCO₂ and chlorophyll fluorescence from GOSAT, *Geophys. Res. Lett.*, 40, 2829–2833, <https://doi.org/10.1002/grl.50452>, 2013.
- Parinussa, R., Meesters, A., Liu, Y., Dorigo, W., Wagner, W., and De Jeu, R.: An analytical solution to estimate the error structure of a global soil moisture data set, *IEEE Geosci. Remote S.*, 8, 779–783, 2011.
- Parker, R., Boesch, H., Cogan, A., Fraser, A., Feng, L., Palmer, P. I., Messerschmidt, J., Deutscher, N., Griffith, D. W. T., Notholt, J., Wennberg, P. O., and Wunch, D.: Methane observations from the Greenhouse Gases Observing SATellite: Comparison to ground-based TCCON data and model calculations, *Geophys. Res. Lett.*, 38, L15807, <https://doi.org/10.1029/2011GL047871>, 2011.
- Pastorello, G. Z., Papale, D., Chu, H., Trotta, C., Agarwal, D. A., Canfora, E., Baldocchi, D. D., and Torn, M. S.: A new data set to keep a sharper eye on land-air exchanges, *EOS*, 98, <https://doi.org/10.1029/2017EO071597>, 2017.
- Peylin, P., Law, R. M., Gurney, K. R., Chevallier, F., Jacobson, A. R., Maki, T., Niwa, Y., Patra, P. K., Peters, W., Rayner, P. J., Rödenbeck, C., van der Laan-Luijkx, I. T., and Zhang, X.: Global atmospheric carbon budget: results from an ensemble of atmospheric CO₂ inversions, *Biogeosciences*, 10, 6699–6720, <https://doi.org/10.5194/bg-10-6699-2013>, 2013.
- Peylin, P., Bacour, C., MacBean, N., Leonard, S., Rayner, P., Kuppel, S., Koffi, E., Kane, A., Maignan, F., Chevallier, F., Ciais, P., and Prunet, P.: A new stepwise carbon cycle data assimilation system using multiple data streams to constrain the simulated land surface carbon cycle, *Geosci. Model Dev.*, 9, 3321–3346, <https://doi.org/10.5194/gmd-9-3321-2016>, 2016.
- Pickett-Heaps, C. A., Canadell, J. G., Briggs, P. R., Gobron, N., Haverd, V., Paget, M. J., Pinty, B., and Raupach, M. R.: Evaluation of six satellite-derived Fraction of Absorbed Photosynthetic Active Radiation (FAPAR) products across the Australian continent, *Remote Sens. Environ.*, 140, 241–256, <https://doi.org/10.1016/j.rse.2013.08.037>, 2014.
- Pinty, B. and Verstraete, M.: GEMI: A non-linear index to monitor global vegetation from satellites, *Vegetatio*, 101, 1335–1372, 1992.
- Pinty, B., Leprieur, C., and Verstraete, M. M.: Towards a quantitative interpretation of vegetation indices Part 1: Biophysical canopy properties and classical indices, *Remote Sensing Reviews*, 7, 127–150, <https://doi.org/10.1080/02757259309532171>, 1993.
- Pinty, B., Lavergne, T., Dickinson, R. E., Widlowski, J.-L., Gobron, N., and Verstraete, M. M.: Simplifying the Interaction of Land Surfaces with Radiation for Relating Remote Sensing Products to Climate Models, *J. Geophys. Res.-Atmos.*, 111, <https://doi.org/10.1029/2005JD005952>, 2006.
- Pinty, B., Lavergne, T., Voßbeck, M., Kaminski, T., Aussedat, O., Giering, R., Gobron, N., Taberner, M., Verstraete, M. M., and Widlowski, J.-L.: Retrieving surface parameters for climate models from Moderate Resolution Imaging Spectroradiometer (MODIS)-Multiangle Imaging Spectroradiometer (MISR) albedo products, *J. Geophys. Res.-Atmos.*, 112, D10116, <https://doi.org/10.1029/2006JD008105>, 2007.
- Pinty, B., Lavergne, T., Kaminski, T., Aussedat, O., Giering, R., Gobron, N., Taberner, M., Verstraete, M. M., Voßbeck, M., and Widlowski, J.-L.: Partitioning the solar radiant fluxes in forest canopies in the presence of snow, *J. Geophys. Res.-Atmos.*, 113, D04104, <https://doi.org/10.1029/2007JD009096>, 2008.

- Pinty, B., Lavergne, T., Widlowski, J.-L., Gobron, N., and Verstraete, M.: On the need to observe vegetation canopies in the near-infrared to estimate visible light absorption, *Remote Sens. Environ.*, 113, 10–23, <https://doi.org/10.1016/j.rse.2008.08.017>, 2009.
- Pinty, B., Andredakis, I., Clerici, M., Kaminski, T., Taberner, M., Verstraete, M. M., Gobron, N., Plummer, S., and Widlowski, J.-L.: Exploiting the MODIS albedos with the Two-stream Inversion Package (JRC-TIP): 1. Effective leaf area index, vegetation, and soil properties, *J. Geophys. Res.-Atmos.*, 116, D09105, <https://doi.org/10.1029/2010JD015372>, 2011a.
- Pinty, B., Clerici, M., Andredakis, I., Kaminski, T., Taberner, M., Verstraete, M. M., Gobron, N., Plummer, S., and Widlowski, J.-L.: Exploiting the MODIS albedos with the Two-stream Inversion Package (JRC-TIP): 2. Fractions of transmitted and absorbed fluxes in the vegetation and soil layers, *J. Geophys. Res.-Atmos.*, 116, D09106, <https://doi.org/10.1029/2010JD015373>, 2011b.
- Pinty, B., Clerici, M., Andredakis, I., Kaminski, T., Taberner, M., Verstraete, M. M., Gobron, N., Plummer, S., and Widlowski, J.-L.: Exploiting the MODIS albedos with the Two-stream Inversion Package (JRC-TIP): 2. Fractions of transmitted and absorbed fluxes in the vegetation and soil layers, *J. Geophys. Res.-Atmos.*, 116, <https://doi.org/10.1029/2010JD015373>, 2011c.
- Porcar-Castell, A., Tyystjärvi, E., Atherton, J., van der Tol, C., Flexas, J., Pfundel, E. E., Moreno, J., Frankenberg, C., and Berry, J. A.: Linking chlorophyll-*a* fluorescence to photosynthesis for remote sensing applications: mechanisms and challenges, *J. Exp. Bot.*, 65, 4065–4095, <https://doi.org/10.1093/jxb/eru191>, 2014.
- Post, H., Vrugt, J. A., Fox, A., Vereecken, H., and Hendriks Franssen, H.-J.: Estimation of Community Land Model parameters for an improved assessment of net carbon fluxes at European sites, *J. Geophys. Res.-Biogeosci.*, 122, 661–689, <https://doi.org/10.1002/2015JG003297>, 2017.
- Prentice, I. C., Liang, X., Medlyn, B. E., and Wang, Y.-P.: Reliable, robust and realistic: the three R's of next-generation land-surface modelling, *Atmos. Chem. Phys.*, 15, 5987–6005, <https://doi.org/10.5194/acp-15-5987-2015>, 2015.
- Raj, R., Hamm, N. A. S., Tol, C. V. D., and Stein, A.: Uncertainty analysis of gross primary production partitioned from net ecosystem exchange measurements, *Biogeosciences*, 13, 1409–1422, <https://doi.org/10.5194/bg-13-1409-2016>, 2016.
- Randerson, J. T., Hoffman, F. M., Thornton, P. E., Mahowald, N. M., Lindsay, K., Lee, Y.-H., Nevison, C. D., Doney, S. C., Bonan, G., Stockli, R., Covey, C., Running, S. W., and Fung, I. Y.: Systematic assessment of terrestrial biogeochemistry in coupled climate-carbon models, *Glob. Change Biol.*, 15, 2462–2484, <https://doi.org/10.1111/j.1365-2486.2009.01912.x>, 2009.
- Raoult, N. M., Jupp, T. E., Cox, P. M., and Luke, C. M.: Land-surface parameter optimisation using data assimilation techniques: the adJULES system V1.0, *Geosci. Model Dev.*, 9, 2833–2852, <https://doi.org/10.5194/gmd-9-2833-2016>, 2016.
- Raupach, M. R., Rayner, P. J., Barrett, D. J., DeFries, R. S., Heimann, M., Ojima, D. S., Quegan, S., and Schmullius, C. C.: Model-data synthesis in terrestrial carbon observation: methods, data requirements and data uncertainty specifications, *Glob. Change Biol.*, 11, 378–397, <https://doi.org/10.1111/j.1365-2486.2005.00917.x>, 2005.
- Rayner, P., Scholze, M., Knorr, W., Kaminski, T., Giering, R., and Widmann, H.: Two decades of terrestrial Carbon fluxes from a Carbon Cycle Data Assimilation System (CCDAS), *Global Biogeochem. Cy.*, 19, GB2026, <https://doi.org/10.1029/2004GB002254>, 2005.
- Rayner, P., Michalak, A. M., and Chevallier, F.: Fundamentals of Data Assimilation, *Geosci. Model Dev. Discuss.*, <https://doi.org/10.5194/gmd-2016-148>, in review, 2016.
- Reuter, M., Buchwitz, M., Schneising, O., Heymann, J., Bovensmann, H., and Burrows, J. P.: A method for improved SCIAMACHY CO₂ retrieval in the presence of optically thin clouds, *Atmos. Meas. Tech.*, 3, 209–232, <https://doi.org/10.5194/amt-3-209-2010>, 2010.
- Reuter, M., Bovensmann, H., Buchwitz, M., Burrows, J. P., Connor, B. J., Deutscher, N. M., Griffith, D. W. T., Heymann, J., Keppel-Aleks, G., Messerschmidt, J., Notholt, J., Petri, C., Robinson, J., Schneising, O., Sherlock, V., Velazco, V., Warneke, T., Wennberg, P. O., and Wunch, D.: Retrieval of atmospheric CO₂ with enhanced accuracy and precision from SCIAMACHY: Validation with FTS measurements and comparison with model results, *J. Geophys. Res.-Atmos.*, 116, D04301, <https://doi.org/10.1029/2010JD015047>, 2011.
- Reuter, M., Bösch, H., Bovensmann, H., Bril, A., Buchwitz, M., Butz, A., Burrows, J. P., O'Dell, C. W., Guerlet, S., Hasekamp, O., Heymann, J., Kikuchi, N., Oshchepkov, S., Parker, R., Pfeifer, S., Schneising, O., Yokota, T., and Yoshida, Y.: A joint effort to deliver satellite retrieved atmospheric CO₂ concentrations for surface flux inversions: the ensemble median algorithm EMMA, *Atmos. Chem. Phys.*, 13, 1771–1780, <https://doi.org/10.5194/acp-13-1771-2013>, 2013.
- Reuter, M., Buchwitz, M., Hilker, M., Heymann, J., Schneising, O., Pillai, D., Bovensmann, H., Burrows, J. P., Bösch, H., Parker, R., Butz, A., Hasekamp, O., O'Dell, C. W., Yoshida, Y., Gerbig, C., Nehrkorn, T., Deutscher, N. M., Warneke, T., Notholt, J., Hase, F., Kivi, R., Sussmann, R., Machida, T., Matsueda, H., and Sawa, Y.: Satellite-inferred European carbon sink larger than expected, *Atmos. Chem. Phys.*, 14, 13739–13753, <https://doi.org/10.5194/acp-14-13739-2014>, 2014.
- Reuter, M., Hilker, M., Schneising, O., Buchwitz, M., and Heymann, J.: ESA Climate Change Initiative (CCI) Comprehensive Error Characterisation Report: BESD full-physics retrieval algorithm for XCO₂ for the Essential Climate Variable (ECV) Greenhouse Gases (GHG), Version 2.0, available at: http://www.esa-ghg-cci.org/webfm_send/284(last access: 14 July 2017), 2016.
- Ricciuto, D. M., Davis, K. J., and Keller, K.: A Bayesian calibration of a simple carbon cycle model: The role of observations in estimating and reducing uncertainty, *Global Biogeochem. Cy.*, 22, GB2030, <https://doi.org/10.1029/2006GB002908>, 2008.
- Richardson, A. D., Mahecha, M. D., Falge, E., Kattge, J., Moffat, A. M., Papale, D., Reichstein, M., Stauch, V. J., Braswell, B. H., Churkina, G., Kruijt, B., and Hollinger, D. Y.: Statistical properties of random CO₂ flux measurement uncertainty inferred from model residuals, *Agr. Forest Meteorol.*, 148, 38–50, <https://doi.org/10.1016/j.agrformet.2007.09.001>, 2008.
- Richardson, A. D., Williams, M., Hollinger, D. Y., Moore, D. J. P., Dail, D. B., Davidson, E. A., Scott, N. A., Evans, R. S., Hughes, H., Lee, J. T., Rodrigues, C., and Savage, K.: Estimating parameters of a forest ecosystem C model with measurements of

- stocks and fluxes as joint constraints, *Oecologia*, 164, 25–40, <https://doi.org/10.1007/s00442-010-1628-y>, 2010.
- Rodell, M., Velicogna, I., and Famiglietti, J. S.: Satellite-based estimates of groundwater depletion in India, *Nature*, 460, 999–1002, <https://doi.org/10.1038/nature08238>, 2009.
- Rodriguez-Fernandez, N. J., Aires, F., Richaume, P., Kerr, Y. H., Prigent, C., Kolassa, J., Cabot, F., Jimenez, C., Mahmoodi, A., and Drusch, M.: Soil Moisture Retrieval Using Neural Networks: Application to SMOS, *IEEE T. Geosci. Remote*, 53, 5991–6007, <https://doi.org/10.1109/TGRS.2015.2430845>, 2015.
- Rogers, C. D.: Inverse Methods for Atmospheric Sounding: Theory and Practice, World Scientific Publishing, 2000.
- Saatchi, S., Mascaro, J., Xu, L., Keller, M., Yang, Y., Duffy, P., Espirito-Santo, F., Baccini, A., Chambers, J., and Schimel, D.: Seeing the forest beyond the trees, *Global Ecol. Biogeogr.*, 24, 606–610, <https://doi.org/10.1111/geb.12256>, 2015.
- Saatchi, S. S., Harris, N. L., Brown, S., Lefsky, M., Mitchard, E. T. A., Salas, W., Zutta, B. R., Buermann, W., Lewis, S. L., Hagen, S., Petrova, S., White, L., Silman, M., and Morel, A.: Benchmark map of forest carbon stocks in tropical regions across three continents, *P. Natl. Acad. Sci. USA*, 108, 9899–9904, <https://doi.org/10.1073/pnas.1019576108>, 2011.
- Santoro, M., Beer, C., Cartus, O., Schmullius, C., Shvidenko, A., McCallum, I., Wegmüller, U., and Wiesmann, A.: Retrieval of growing stock volume in boreal forest using hyper-temporal series of Envisat ASA ScanSAR backscatter measurements, *Remote Sens. Environ.*, 115, 490–507, <https://doi.org/10.1016/j.rse.2010.09.018>, 2011.
- Santoro, M., Cartus, O., Fransson, J. E., Shvidenko, A., McCallum, I., Hall, R. J., Beaudoin, A., Beer, C., and Schmullius, C.: Estimates of Forest Growing Stock Volume for Sweden, Central Siberia, and Quebec using Envisat Advanced Synthetic Aperture Radar Backscatter Data, *Remote Sensing*, 5, 4503–4532, <https://doi.org/10.3390/rs5094503>, 2013.
- Santoro, M., Beaudoin, A., Beer, C., Cartus, O., Fransson, J. E., Hall, R. J., Pathe, C., Schmullius, C., Schepaschenko, D., Shvidenko, A., Thurner, M., and Wegmueller, U.: Forest growing stock volume of the northern hemisphere: Spatially explicit estimates for 2010 derived from Envisat ASAR, *Remote Sens. Environ.*, 168, 316–334, <https://doi.org/10.1016/j.rse.2015.07.005>, 2015.
- Schneising, O., Buchwitz, M., Burrows, J. P., Bovensmann, H., Reuter, M., Notholt, J., Macatangay, R., and Warneke, T.: Three years of greenhouse gas column-averaged dry air mole fractions retrieved from satellite – Part 1: Carbon dioxide, *Atmos. Chem. Phys.*, 8, 3827–3853, <https://doi.org/10.5194/acp-8-3827-2008>, 2008.
- Schneising, O., Buchwitz, M., Burrows, J. P., Bovensmann, H., Bergamaschi, P., and Peters, W.: Three years of greenhouse gas column-averaged dry air mole fractions retrieved from satellite – Part 2: Methane, *Atmos. Chem. Phys.*, 9, 443–465, <https://doi.org/10.5194/acp-9-443-2009>, 2009.
- Schneising, O., Buchwitz, M., Reuter, M., Heymann, J., Bovensmann, H., and Burrows, J. P.: Long-term analysis of carbon dioxide and methane column-averaged mole fractions retrieved from SCIAMACHY, *Atmos. Chem. Phys.*, 11, 2863–2880, <https://doi.org/10.5194/acp-11-2863-2011>, 2011.
- Schneising, O., Bergamaschi, P., Bovensmann, H., Buchwitz, M., Burrows, J. P., Deutscher, N. M., Griffith, D. W. T., Heymann, J., Macatangay, R., Messerschmidt, J., Notholt, J., Rettinger, M., Reuter, M., Sussmann, R., Velazco, V. A., Warneke, T., Wennberg, P. O., and Wunch, D.: Atmospheric greenhouse gases retrieved from SCIAMACHY: comparison to ground-based FTS measurements and model results, *Atmos. Chem. Phys.*, 12, 1527–1540, <https://doi.org/10.5194/acp-12-1527-2012>, 2012.
- Scholze, M., Kaminski, T., Rayner, P., Knorr, W., and Giering, R.: Propagating uncertainty through prognostic CCDAS simulations, *J. Geophys. Res.*, 112, D17305, <https://doi.org/10.1029/2007JD008642>, 2007.
- Scholze, M., Allen, I., Bill Collins, B., Cornell, S., Huntingford, C., Joshi, M., Lowe, J., Smith, R., Ridgwell, A., and Wild, O.: Understanding the Earth System – Global Change Science for Application, chap. 5 Earth System Models: a tool to understand changes in the Earth System, Cambridge University Press, Cambridge, UK, 2012.
- Scholze, M., Kaminski, T., Knorr, W., Blessing, S., Vossbeck, M., Grant, J., and Scipal, K.: Simultaneous assimilation of {SMOS} soil moisture and atmospheric {CO₂} in-situ observations to constrain the global terrestrial carbon cycle, *Remote Sens. Environ.*, 180, 334–345, <https://doi.org/10.1016/j.rse.2016.02.058>, 2016.
- Schürmann, G. J., Kaminski, T., Köstler, C., Carvalhais, N., Voßbeck, M., Kattge, J., Giering, R., Rödenbeck, C., Heimann, M., and Zaehle, S.: Constraining a land-surface model with multiple observations by application of the MPI-Carbon Cycle Data Assimilation System V1.0, *Geosci. Model Dev.*, 9, 2999–3026, <https://doi.org/10.5194/gmd-9-2999-2016>, 2016.
- Scipal, K., Holmes, T., de Jeu, R., Naeimi, V., and Wagner, W.: A possible solution for the problem of estimating the error structure of global soil moisture data sets, *Geophys. Res. Lett.*, 35, L24403, <https://doi.org/10.1029/2008gl035599>, 2008.
- Tao, X., Liang, S., and Wang, D.: Assessment of five global satellite products of fraction of absorbed photosynthetically active radiation: Intercomparison and direct validation against ground-based data, *Remote Sens. Environ.*, 163, 270–285, <https://doi.org/10.1016/j.rse.2015.03.025>, 2015.
- Tarantola, A.: Inverse Problem Theory and methods for model parameter estimation, SIAM, Philadelphia, 2005.
- Thum, T., MacBean, N., Peylin, P., Bacour, C., Santaren, D., Longdoz, B., Loustau, D., and Ciais, P.: The potential benefit of using forest biomass data in addition to carbon and water fluxes measurements to constrain ecosystem model parameters: case studies at two temperate forest sites, *Agr. Forest Meteorol.*, 234, 48–65, 2017.
- Thurner, M., Beer, C., Santoro, M., Carvalhais, N., Wutzler, T., Schepaschenko, D., Shvidenko, A., Kompter, E., Ahrens, B., Levick, S. R., and Schmullius, C.: Carbon stock and density of northern boreal and temperate forests, *Global Ecol. Biogeogr.*, 23, 297–310, <https://doi.org/10.1111/geb.12125>, 2014.
- Trudinger, C. M., Raupach, M. R., Rayner, P. J., Kattge, J., Liu, Q., Pak, B., Reichstein, M., Renzullo, L., Richardson, A. D., Roxburgh, S. H., Styles, J., Wang, Y. P., Briggs, P., Barrett, D., and Nikolova, S.: OptIC project: An intercomparison of optimization techniques for parameter estimation in terrestrial biogeochemical models, *J. Geophys. Res.-Biogeo.*, 112, G02027, <https://doi.org/10.1029/2006JG000367>, 2007.
- Van der Molen, M. K., de Jeu, R. A. M., Wagner, W., van der Velde, I. R., Kolari, P., Kurbatova, J., Varlagin, A., Maximov, T. C.,

- Kononov, A. V., Ohta, T., Kotani, A., Krol, M. C., and Peters, W.: The effect of assimilating satellite-derived soil moisture data in SiBCASA on simulated carbon fluxes in Boreal Eurasia, *Hydrol. Earth Syst. Sci.*, 20, 605–624, <https://doi.org/10.5194/hess-20-605-2016>, 2016.
- Veefkind, J., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H., de Haan, J., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visscher, H., and Levelt, P.: {TROPOMI} on the {ESA} Sentinel-5 Precursor: A {GMES} mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote Sens. Environ.*, 120, 70–83, <https://doi.org/10.1016/j.rse.2011.09.027>, 2012.
- Villard, L. and Toan, T. L.: Relating P-Band SAR Intensity to Biomass for Tropical Dense Forests in Hilly Terrain: γ^0 or t^0 ?, *IEEE J. Sel. Top. Appl.*, 8, 214–223, <https://doi.org/10.1109/JSTARS.2014.2359231>, 2015.
- Wagner, W., Lemoine, G., and Rott, H.: A method for estimating soil moisture from ERS scatterometer and soil data, *Remote Sens. Environ.*, 70, 191–207, 1999.
- Wagner, W., Hahn, S., Kidd, R., Melzer, T., Bartalis, Z., Hassenauer, S., Figa-Saldaña, J., de Rosnay, P., Jann, A., Schneider, S., Komma, J., Kubu, G., Brugger, K., Aubrecht, C., Z'uger, J., Gangkofner, U., Kienberger, S., Brocca, L., Wang, Y., Bl'oschl, G., Eitzinger, J., and Steinnocher, K.: The ASCAT Soil Moisture Product: A Review of its Specifications, Validation Results, and Emerging Applications, *Meteorol. Z.*, 22, 5–33, <https://doi.org/10.1127/0941-2948/2013/0399>, 2013.
- Walther, S., Voigt, M., Thum, T., Gonsamo, A., Zhang, Y., Koehler, P., Jung, M., Varlagin, A., and Guanter, L.: Satellite chlorophyll fluorescence measurements reveal large-scale decoupling of photosynthesis and greenness dynamics in boreal evergreen forests, *Glob. Change Biol.*, 2979–2996, <https://doi.org/10.1111/gcb.13200>, 2015.
- Wang, Y. P., Leuning, R., Cleugh, H., and Coppin, P. A.: Parameter estimation in surface exchange models using non-linear inversion: How many parameters can we estimate and which measurements are most useful?, *Glob. Change Biol.*, 7, 495–510, 2001.
- Widlowski, J.-L.: On the bias of instantaneous {FAPAR} estimates in open-canopy forests, *Agr. Forest Meteorol.*, 150, 1501–1522, <https://doi.org/j.agrformet.2010.07.011>, 2010.
- Williams, M., Schwarz, P. A., Law, B. E., Irvine, J., and Kurpius, M. R.: An improved analysis of forest carbon dynamics using data assimilation, *Glob. Change Biol.*, 11, 89–105, <https://doi.org/10.1111/j.1365-2486.2004.00891.x>, 2005.
- WMO: Greenhouse Gas Bulletin, The State of Greenhouse Gases in the Atmosphere Based on Global Observations through 2014, World Meteorological Organization, No. 11, 9 November, 2015.
- Wolanin, A., Rozanov, V., Dinter, T., Noël, S., Vountas, M., Burrows, J., and Bracher, A.: Global retrieval of marine and terrestrial chlorophyll fluorescence at its red peak using hyperspectral top of atmosphere radiance measurements: Feasibility study and first results, *Remote Sens. Environ.*, 166, 243–261, <https://doi.org/10.1016/j.rse.2015.05.018>, 2015.
- Wunch, D., Toon, G. C., Wennberg, P. O., Wofsy, S. C., Stephens, B. B., Fischer, M. L., Uchino, O., Abshire, J. B., Bernath, P., Biraud, S. C., Blavier, J.-F. L., Boone, C., Bowman, K. P., Brownell, E. V., Campos, T., Connor, B. J., Daube, B. C., Deutscher, N. M., Diao, M., Elkins, J. W., Gerbig, C., Gottlieb, E., Griffith, D. W. T., Hurst, D. F., Jiménez, R., Keppel-Aleks, G., Kort, E. A., Macatangay, R., Machida, T., Matsueda, H., Moore, F., Morino, I., Park, S., Robinson, J., Roehl, C. M., Sawa, Y., Sherlock, V., Sweeney, C., Tanaka, T., and Zondlo, M. A.: Calibration of the Total Carbon Column Observing Network using aircraft profile data, *Atmos. Meas. Tech.*, 3, 1351–1362, <https://doi.org/10.5194/amt-3-1351-2010>, 2010.
- Wunch, D., Toon, G. C., Blavier, J.-F. L., Washenfelder, R. A., Notholt, J., Connor, B. J., Griffith, D. W. T., Sherlock, V., and Wennberg, P. O.: The Total Carbon Column Observing Network, *Philos. T. Roy. Soc. A*, 369, 2087–2112, <https://doi.org/10.1098/rsta.2010.0240>, 2011.
- Xiong, X., Barnet, C., Maddy, E., Wofsy, S., Chen, L., Karion, A., and Sweeney, C.: Detection of methane depletion associated with stratospheric intrusion by atmospheric infrared sounder (AIRS), *Geophys. Res. Lett.*, 40, 2455–2459, <https://doi.org/10.1002/grl.50476>, 2013.
- Yoshida, Y., Kikuchi, N., Morino, I., Uchino, O., Oshchepkov, S., Bril, A., Saeki, T., Schutgens, N., Toon, G. C., Wunch, D., Roehl, C. M., Wennberg, P. O., Griffith, D. W. T., Deutscher, N. M., Warneke, T., Notholt, J., Robinson, J., Sherlock, V., Connor, B., Rettinger, M., Sussmann, R., Ahonen, P., Heikkinen, P., Kyrö, E., Mendonca, J., Strong, K., Hase, F., Dohe, S., and Yokota, T.: Improvement of the retrieval algorithm for GOSAT SWIR XCO₂ and XCH₄ and their validation using TCCON data, *Atmos. Meas. Tech.*, 6, 1533–1547, <https://doi.org/10.5194/amt-6-1533-2013>, 2013..
- Zwieback, S., Su, C.-H., Gruber, A., Dorigo, W. A., and Wagner, W.: The Impact of Quadratic Nonlinear Relations between Soil Moisture Products on Uncertainty Estimates from Triple Collocation Analysis and Two Quadratic Extensions, *J. Hydrometeorol.*, 17, 1725–1743, <https://doi.org/10.1175/JHM-D-15-0213.1>, 2016.