

J. Blitzstein's & Jessica Hwang's Introduction to Probability - A partial
solutions manual

George Lydakis

June 3, 2026

Contents

1	Probability and counting	5
1.1	Counting	5
1.2	Story Proofs	7
1.3	Naive definition of probability	10

Chapter 1

Probability and counting

1.1 Counting

Exercise 1

How many ways are there to permute the letters in the word MISSISSIPPI?

Solution.

The word has 11 letters and the following letter frequencies. M: 1, I: 4, S: 4, P: 2. We can find the number of permutations by counting as follows: pick the places for the M's first, then out of the remaining places pick the places for P's, then for S's, and then the places for I's are determined uniquely. Then, apply the multiplication rule. Note that within each selection of places, the order doesn't matter since we don't distinguish between "instances" of the same letter. We thus have that the number of permutations is:

$$\binom{11}{1} \cdot \binom{10}{2} \cdot \binom{8}{4} = 11 \cdot \frac{10!}{8! \cdot 2!} \cdot \frac{8!}{4! \cdot 4!} = 11 \cdot \frac{10!}{4! \cdot 4! \cdot 2!} = 11 \cdot \frac{10 \cdot 9 \dots 5}{4! \cdot 2} = 34,650$$

Exercise 4

A *round-robin* tournament is being held with n tennis players; this means that every player will play against every other player exactly once.

- (a) How many possible outcomes are there for the tournament (the outcome lists out who won and who lost for each game)?
- (b) How many games are played in total?

Solution.

(a) Consider ordering the players in some arbitrary but fixed order. Then we can count the number of outcomes as follows. For the first player, there are 2 possible outcomes (win/lose) against all other $n - 1$ players. This is equivalent to sampling with replacement $n - 1$ times from a set of two elements. For the second player in the ordering, the outcome against the first player has already been decided/accounted for, so there are $n - 2$ outcomes against all other players. We continue this way and apply the multiplication rule to obtain the total number of outcomes:

$$\prod_{k=1}^n 2^{n-k} = 2^{\sum_{k=1}^n n-k} = 2^{\sum_{k=1}^{n-1} k} = 2^{\frac{n(n-1)}{2}}$$

(b) The first player in the ordering plays in $n - 1$ games. The second player plays in $n - 2$ games *more*, since the match against player 1 was already accounted for. We continue this way to obtain the total number of games:

$$\sum_{k=1}^n (n - k) = \frac{n(n - 1)}{2}$$

Notice that this is consistent with the answer from (a): another way of getting the number of possible outcomes is listing the number of different games between two players, and assigning a win/lose outcome to each.

Exercise 5

A *knock-out tournament* is being held with 2^n tennis players. This means that for each round, the winners move on to the next round and the losers are eliminated, until only one person remains. For example, if initially there were $2^4 = 16$ players, then there are 8 games on the first round, then the 8 winners move on to round 2, then the 4 winners move on to round 3, then the 2 winners move on to round 4, the winner of which is declared the winner of the tournament. (There are various systems for determining who plays whom within a round, but these do not matter for this problem.)

- (a) How many rounds are there?
 - (b) Count how many games in total are played, by adding up the numbers of games played in each round.
 - (c) Count how many games in total are played, this time by directly thinking about it without doing almost any calculation.
- Hint: How many players need to be eliminated?

Solution.

- (a) The number of players per round is halved every time, until two players remain, or, in other words, until the exponent of 2^k drops to 1. This corresponds of course to $n - 1$ rounds.
- (b) In each round, every winner plays exactly one game against exactly one loser. In other words, out of 2^k players of the round, 2^{k-1} games are formed. Since there are $n - 1$ rounds, the total number of games is:

$$\sum_{k=1}^n 2^{k-1} = \sum_{k=0}^{n-1} 2^k = 2^n - 1$$

- (c) In order to obtain the final winner, $2^n - 1$ players will need to be eliminated (all but one). Every game in the tournament has exactly one loser, and each time someone loses they never play again, which means that the number of games equals the number of eliminated players, and so we obtain $2^n - 1$ once again.

Exercise 10

To fulfill the requirements for a certain degree, a student can choose to take any 7 out of a list of 20 courses, with the constraint that at least 1 of the 7 courses must be a statistics course. Suppose that 5 out of the 20 courses are statistics courses.

- (a) How many choices are there for which 7 courses to take?
- (b) Explain intuitively why the answer to (a) is not $\binom{5}{1} \cdot \binom{19}{6}$.

Solution.

- (a) A way to compute this is to compute the possible ways in which a student can pick 7 courses, *none* of which is statistics courses, and then subtracting it from the total number of choices. Picking 7 courses with no statistics ones means selecting 7 distinct courses from a pool of $20 - 5 = 15$. The total number of choices —without constraints— can be computed as choosing 7 out of 20. This means that the number of choices for completing the degree is:

$$\binom{20}{7} - \binom{15}{7} = \frac{1}{7!} \left(\frac{20!}{13!} - \frac{15!}{8!} \right) = \frac{1}{7!} (20 \cdot 19 \cdot \dots \cdot 14 - 15 \cdot 14 \cdot \dots \cdot 9) = 71,085$$

- (b) The expression here arises from the following line of thought: first, pick one statistics course ($\binom{5}{1}$ ways to do this), and then pick 6 more courses out of the remaining 19 ones. However, this results in double counting: order does not matter here, but picking, for example, statistics course #1 as the mandatory one, and then statistics course #2 as the first elective will be counted separately from picking statistics course #2 as the mandatory one and statistics course #1 as the first elective.

Exercise 12

Four players, named A, B, C, D, are playing a card game. A standard, well-shuffled deck of cards is dealt to the players (so each player receives a 13-card hand).

- (a) How many possibilities are there for the hand that player A will get? (Within a hand, the order in which cards were received doesn't matter).
 (b) How many possibilities are there overall for what hands everyone will get, assuming that it matters which player gets which hand, but not the order of cards within a hand?

Solution.

- (a) This is simply $\binom{52}{13}$: we choose 13 cards out of 52, without repetition, and without the order mattering.
 (b) We can compute this by fixing the order in which we examine the players as, say, A, B, C, D. For player A, there are $\binom{52}{13}$ possible hands. For each of these hands, there remain $52 - 13 = 39$ cards out of which the 13 cards for player B are picked. We continue this way (and using the multiplication rule) to compute the possible number of hands for everyone as:

$$\binom{52}{13} \cdot \binom{39}{13} \cdot \binom{26}{13} \cdot \binom{13}{13} = \binom{52}{13} \cdot \binom{39}{13} \cdot \binom{26}{13}$$

Exercise 13

A certain casino uses 10 standard decks of cards mixed together into one big deck, which we will call a *superdeck*. Thus, the superdeck has $52 \cdot 10 = 520$ cards, with 10 copies of each card. How many different 10-card hands can be dealt from the superdeck? The order of the cards does not matter, nor does it matter which of the original 10 decks the cards came from. Express your answer as a binomial coefficient. [Hint: Bose-Einstein.]

Solution.

The Bose-Einstein counting problem asks how many ways are there to choose k times from a set of n objects with replacement, if order doesn't matter. Consider thinking about the problem here as drawing 10 times from a deck of 52 cards, noting down the card we drew and then putting it back for the next draw. We can then map results of this experiment bijectively to results of the experiment in the exercise: drawing card i from deck j corresponds to drawing card i on the j -th trial. We then see that the number of possible hands corresponds to choosing 10 times from the deck, *with* replacement, and *without* order mattering, which by the Bose-Einstein solution gives:

$$\binom{52 + 10 - 1}{10}$$

1.2 Story Proofs

Exercise 15

Give a story proof that $\sum_{k=0}^n \binom{n}{k} = 2^n$.

Solution.

Imagine having a set of n distinct objects and wanting to count the number of possible subsets. As we already know, this is equal to 2^n (for each object, choose to either include it or not in a subset). Another equivalent way of counting this would be to count all possible subsets of size $0, 1, 2, \dots, n$ and sum the resulting quantities. We don't care about the order of object selection, so the number of ways to choose a subset of size k is $\binom{n}{k}$, which means that summing $\sum_{k=0}^n \binom{n}{k}$ must equal 2^n .

Exercise 16

Show that for all positive integers n and k with $n \geq k$,

$$\binom{n}{k} + \binom{n}{k-1} = \binom{n+1}{k},$$

doing this in two ways: (a) algebraically and (b) with a story, giving an interpretation for why both sides count the same thing.

Hint for the story proof: Imagine an organization consisting of $n+1$ people, with one of them pre-designated as the president of the organization.

Solution.

(a) For an algebraic proof, notice that, from previous identities and some calculations:

$$\begin{aligned} \binom{n}{k} + \binom{n}{k-1} &= \binom{n}{n-k} + \binom{n}{n-k+1} = \frac{n!}{(n-k)! \cdot k!} + \frac{n!}{(k-1)! \cdot (n-k+1)!} \\ &= \frac{n!}{(k-1)!} \left(\frac{1}{k(n-k)!} + \frac{1}{(n-k+1)!} \right) = \frac{n!}{(k-1)!(n-k)!} \left(\frac{1}{k} + \frac{1}{n-k+1} \right) \\ &= \frac{n!}{(k-1)!(n-k)!} \cdot \frac{n+1}{k(n-k+1)} = \frac{(n+1)!}{k! \cdot (n-k+1)!} = \binom{n+1}{k} \end{aligned}$$

(b) For a story proof, think about it as follows. We want to choose a committee of k people from a total of $n+1$, which is clearly equal to the RHS. Another way of counting the possible ways to do this, is by choosing a pre-designated “president” of the committee, and the remaining $k-1$ members separately. Fix any president, which results in having to choose $k-1$ members from $n-1$ people ($\binom{n}{k-1}$ ways). Notice that this misses all committees in which this particular person is not present at all: this is equal to the number of ways to choose k members from the remaining n people ($\binom{n}{k}$). The sum of these two counts must equal precisely the number of possible committees. Notice, also, that in the second count we didn’t have to utilize the “president” argument.

Exercise 20

(a) Show using a story proof that

$$\binom{k}{k} + \binom{k+1}{k} + \binom{k+2}{k} + \dots + \binom{n}{k} = \binom{n+1}{k+1},$$

where n, k are positive integers with $n \geq k$. This is called the *hockey stick identity*. Hint: Imagine arranging a group of people by age, and then think about the oldest person in a chosen subgroup.

(b) Suppose that a large pack of Haribo gummi bears can have anywhere between 30 and 50 gummi bears. There are 5 delicious flavors: pineapple (clear), raspberry (red), orange (orange), strawberry (green, mysteriously), and lemon (yellow). There are 0 non-delicious flavors. How many possibilities are there for the composition of such a pack of gummi bears? You can leave your answer in terms of a couple of binomial coefficients, but not a sum of lots of binomial coefficients.

Solution.

(a) Suppose we have a group of $n+1$ people, from which we want to choose $k+1$ without caring about the order. We know that there are $\binom{n+1}{k+1}$ ways to do this. We will show that this equals the LHS by describing another way of getting all such possible choices. Arrange the people in the group by age, labeling them $\#1, \#2, \dots, \#n+1$. Then, every possible combination of people of size $k+1$ has an oldest person. If the oldest person is $\#1$, then there are $\binom{n}{k}$ ways to choose all remaining others. If the oldest person is $\#2$, then there are $\binom{n-1}{k}$ ways to choose all the others. This holds all the way to the oldest being $\#n-k$, in which case there is a unique choice of the rest ($\binom{k}{k}$). Since the group has size $k+1$, it cannot be that the oldest person is anyone after that point. Notice that this also does not “double-count” anything, since by choosing the oldest person first we guarantee that we never encounter the same combination again. This leads us to conclude that the resulting sum $\binom{n}{k} + \binom{n-1}{k} + \dots + \binom{k+1}{k} + \binom{k}{k}$ must equal $\binom{n+1}{k+1}$.

(b) We begin by noting that this is similar to the Bose-Einstein setting. If we think of having five objects (the gummi bear flavors), we are selecting, with replacement, between 30 and 50 from them in order to make a pack, and we only care about how many times each flavor is selected. Therefore, $k = 5$ always, and n varies between 30 and 50. The total number of possible packs must then equal:

$$\binom{5+30-1}{30} + \binom{5+31-1}{31} + \dots + \binom{5+50-1}{50} = \binom{34}{4} + \binom{35}{4} + \dots + \binom{54}{4},$$

where we use Example 1.5.1, $\binom{n}{k} = \binom{n}{n-k}$. We can simplify further by using the hockey stick identity from part (a):

$$\begin{aligned} \binom{34}{4} + \binom{35}{4} + \dots + \binom{54}{4} &= \binom{4}{4} + \binom{5}{4} + \dots + \binom{34}{4} + \dots + \binom{54}{4} - \left(\binom{4}{4} + \dots + \binom{33}{4} \right) \\ &= \binom{54+1}{4+1} - \binom{33+1}{4+1} = \binom{55}{5} - \binom{34}{5} \end{aligned}$$

Exercise 22

The Dutch mathematician R. J. Stroeker remarked:

Every beginning student of number theory surely must have marveled at the miraculous fact that for each natural number n the sum of the first n consecutive cubes is a perfect square.

Furthermore, it is the square of the sum of the first n positive integers! That is,

$$1^3 + 2^3 + \dots + n^3 = (1 + 2 + \dots + n)^2$$

Usually this identity is proven by induction, but that does not give much insight into why the result is true, nor does it help much if we wanted to compute the left-hand side but didn't already know this result. In this problem, you will give a story proof of the identity.

(a) Give a story proof of the identity

$$1 + 2 + \dots + n = \binom{n+1}{2}$$

Hint: Consider a round-robin tournament (see Exercise 4).

(b) Give a story proof of the identity

$$1^3 + 2^3 + \dots + n^3 = 6 \binom{n+1}{4} + 6 \binom{n+1}{3} + \binom{n+1}{2}$$

It is then just basic algebra (not required for this problem) to check that the square of the RHS in (a) is the RHS in (b).

Hint: Imagine choosing a number between 1 and n and then choosing 3 numbers between 0 and n smaller than the original number, with replacement. Then consider cases based on how many distinct numbers were chosen.

Solution.

(a) Imagine having $n + 1$ tennis players in a round-robin tournament, and wanting to count the total number of games that will be played (regardless of win-lose). This equals $\binom{n+1}{2}$, since every pair of players must play against each other. Another way of counting this is labeling the players from 1, 2, $\dots$, $n + 1$, and counting all games in which each player will play. For player 1, this equals n games, for player 2, it equals $n - 1$ (since we've already counted the game against 1), and so on. This means that the total number of games also equals $n + n - 1 + \dots + 2 + 1$, thus showing the required identity.

(b) Suppose we choose a number k between 1 and n , inclusive, and then sample 3 times, with replacement, a number between 0 and $k - 1$, inclusive. Suppose we also want to record and count results in the form "chosen number - first sample - second sample - third sample", where the order matters. For any given k , there are k^3 possible sampling outcomes, and so the total count we are interested in is $1^3 + 2^3 + \dots + n^3$.

Now consider a different way of counting this, based on how many distinct numbers are chosen. Clearly, there are three cases: one, two, and three distinct numbers. Choosing one distinct number is feasible whenever $k = 1, 2, \dots, n$, and by thinking about the fact that for a given k there are k choices, this can be counted as:

$$\binom{1}{1} + \binom{2}{1} + \dots + \binom{n}{1} = \binom{n+1}{2},$$

where we make use of the hockey stick identity from Exercise 1.20.

Similarly, choosing two distinct numbers is feasible whenever $k = 2, 3, \dots, n$, and by a similar observation yields:

$$\binom{2}{2} + \binom{3}{2} + \dots + \binom{n}{2} = \binom{n+1}{3}$$

Note, however, that since we're dealing with combinations, we do not account for the fact that in our count here *order matters*. To account for this, note that given a combination of 3 items, 6 permutations can be made, and therefore we need to multiply the above count by 6. Finally, choosing three distinct numbers is feasible when $k = 3, 4, \dots, n$ and yields:

$$\binom{3}{3} + \binom{4}{3} + \dots + \binom{n}{3} = \binom{n+1}{4},$$

where for the same reason as above we are under-counting by a factor of 6. Putting it all together, we end up with the equality:

$$1^3 + 2^3 + \dots + n^3 = 6\binom{n+1}{4} + 6\binom{n+1}{3} + \binom{n+1}{2}$$

1.3 Naive definition of probability

Exercise 23

Three people get into an empty elevator at the first floor of a building that has 10 floors. Each presses the button for their desired floor (unless one of the others has already pressed that button). Assume that they are equally likely to want to go to floors 2 through 10 (independently of each other). What is the probability that the buttons for 3 consecutive floors are pressed?

Solution.

Let us first examine the number of possible outcomes. We can think of an outcome as a sequence of three numbers, each from 2 to 10, inclusive. These can be repeated, since e.g. 223 would mean that the first person pressed the button for floor 2, the second person also wanted the second one and thus pressed no button, and the third pressed the button for floor 3. In total, there are then 9^3 events (sampling 3 times with replacement from a set of 9 elements). We can count the number of favorable outcomes by observing that if we take an outcome and sort it by floor numbers, we end up with 7 possible ways to get 3 consecutive floors:

$$(2, 3, 4), (3, 4, 5), \dots, (8, 9, 10)$$

Now, each of these sorted lists can originate from $3!$ outcomes, its permutations. This means that the favorable outcomes are $7 \cdot 3! = 42$, and the probability of the event we're interested in is:

$$\frac{42}{9^3} = \frac{42}{729} \approx 0.0576$$

Exercise 26

A survey is being conducted in a city with 1 million residents. It would be far too expensive to survey all of the residents, so a random sample of size 1000 is chosen (in practice, there are many challenges with sampling, such as obtaining a complete list of everyone in the city, and dealing with people who refuse to participate). The survey is conducted by choosing people one at a time, *with* replacement and with equal probabilities.

- (a) Explain how sampling with vs. without replacement here relates to the birthday problem.
- (b) Find the probability that at least one person will get chosen more than once.

Solution.

(a) Imagine that a year had, instead of 365 days, 1 million ones. Then, choosing 1,000 people *with* replacement is in essence an instance of the “general” birthday problem in a setting where a room has 1,000 people and a year has 1 million days. If instead we sample without replacement, it’s as if we’re choosing $1,000 < 10^6$ unique birthdays to create a room where no two people have the same birthday.

(b) Using the birthday problem result with the adjustments described in (a) we obtain the desired probability as:

$$1 - \frac{10^6 \cdot (10^6 - 1) \cdot \dots \cdot (10^6 - 1,000 + 1)}{10^{6 \cdot 1000}} = 1 - 1 \cdot \left(1 - \frac{1}{10^6}\right) \cdot \dots \cdot \left(1 - \frac{999}{10^6}\right) \approx 0.393$$

Exercise 31

Solution.

There are $\binom{N}{m}$ ways of choosing the second sample. We can think of forming a sample that satisfies our constraints as follows. First, select k out of the n tagged elk. There are $\binom{n}{k}$ ways of doing this. Then, select $m - k$ out of the $N - n$ untagged elk. There are $\binom{N-n}{m-k}$ ways of doing this. Then the total number of samples that satisfy our constraints is $\binom{N}{k} \cdot \binom{N-n}{m-k}$, one for each combination of two subsamples. We thus conclude that the the probability of choosing exactly k previously tagged elk in our new sample of m elk is:

$$\frac{\binom{n}{k} \cdot \binom{N-n}{m-k}}{\binom{N}{m}}$$

Exercise 40

There are n balls in a jar, labeled with the numbers $1, 2, \dots, n$. A total of k balls are drawn, one by one *with replacement*, to obtain a sequence of numbers.

- (a) What is the probability that the sequence obtained is strictly increasing?
- (b) What is the probability that the sequence obtained is increasing (but not necessarily strictly increasing, i.e., there can be repetitions)?

(a) We begin by noting that there n^k possible sequences in total. To count the number of strictly increasing ones, we note that for each k balls that all different from each other, there exists precisely one sequence that contains all of them and is strictly increasing: the sequence resulting from sorting them. The converse is of course also true, i.e., a strictly increasing sequence can be mapped uniquely to k balls selected *without* replacement and without caring about the order. We conclude that the problem is equivalent to counting the number of unique combinations of k out of n balls, which of course means that the requested probability is

$$\frac{\binom{n}{k}}{n^k}$$

(b) The reasoning here is similar. First, any increasing sequence can be mapped uniquely to a selection of k out of n balls where we count the number of times each ball was selected (corresponding to equalities in

the result). Conversely, any selection of k out of n balls with replacement, and where we count the number of times each ball is drawn can be uniquely mapped to an increasing sequence. This is again done by sorting the balls, except that now equality of adjacent elements in the result is allowed, and corresponds to a ball being selected more than one time. We conclude that the problem is equivalent to the Bose-Einstein counting problem, and so the requested probability is

$$\frac{\binom{n+k-1}{k}}{n^k}$$

Exercise 43

Show that for any events A, B ,

$$P(A) + P(B) - 1 \leq P(A \cap B) \leq P(A \cup B) \leq P(A) + P(B)$$

For each of these three inequalities, give a simple criterion for when the inequality is actually an equality (e.g., give a simple condition such that $P(A \cap B) = P(A \cup B)$ if only if the condition holds).

Solution.

Recall that $P(A \cup B) = P(A) + P(B) - P(A \cap B)$, and that probabilities are non-negative, meaning that

$$P(A \cup B) = P(A) + P(B) - P(A \cap B) \leq P(A) + P(B)$$

This becomes an equality precisely whenever $P(A \cap B) = 0$. We also know that $A \cap B \subset A \implies P(A \cap B) \leq P(A)$, and similarly $A \cap B \subset B \implies P(A \cap B) \leq P(B)$. Therefore,

$$2P(A \cap B) \leq P(A) + P(B) \implies P(A \cap B) \leq P(A) + P(B) - P(A \cap B) = P(A \cup B)$$

This becomes an equality precisely when the first two inequalities become equalities, i.e., when $P(A \cap B) = P(A) = P(B)$. Finally, we have that

$$P(A \cup B) \leq 1 \implies P(A) + P(B) - P(A \cap B) \leq 1 \implies P(A) + P(B) - 1 \leq P(A \cap B)$$

This becomes an equality precisely when $P(A \cup B) = 1$.

Exercise 44

Let A, B be events. The *difference* $B - A$ is defined to be the set of all elements of B that are not in A . Show that if $A \subset B$, then

$$P(B - A) = P(B) - P(A),$$

directly using the axioms of probability.

Solution.

If $A \subset B$, we can write $B = (B - A) \cup A$, where $A, B - A$ are disjoint. Then, using the second axiom of probability, we obtain that

$$P(B) = P((B - A) \cup A) = P(B - A) + P(A) \implies P(B - A) = P(B) - P(A)$$

Exercise 48

Arby has a belief system assigning a number $P_{Ar}(A)$ between 0 and 1 to every event A (for some sample space). This represents Arby's degree of belief about how likely A is to occur. For any event A , Arby is willing to pay a price of $1000 \cdot P_{Ar}(A)$ dollars to buy a certificate such as the one shown below:

Certificate:

The owner of this certificate can redeem it for \$1000 if A occurs. No value if A does not occur, except as required by federal, state, or local law. No expiration date.

Likewise, Arby is willing to sell such a certificate at the same price. Indeed, Arby is willing to buy or sell any number of certificates at this price, as Arby considers it the “fair” price. Arby stubbornly refuses to accept the axioms of probability. In particular, suppose that there are two *disjoint* events A, B with

$$P_{Ar}(A \cup B) \neq P_{Ar}(A) + P_{Ar}(B)$$

Show how to make Arby go bankrupt, by giving a list of transactions Arby is willing to make that will *guarantee* that Arby will lose money (you can assume it will be known whether A occurred and whether B occurred the day after any certificates are bought/sold).

Solution.

Suppose first that $P_{Ar}(A \cup B) > P_{Ar}(A) + P_{Ar}(B)$. Consider then the following transactions:

- We, as an opposing party, offer to sell Arby a certificate for $A \cup B$ at his nominated price.
- We then offer to buy from him a certificate for A and a certificate for B .
- In total, this means that Arby has incurred a loss of $-P_{Ar}(A \cup B) + P_{Ar}(A) + P_{Ar}(B) < 0$, in the hopes, of course, of making money when the events are realized.

The following day, and since A, B are disjoint, there are only three possible outcomes:

- Neither A nor B occur. In this case, Arby neither makes nor loses any more money.
- A occurs, in which case we have to pay Arby \$1000, but he also has to pay us \$1000. This therefore means that he again has no profit or loss.
- B occurs, in which case exactly the same observations apply.

We therefore conclude that this series of transactions is *guaranteed* to decrease Arby's balance, which means that if we were to repeat it an appropriate number times (by the problem setting, “Arby is willing to buy or sell any number of certificates”), Arby goes bankrupt regardless of his initial money. If, instead, it was the case that $P_{Ar}(A \cup B) < P_{Ar}(A) + P_{Ar}(B)$, we have but to reverse the strategy presented above, buying certificates for $A \cup B$ and selling certificates for A and B . Arby's reasoning will therefore cause him to go bankrupt in all cases.