

# **Probability and Statistics**

UG2, Core course, IIIT,H

Pawan Kumar

IIIT, Hyderabad

November 16, 2021

- ① Joint Distributions: Two Random Variables
  - Conditional PMF and Conditional CDF
  - Independent Random Variables
  - Conditional Expectation
  - Functions of Two Random Variables
- ② Joint Continuous Random Variables
  - Solved Problems
- ③ Multiple Random Variables

- Joint PDF and Joint CDF of Multiple Random Variables
- Sums of Random Variables
- Random Vectors
- Functions of Random Vectors and Method of Transformations
- ④ Law of Large Numbers
  - Sample Mean, Expectation and Variance
  - Weak Law of Large Numbers

## Outline

### ① Joint Distributions: Two Random Variables

Conditional PMF and Conditional CDF

Independent Random Variables

Conditional Expectation

Functions of Two Random Variables

### ② Joint Continuous Random Variables

### ③ Multiple Random Variables

### ④ Law of Large Numbers

## Conditional PMF and Conditional CDF...

## Conditional PMF and Conditional CDF...

Example Motivation for Conditional PMF and CDF

## Conditional PMF and Conditional CDF...

### Example Motivation for Conditional PMF and CDF

I roll a fair die.

## Conditional PMF and Conditional CDF...

### Example Motivation for Conditional PMF and CDF

I roll a fair die. Let  $X$  be the observed number.

## Conditional PMF and Conditional CDF...

### Example Motivation for Conditional PMF and CDF

I roll a fair die. Let  $X$  be the observed number. Find the conditional PMF of  $X$  given that we know the observed number was less than 5.

Solution:

$$R_X = \{1, 2, 3, 4, 5, 6\}.$$

$$A = \{x < 5\}$$

$$P_{X|A}(1) = P(X=1 | A) = \frac{P(X=1, A)}{P(A)} = \frac{P(X=1, X < 5)}{P(A)}$$

$$P_{X|A}(2) = P_{X|A}(3) = P_{X|A}(4) = \frac{P(X=1)}{P(X < 5)} = \frac{1/6}{4/6} = 1/4$$
$$P_{X|A}(5) = P_{X|A}(6) = 0$$

## Conditional PMF and Conditional CDF...

## Conditional PMF and Conditional CDF...

Definition of Conditional PMF and Conditional CDF

## Conditional PMF and Conditional CDF...

### Definition of Conditional PMF and Conditional CDF

Let  $X$  be a **discrete** random variable and  $A$  be **any** event.

## Conditional PMF and Conditional CDF...

### Definition of Conditional PMF and Conditional CDF

Let  $X$  be a **discrete** random variable and  $A$  be **any** event. The **conditional PMF** of  $X$  given  $A$  is defined as

## Conditional PMF and Conditional CDF...

### Definition of Conditional PMF and Conditional CDF

Let  $X$  be a **discrete** random variable and  $A$  be **any** event. The **conditional PMF** of  $X$  given  $A$  is defined as

$$P_{X|A}(x_i) = P(X = x_i | A)$$

## Conditional PMF and Conditional CDF...

### Definition of Conditional PMF and Conditional CDF

Let  $X$  be a discrete random variable and  $A$  be any event. The conditional PMF of  $X$  given  $A$  is defined as

$$\begin{aligned} P_{X|A}(x_i) &= P(X = x_i | A) \\ &= \frac{P(X = x_i \text{ and } A)}{P(A)}, \quad \text{for any } x_i \in R_X \end{aligned}$$

## Conditional PMF and Conditional CDF...

### Definition of Conditional PMF and Conditional CDF

Let  $X$  be a discrete random variable and  $A$  be any event. The conditional PMF of  $X$  given  $A$  is defined as

$$\begin{aligned} P_{X|A}(x_i) &= P(X = x_i | A) \\ &= \frac{P(X = x_i \text{ and } A)}{P(A)}, \quad \text{for any } x_i \in Rx \end{aligned}$$

The conditional CDF of  $X$  is given by

## Conditional PMF and Conditional CDF...

### Definition of Conditional PMF and Conditional CDF

Let  $X$  be a **discrete** random variable and  $A$  be **any** event. The **conditional PMF** of  $X$  given  $A$  is defined as

$$\begin{aligned} P_{X|A}(x_i) &= P(X = x_i | A) \\ &= \frac{P(X = x_i \text{ and } A)}{P(A)}, \quad \text{for any } x_i \in R_X \end{aligned}$$

The **conditional CDF** of  $X$  is given by

$$F_{X|A}(x) = P(X \leq x | A).$$

Conditional PMF of  $X$  given  $Y \dots$

Conditional PMF of  $X$  given  $Y \dots$

Conditional PMF of  $X$  and  $Y$

## Conditional PMF of $X$ given $Y$ ...

### Conditional PMF of $X$ and $Y$

For **discrete** random variables  $X$  and  $Y$ ,

## Conditional PMF of $X$ given $Y$ ...

### Conditional PMF of $X$ and $Y$

For **discrete** random variables  $X$  and  $Y$ , the **conditional PMFs** of  $X$  and  $Y$  is defined as follows

## Conditional PMF of $X$ given $Y$ ...

### Conditional PMF of $X$ and $Y$

For **discrete** random variables  $X$  and  $Y$ , the **conditional PMFs** of  $X$  and  $Y$  is defined as follows

$$P_{X|Y}(x_i, y_j) = \frac{P_{XY}(x_i, y_j)}{P_Y(y_j)}$$

Joint PMF

## Conditional PMF of $X$ given $Y$ ...

### Conditional PMF of $X$ and $Y$

For **discrete** random variables  $X$  and  $Y$ , the **conditional PMFs** of  $X$  and  $Y$  is defined as follows

$$P_{X|Y}(x_i, y_j) = \frac{P_{XY}(x_i, y_j)}{P_Y(y_j)}$$
$$P_{Y|X}(x_i, y_j) = \frac{P_{XY}(x_i, y_j)}{P_X(x_i)}$$

## Conditional PMF of $X$ given $Y$ ...

### Conditional PMF of $X$ and $Y$

For **discrete** random variables  $X$  and  $Y$ , the **conditional PMFs** of  $X$  and  $Y$  is defined as follows

$$P_{X|Y}(x_i, y_j) = \frac{P_{XY}(x_i, y_j)}{P_Y(y_j)}$$

$$P_{Y|X}(x_i, y_j) = \frac{P_{XY}(x_i, y_j)}{P_X(x_i)}$$

for **any**  $x_i \in R_X$  and  $y_j \in R_Y$ .

## Independent Random Variables...

## Independent Random Variables...

Definition of Independent Random Variables

## Independent Random Variables...

### Definition of Independent Random Variables

Let  $X, Y$  be two random variables, then they are **independent** if

## Independent Random Variables...

### Definition of Independent Random Variables

Let  $X, Y$  be two random variables, then they are **independent** if

$$P_{XY}(x, y) = P_X(x)P_Y(y) \quad \text{for all } x, y$$

## Independent Random Variables...

### Definition of Independent Random Variables

Let  $X, Y$  be two random variables, then they are **independent** if

$$P_{XY}(x, y) = P_X(x)P_Y(y) \quad \text{for all } x, y$$

In other words,  $X$  and  $Y$  are **independent** if

## Independent Random Variables...

### Definition of Independent Random Variables

Let  $X, Y$  be two random variables, then they are **independent** if

$$P_{XY}(x, y) = P_X(x)P_Y(y) \quad \text{for all } x, y$$

In other words,  $X$  and  $Y$  are **independent** if

$$\underline{F_{XY}(x, y)} = \underline{F_X(x)} \underline{F_Y(y)}, \quad \text{for all } x, y$$

- If  $X$  and  $Y$  are **independent** if

## Independent Random Variables...

### Definition of Independent Random Variables

Let  $X, Y$  be two random variables, then they are **independent** if

$$P_{XY}(x, y) = P_X(x)P_Y(y) \quad \text{for all } x, y$$

In other words,  $X$  and  $Y$  are **independent** if

$$F_{XY}(x, y) = F_X(x)F_Y(y), \quad \text{for all } x, y$$

- If  $X$  and  $Y$  are **independent** if

$$P_{X|Y}(x_i|y_j) = P(X = x_i | Y = y_j) = \frac{P_{XY}(x_i, y_j)}{P_Y(y_j)} = \frac{P_X(x_i)P_Y(y_j)}{P_Y(y_j)} = P_X(x_i)$$

## Example of Joint and Marginal PDF and Conditional PMF...

## Example of Joint and Marginal PDF and Conditional PMF...

### Example

Consider the set of points in set  $G$  defined as follows

## Example of Joint and Marginal PDF and Conditional PMF...

### Example

Consider the set of points in set  $G$  defined as follows

$$G = \{(x, y) \mid x, y \in \mathbb{Z}, \quad |x| + |y| \leq 2\}.$$

## Example of Joint and Marginal PDF and Conditional PMF...

### Example

Consider the set of points in set  $G$  defined as follows

$$G = \{(x, y) \mid x, y \in \mathbb{Z}, \quad |x| + |y| \leq 2\}.$$

If we pick a point  $(X, Y)$  from this grid at random,

## Example of Joint and Marginal PDF and Conditional PMF...

### Example

Consider the set of points in set  $G$  defined as follows

$$G = \{(x, y) \mid x, y \in \mathbb{Z}, |x| + |y| \leq 2\}.$$

If we pick a point  $(X, Y)$  from this grid at random, then the probability of choosing a point is  $1/13$ .

$$G = \{(0,0), (0,1), (1,1), (1,-1), \dots\}$$
$$|G| = 13 \quad (\text{Ex})$$

## Example of Joint and Marginal PDF and Conditional PMF...

### Example

Consider the set of points in set  $G$  defined as follows

$$G = \{(x, y) \mid x, y \in \mathbb{Z}, \quad |x| + |y| \leq 2\}.$$

If we pick a point  $(X, Y)$  from this grid at random, then the probability of choosing a point is  $1/13$ .

- 1 Find the joint and marginal PMFs of  $X$  and  $Y$ .

## Example of Joint and Marginal PDF and Conditional PMF...

### Example

Consider the set of points in set  $G$  defined as follows

$$G = \{(x, y) \mid x, y \in \mathbb{Z}, \quad |x| + |y| \leq 2\}.$$

If we pick a point  $(X, Y)$  from this grid at random, then the probability of choosing a point is  $1/13$ .

- 1 Find the joint and marginal PMFs of  $X$  and  $Y$ .
- 2 Find the conditional PMF of  $X$  given  $Y = 1$ .

## Example of Joint and Marginal PDF and Conditional PMF...

### Example

Consider the set of points in set  $G$  defined as follows

$$G = \{(x, y) \mid x, y \in \mathbb{Z}, |x| + |y| \leq 2\}.$$

If we pick a point  $(X, Y)$  from this grid at random, then the probability of choosing a point is  $1/13$ .

- 1 Find the joint and marginal PMFs of  $X$  and  $Y$ .
- 2 Find the conditional PMF of  $X$  given  $Y = 1$ .
- 3 Are  $X$  and  $Y$  independent?

① Joint PMF

$$P_{XY}(x, y) = \begin{cases} \frac{1}{13}, & (x, y) \in G \\ 0, & \text{otherwise} \end{cases}$$

$$P_Y = \sum_{x_i} P_{XY}(x_i, y), \quad P_X = \sum_{y_i} P_{XY}(x, y_i)$$

Answer to previous problem...

$$P_X(-2) = P_{XY}(-2, 0) = \gamma_B$$

$$\begin{aligned} P_X(-1) &= P_{XY}(-1, -1) + P_{XY}(-1, 0) \\ &\quad + P_{XY}(0, -1) = \frac{3}{13} \\ P_X(0) &= - - - \\ &\quad - - - \end{aligned}$$

$$\begin{aligned} P_Y(0) &= P_{XY}(0, 0), P_{XY}(0, -1) + P_{XY}(0, 1) \\ P_X(0) &= P_{XY}(0, 0) + P_{XY}(0, -1) + P_{XY}(0, 1) \\ &+ P_{XY}(1, 0) + P_{XY}(1, -1) = \frac{5}{13} \\ &\neq \frac{1}{13} = P_X(1) \end{aligned}$$

② Given  $Y = 1$   
possible  $X = -1, 0, 1$

$$\begin{aligned} P_{XY}(i, 1) &= \frac{P_{XY}(i, 1)}{P_Y(1)} \\ &= \frac{\gamma_B}{\frac{3}{13}} \quad [P_Y(1) = P_X(-1)] \end{aligned}$$

$$\begin{aligned} P_{XY}(i|1) &= \frac{\gamma_B}{\frac{3}{13}} \\ &= \frac{\gamma_B}{3/13} \quad \text{for } i = -1, 0, 1 \\ &\text{otherwise } 0 \end{aligned}$$

Answer to previous problem...

③ Are  $X, Y$  ind?

$$P_{X|Y}(i, i) = P_X(i)$$

Not true

## Definition of Conditional Expectation...

## Definition of Conditional Expectation...

### Definition of Conditional Expectation

Let  $A$  be any event.

## Definition of Conditional Expectation...

### Definition of Conditional Expectation

Let  $A$  be any event. Let  $X$  and  $Y$  be two random variables with ranges  $R_X$  and  $R_Y$  respectively.

## Definition of Conditional Expectation...

### Definition of Conditional Expectation

Let  $A$  be any event. Let  $X$  and  $Y$  be two random variables with ranges  $R_X$  and  $R_Y$  respectively. Then the **conditional expectations** are defined as follows

## Definition of Conditional Expectation...

### Definition of Conditional Expectation

Let  $A$  be any event. Let  $X$  and  $Y$  be two random variables with ranges  $R_X$  and  $R_Y$  respectively. Then the **conditional expectations** are defined as follows

$$E[X | A] = \sum_{x_i \in R_X} x_i P_{X|A}(x_i)$$

$$E[X | Y = y_j] = \sum_{x_i \in R_X} x_i P_{X|Y}(x_i | y_j)$$

## Example of Conditional Expectation

## Example of Conditional Expectation

### Example

Consider the set of points in set  $G$  defined as follows

## Example of Conditional Expectation

### Example

Consider the set of points in set  $G$  defined as follows

$$G = \{(x, y) \mid x, y \in \mathbb{Z}, \quad |x| + |y| \leq 2\}.$$

## Example of Conditional Expectation

### Example

Consider the set of points in set  $G$  defined as follows

$$G = \{(x, y) \mid x, y \in \mathbb{Z}, \quad |x| + |y| \leq 2\}.$$

If we pick a point  $(X, Y)$  from this grid at random,

## Example of Conditional Expectation

### Example

Consider the set of points in set  $G$  defined as follows

$$G = \{(x, y) \mid x, y \in \mathbb{Z}, \quad |x| + |y| \leq 2\}.$$

If we pick a point  $(X, Y)$  from this grid at random, then the probability of choosing a point is  $1/13$ .

## Example of Conditional Expectation

When  $\gamma=1$ , possible for  
 $x = -1, \underline{0}, 1$

### Example

Consider the set of points in set  $G$  defined as follows

$$G = \{(x, y) \mid x, y \in \mathbb{Z}, |x| + |y| \leq 2\}.$$

If we pick a point  $(X, Y)$  from this grid at random, then the probability of choosing a point is  $1/13$ .

Find  $E[X \mid Y = 1]$

we have from before

$$P_{X|Y=1} = \begin{cases} \frac{1}{3}, & X = -1, 0, 1 \\ 0, & \text{otherwise} \end{cases}$$

$$\begin{aligned} \Rightarrow E[X \mid Y=1] &= (-1) P_{X|Y=1}(-1) + 0 P_{X|Y=1}(0) + 1 \cdot P_{X|Y=1}(1) \\ &= -1 \cdot \frac{1}{3} + 0 \cdot \frac{1}{3} + 1 \cdot \frac{1}{3} = 0 \end{aligned}$$

## Example of Conditional Expectation

$$P(-2) = \frac{13}{8} P(X=-2|A)$$

Example

$$\text{If } -1 < Y < 2 \Rightarrow X = -2, -1, 0 \quad = \frac{13}{8} P(-2|A) = \frac{13}{8} \cdot \frac{1}{13} = \underline{\underline{\frac{1}{8}}}$$

Consider the set of points in set  $G$  defined as follows

$$G = \{(x, y) \mid x, y \in \mathbb{Z}, |x| + |y| \leq 2\}$$

If we pick a point  $(X, Y)$  from this grid at random, then the probability of choosing a point is  $1/13$ .

1 Find  $E[X \mid Y = 1]$

2 Find  $E[X \mid -1 < Y < 2]$   $\leftarrow P_{X|Y}$   
 $-1 < Y < 2 \Rightarrow Y = 0, 1, Y \in \{0, 1\} = A$

$$P(-1 < Y < 2) = P(A) = P_y(0) + P_y(1) = \frac{5}{13} + \frac{3}{13} = \frac{8}{13}$$

$$P_{X|A}(k) = \frac{P(X=k, A)}{P(A)} = \frac{13}{8} \frac{P(X=k, A)}{13} \rightarrow$$

## Example of Conditional Expectation

### Example

Consider the set of points in set  $G$  defined as follows

$$G = \{(x, y) \mid x, y \in \mathbb{Z}, \quad |x| + |y| \leq 2\}.$$

If we pick a point  $(X, Y)$  from this grid at random, then the probability of choosing a point is  $1/13$ .

- 1 Find  $E[X \mid Y = 1]$
- 2 Find  $E[X \mid -1 < Y < 2]$  
- 3 Find  $E[|X| \mid -1 < Y < 2]$  

Answer to previous problem...

$$X = \{ -2, 0, 1, 2 \}$$

$$P_{X|A}(-1) = \frac{13}{16} P(X=-1, A)$$

$$= \frac{13}{16} P_{XY}(-1, 0) + \frac{13}{16} P_{XY}(-1, 1)$$

$$= \frac{13}{16} \cdot \frac{1}{5} + \frac{13}{16} \cdot \frac{1}{3}$$
$$= \frac{1}{4}$$

Similarly

$$P_{X|A}(0) = \frac{13}{16} P_{XY}(0, 0) + \frac{13}{16} P_{XY}(0, 1)$$
$$= \frac{1}{4}$$

$$P_{X|A}(2) = \frac{1}{8}$$

$$E[X|A]$$

$$= \sum_i x_i P_{X|A}(x_i)$$

$$= -2 \cdot \frac{1}{8} + (-1) \cdot \frac{1}{4} + 0 \cdot \frac{1}{4}$$

$$+ 1 \cdot \frac{1}{4} + 2 \cdot \frac{1}{8}$$
$$= -$$

Answer to previous problem...



## Law of Total Probability...

## Law of Total Probability...

Law of Total Probability

## Law of Total Probability...

### Law of Total Probability

- Law of Total Probability

## Law of Total Probability...

### Law of Total Probability

- Law of Total Probability

$$\underbrace{P(X \in A)}_{y_j \in R_Y} = \sum \underbrace{P(X \in A | Y = y_j) P_Y(y_j)}_{\text{for any set } A},$$

## Law of Total Probability...

### Law of Total Probability

- Law of Total Probability

$$P(X \in A) = \sum_{y_j \in R_Y} P(X \in A \mid Y = y_j) P_Y(y_j), \quad \text{for any set } A$$

$y_j \in R_Y$

- Law of Total Expectation:

## Law of Total Probability...

### Law of Total Probability

- Law of Total Probability

$$P(X \in A) = \sum_{y_j \in R_Y} P(X \in A \mid Y = y_j) P_Y(y_j), \quad \text{for any set } A$$

- Law of Total Expectation:

1 If  $B_1, B_2, \dots$  is a partition of sample space  $S$

## Law of Total Probability...

### Law of Total Probability

- Law of Total Probability

$$P(X \in A) = \sum_{y_j \in R_Y} P(X \in A | Y = y_j) P_Y(y_j), \quad \text{for any set } A$$

- Law of Total Expectation:

1 If  $B_1, B_2, \dots$  is a partition of sample space  $S$

$$E[X] = \sum_i E[X | B_i] P(B_i)$$

## Law of Total Probability...

### Law of Total Probability

- Law of Total Probability

$$P(X \in A) = \sum_{y_j \in R_Y} P(X \in A \mid Y = y_j) P_Y(y_j), \quad \text{for any set } A$$

- Law of Total Expectation:

- 1 If  $B_1, B_2, \dots$  is a partition of sample space  $S$

$$E[X] = \sum_i E[X \mid B_i] P(B_i)$$

- 2 For a RV  $X$  and a discrete RV  $Y$

## Law of Total Probability...


### Law of Total Probability

- Law of Total Probability

$$P(X \in A) = \sum_{y_j \in R_Y} P(X \in A | Y = y_j) P_Y(y_j), \quad \text{for any set } A$$

- Law of Total Expectation:

1 If  $B_1, B_2, \dots$  is a partition of sample space  $S$


$$E[X] = \sum_i E[X | B_i] P(B_i)$$

2 For a RV  $X$  and a discrete RV  $Y$

$$E[Y] = \sum_{y_j \in R_Y} E[X | Y = y_j] P_Y(y_j)$$

## Solved Example

## Solved Example

X

$$\sum x_i p(x_i)$$

### Solved Example 1

Let  $X \sim \text{Geometric}(p)$ . Find  $E[X]$ . [Hint: condition on first coin toss.]

Geometric : number of times I toss to get 1st head.  
 $p(H) = p$ ,  $p(T) = 1-p$ .  
 $X$  = total no. of tosses to get 1st H.

Two outcomes :  $\{H\} \cup \{T\}$

Using the law of total expectation.

$$\begin{aligned} E[X] &= E[X|H] p(H) + E[X|T] p(T) \\ &= p E[X|H] + (1-p) E[X|T] \\ &= p \cdot 1 + (1-p) (1 + E[X]) \end{aligned}$$

$$\begin{aligned} E[X|T] &= 1 + E[X] \\ 1 + E[X] &= 1 + E[X] \\ \text{Solve for } E[X]. & \\ 1/p & \quad (\text{dual}) \end{aligned}$$

## Solved Example

## Solved Example

### Solved Example 2

Number of customers  $N$  visiting a fast food restaurant follows Poisson distribution  $N \sim \text{Poisson}(\lambda)$ .

## Solved Example

### Solved Example 2

Number of customers  $N$  visiting a fast food restaurant follows Poisson distribution  $N \sim \text{Poisson}(\lambda)$ . Each customer arriving in this restaurant purchases a drink with probability  $p$ ,

## Solved Example

### Solved Example 2


Number of customers  $N$  visiting a fast food restaurant follows Poisson distribution  $N \sim \text{Poisson}(\lambda)$ . Each customer arriving in this restaurant purchases a drink with probability  $p$ , which is independent from other customers.

## Solved Example



## Solved Example 2

Number of customers  $N$  visiting a fast food restaurant follows Poisson distribution  $N \sim \text{Poisson}(\lambda)$ . Each customer arriving in this restaurant purchases a drink with probability  $p$ , which is independent from other customers. What is the average number of customers who purchase drinks?



Answer to previous problem...

Given  $N=n$ ,  $X$  is a sum of  $n$  independent Bernoulli( $p$ ), which means

$$X|N=n \sim \text{Binomial}(n, p)$$

$$P_{X|N}(k|n) = \binom{n}{k} p^k (1-p)^{n-k}.$$

$$E[X|N=n] = np.$$

We want:  $E[X]$

= exp. no. of customers purchasing drinks.

$$\begin{aligned} & L.O.T.E \sum_{n=0}^{\infty} E[X|N=n] P_N(n) \\ &= \sum_{n=0}^{\infty} np \cdot P_N(n) \\ &= p \sum_{n=0}^{\infty} n P_N(n) \\ &= p \lambda \quad \begin{cases} \text{PMF} \\ \text{Pois.} \end{cases} \\ & \uparrow \quad \text{Exp. of Pois.} \end{aligned}$$

## PMF and Expectation of Two Random Variables...

## PMF and Expectation of Two Random Variables...

### PMF and Expectations of Two Random Variables

- Let  $X, Y$  be two RVs and suppose  $Z = \underline{g(X, Y)}$ ,  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ .

## PMF and Expectation of Two Random Variables...

### PMF and Expectations of Two Random Variables

- Let  $X, Y$  be two RVs and suppose  $Z = g(X, Y), g : \mathbb{R}^2 \rightarrow \mathbb{R}$ .  
Then the PMF of  $Z$  is

## PMF and Expectation of Two Random Variables...

### PMF and Expectations of Two Random Variables

- Let  $X, Y$  be two RVs and suppose  $Z = g(X, Y), g : \mathbb{R}^2 \rightarrow \mathbb{R}$ .  
Then the PMF of  $Z$  is

$$P_Z(z) = P(g(X, Y) = z) = \sum_{(x_i, y_j) \in A_z} P_{XY}(x_i, y_j),$$

## PMF and Expectation of Two Random Variables...

### PMF and Expectations of Two Random Variables

- Let  $X, Y$  be two RVs and suppose  $Z = g(X, Y), g : \mathbb{R}^2 \rightarrow \mathbb{R}$ .  
Then the PMF of  $Z$  is

$$P_Z(z) = P(g(X, Y) = z) = \sum_{(x_i, y_j) \in A_z} P_{XY}(x_i, y_j),$$

where  $A_z = \{(x_i, y_j) \in R_{XY} : g(x_i, y_j) = z\}$

more than one  
for given  $z$   
if  $g$  is not 1-1

## PMF and Expectation of Two Random Variables...

### PMF and Expectations of Two Random Variables

- Let  $X, Y$  be two RVs and suppose  $Z = g(X, Y), g : \mathbb{R}^2 \rightarrow \mathbb{R}$ .  
Then the PMF of  $Z$  is

$$P_Z(z) = P(g(X, Y) = z) = \sum_{(x_i, y_j) \in A_z} P_{XY}(x_i, y_j),$$

where  $A_z = \{(x_i, y_j) \in R_{XY} : g(x_i, y_j) = z\}$

- The expectation is given as follows

## PMF and Expectation of Two Random Variables...

### PMF and Expectations of Two Random Variables

- Let  $X, Y$  be two RVs and suppose  $Z = g(X, Y), g : \mathbb{R}^2 \rightarrow \mathbb{R}$ .  
Then the PMF of  $Z$  is

$$P_Z(z) = P(g(X, Y) = z) = \sum_{(x_i, y_j) \in A_z} P_{XY}(x_i, y_j),$$

$g(x) P(\tau)$

where  $A_z = \{(x_i, y_j) \in R_{XY} : g(x_i, y_j) = z\}$

- The expectation is given as follows

$$E[g(X, Y)] = \sum_{(x_i, y_j) \in R_{XY}} \underbrace{g(x_i, y_j)}_{g(x) P(\tau)} P_{XY}(x_i, y_j)$$

## Linearity of Expectation for Two Random Variable...

## Linearity of Expectation for Two Random Variable...

### Linearity of Expectation for Two RV

Let  $X, Y$  be two discrete RVs. Then  $E[X + Y] = E[X] + E[Y]$ .

$$\sum$$

## PMF of Difference of Two Geometric Distributions...

## PMF of Difference of Two Geometric Distributions...

PMF of Difference

Let  $X, Y \sim \text{Geometric}(p)$  be two random variables. Let  $Z = X - Y$ . Find the PMF of  $Z$ .

$$R_X, R_Y = \{1, 2, 3, \dots\} = \mathbb{N}$$

$$\Rightarrow R_Z = \{\dots, -2, -1, 0, 1, 2, \dots\} = \mathbb{Z}.$$

Since  $X, Y$  Geometric

$$P_X(k) = P_Y(k) = p^k q^{k-1}, \quad k=1, 2, 3, \dots$$

$$P_Z(k) = P(Z=k) = P(X-Y=k) = P(X=\underline{Y+k})$$

$\xrightarrow{\text{L.O.T.P}}$

$$= \sum_{j=1}^{\infty} P(X=\underline{Y+k} | Y=j) P(Y=j)$$

$\circlearrowleft + Y \text{ ind}$

$$= \sum_{j=1}^{\infty} P(X=\underline{j+k}) P(Y=j)$$

Answer to previous problem...


$$P_2(k) = \sum_{j=1}^{\infty} p^j q^{j+k-1} \cdot p^{j-1}$$

$$\frac{k \gamma, 0}{\longrightarrow}$$

For  $k < 0$

$$P_2(k) = \sum_{j=-k+1}^{\infty} p^j q^{j+k-1} \cdot p^{j-1}$$

Answer to previous problem...



## Conditional Expectation Revisited...

## Conditional Expectation Revisited...

- We recall that the conditional expectation of  $X$  given  $Y = y$  is

## Conditional Expectation Revisited...

- We recall that the **conditional expectation** of  $X$  given  $Y = y$  is

$$E[X | Y = y] = \sum_{x_i \in R_X} x_i P_{X|Y}(x_i | y)$$

## Conditional Expectation Revisited...

- We recall that the conditional expectation of  $X$  given  $Y = y$  is

$$E[X | Y = y] = \sum_{x_i \in R_X} x_i P_{X|Y}(x_i | y)$$

- Here  $E[X | Y = y]$  is a function of  $y$ . That is  $g(y) = E[X | Y = y]$

## Conditional Expectation Revisited...

- We recall that the conditional expectation of  $X$  given  $Y = y$  is

$$E[X | Y = y] = \sum_{x_i \in R_X} x_i P_{X|Y}(x_i | y)$$

- Here  $E[X | Y = y]$  is a function of  $y$ . That is  $g(y) = E[X | Y = y]$
- We can think of  $g(Y) = E[X | Y]$  as a random variable

## Conditional Expectation Revisited...

- We recall that the **conditional expectation** of  $X$  given  $Y = y$  is

$$E[X | Y = y] = \sum_{x_i \in R_X} x_i P_{X|Y}(x_i | y)$$

- Here  $E[X | Y = y]$  is a **function of  $y$** . That is  $g(y) = E[X | Y = y]$
- We can think of  $g(Y) = E[X | Y]$  as a random variable
- If  $Y$  is a RV with range  $R_Y = \{y_1, y_2, \dots\}$ , then

## Conditional Expectation Revisited...

- We recall that the **conditional expectation** of  $X$  given  $Y = y$  is

$$E[X | Y = y] = \sum_{x_i \in R_X} x_i P_{X|Y}(x_i | y)$$

- Here  $E[X | Y = y]$  is a **function of  $y$** . That is  $g(y) = E[X | Y = y]$
- We can think of  $g(Y) = E[X | Y]$  as a random variable
- If  $Y$  is a RV with range  $R_Y = \{y_1, y_2, \dots\}$ , then

$$E[X | Y] = \begin{cases} E[X | Y = y_1] & \text{with probability } P(Y = y_1) \\ E[X | Y = y_2] & \text{with probability } P(Y = y_2) \\ \vdots & \vdots \end{cases}$$

## Conditional Expectation Revisited...

- We recall that the **conditional expectation** of  $X$  given  $Y = y$  is

$$E[X | Y = y] = \sum_{x_i \in R_X} x_i P_{X|Y}(x_i | y)$$

- Here  $E[X | Y = y]$  is a **function of  $y$** . That is  $g(y) = E[X | Y = y]$
- We can think of  $g(Y) = E[X | Y]$  as a random variable
- If  $Y$  is a RV with range  $R_Y = \{y_1, y_2, \dots\}$ , then

$$E[X | Y] = \begin{cases} E[X | Y = y_1] & \text{with probability } P(Y = y_1) \\ E[X | Y = y_2] & \text{with probability } P(Y = y_2) \\ \vdots & \vdots \end{cases}$$

- For an example, let  $X = aY + b$ . Then  $E[X | Y = y] = E[aY + b | Y = y] = ay + b$

## Conditional Expectation Revisited...

- We recall that the conditional expectation of  $X$  given  $Y = y$  is

$$\underline{E[X | Y = y]} = \sum_{x_i \in R_X} x_i \underline{P_{X|Y}(x_i | y)}$$

- Here  $E[X | Y = y]$  is a function of  $y$ . That is  $g(y) = E[X | Y = y]$
- We can think of  $g(Y) = \underline{\underline{E[X | Y]}}$  as a random variable
- If  $Y$  is a RV with range  $R_Y = \{y_1, y_2, \dots\}$ , then

$$E[X | Y] = \begin{cases} E[X | Y = y_1] & \text{with probability } P(Y = y_1) \\ E[X | Y = y_2] & \text{with probability } P(Y = y_2) \\ \vdots & \vdots \end{cases}$$

- For an example, let  $\underline{X = aY + b}$ . Then  $\underline{E[X | Y = y]} = E[aY + b | Y = y] = \boxed{ay + b}$   
 $g(y) = \boxed{ay + b}, \quad E[X | Y] = aY + b$

## Conditional Expectation Revisited...

- We recall that the **conditional expectation** of  $X$  given  $Y = y$  is

$$E[X | Y = y] = \sum_{x_i \in R_X} x_i P_{X|Y}(x_i | y)$$



- Here  $E[X | Y = y]$  is a **function of  $y$** . That is  $g(y) = E[X | Y = y]$
- We can think of  $g(Y) = E[X | Y]$  as a random variable
- If  $Y$  is a RV with range  $R_Y = \{y_1, y_2, \dots\}$ , then

$X$ ,  
 $x$ ,  
 $x$ ,  
.

$$E[X | Y] = \begin{cases} E[X | Y = y_1] & \text{with probability } P(Y = y_1) \\ E[X | Y = y_2] & \text{with probability } P(Y = y_2) \\ \vdots & \vdots \end{cases}$$

- For an example, let  $X = aY + b$ . Then  $E[X | Y = y] = E[aY + b | Y = y] = ay + b$   
 $g(y) = ay + b$ ,  $E[X|Y] = aY + b$
- Since  $E[X | Y]$  is a **RV**, we can find its PMF, CDF, Variance, etc

## Solved Example on Conditional Expectation as a Random Variable...

## Solved Example on Conditional Expectation as a Random Variable...

### Solved Example

Let  $X, Y$  be RV with joint PMF given as follows

## Solved Example on Conditional Expectation as a Random Variable...

### Solved Example

Let  $X, Y$  be RV with joint PMF given as follows

	$Y = 0$	$Y = 1$
$X = 0$	$\frac{1}{5}$	$\frac{2}{5}$
$X = 1$	$\frac{2}{5}$	0

## Solved Example on Conditional Expectation as a Random Variable...

### Solved Example

Let  $X, Y$  be RV with joint PMF given as follows

	$Y = 0$	$Y = 1$
$X = 0$	$\frac{1}{5}$	$\frac{2}{5}$
$X = 1$	$\frac{2}{5}$	0

Let  $Z = E[X | Y]$ .

## Solved Example on Conditional Expectation as a Random Variable...

### Solved Example

Let  $X, Y$  be RV with joint PMF given as follows

	$Y = 0$	$Y = 1$
$X = 0$	$\frac{1}{5}$	$\frac{2}{5}$
$X = 1$	$\frac{2}{5}$	0

Let  $Z = E[X | Y]$ .

- Find the Marginal PMFs of  $X$  and  $Y$ .

## Solved Example on Conditional Expectation as a Random Variable...

### Solved Example

Let  $X, Y$  be RV with joint PMF given as follows

	$Y = 0$	$Y = 1$
$X = 0$	$\frac{1}{5}$	$\frac{2}{5}$
$X = 1$	$\frac{2}{5}$	0

Let  $Z = E[X | Y]$ .

- 1 Find the Marginal PMFs of  $X$  and  $Y$ .
- 2 Find the conditional PMF of  $X$  given  $Y=0$  and  $Y=1$ , i.e., find  $P_{X|Y}(x|0)$  and  $P_{X|Y}(x|1)$

## Solved Example on Conditional Expectation as a Random Variable...

### Solved Example

Let  $X, Y$  be RV with joint PMF given as follows

	$Y = 0$	$Y = 1$
$X = 0$	$\frac{1}{5}$	$\frac{2}{5}$
$X = 1$	$\frac{2}{5}$	0

Let  $Z = E[X | Y]$ .

- 1 Find the Marginal PMFs of  $X$  and  $Y$ .
- 2 Find the conditional PMF of  $X$  given  $Y=0$  and  $Y=1$ , i.e., find  $P_{X|Y}(x|0)$  and  $P_{X|Y}(x|1)$
- 3 Find the PMF of  $Z$

## Solved Example on Conditional Expectation as a Random Variable...

### Solved Example

Let  $X, Y$  be RV with joint PMF given as follows

	$Y = 0$	$Y = 1$
$X = 0$	$\frac{1}{5}$	$\frac{2}{5}$
$X = 1$	$\frac{2}{5}$	0

Let  $Z = E[X | Y]$ .

- 1 Find the Marginal PMFs of  $X$  and  $Y$ .
- 2 Find the conditional PMF of  $X$  given  $Y=0$  and  $Y=1$ , i.e., find  $P_{X|Y}(x|0)$  and  $P_{X|Y}(x|1)$
- 3 Find the PMF of  $Z$
- 4 Find  $E[Z]$ , and check that  $E[Z] = E[X]$

## Solved Example on Conditional Expectation as a Random Variable...

### Solved Example

Let  $X, Y$  be RV with joint PMF given as follows

	$Y = 0$	$Y = 1$
$X = 0$	$\frac{1}{5}$	$\frac{2}{5}$
$X = 1$	$\frac{2}{5}$	0

Let  $Z = E[X | Y]$ .

- 1 Find the Marginal PMFs of  $X$  and  $Y$ .
- 2 Find the conditional PMF of  $X$  given  $Y=0$  and  $Y=1$ , i.e., find  $P_{X|Y}(x|0)$  and  $P_{X|Y}(x|1)$
- 3 Find the PMF of  $Z$
- 4 Find  $E[Z]$ , and check that  $E[Z] = E[X]$
- 5 Find  $\text{Var}(Z)$