

Probability and Statistics

UG2, Core course, IIIT,H

Pawan Kumar

IIIT, Hyderabad

August 27, 2021

- ① Digress: Game Theory
- ② Random Walks
- ③ Conditional Probability, Bayes Theorem


- ④ Define Conditional Probability and Chain Rule
- ⑤ The Monty Hall Problem
- ⑥ Independence
- ⑦ Conditional Independence

0 Computing probabilities...

| 2

Experiment: Throw two dice.

0 Computing probabilities...



| 2

Experiment: Throw two dice.

Event: Bigger number on red than blue.

0 Computing probabilities...

| 2

Experiment: Throw two dice.

Event: Bigger number on red than blue.

Quiz: What is the probability of this event?

0 Computing probabilities...

| 2

Experiment: Throw two dice.

Event: Bigger number on red than blue.

Quiz: What is the probability of this event?

11	12	13	14	15	16
21	22	23	24	25	26
31	32	33	34	35	36
41	42	43	44	45	46
51	52	53	54	55	56
61	62	63	64	65	66

0 Computing probabilities...

| 2

Experiment: Throw two dice.

Event: Bigger number on red than blue.

Quiz: What is the probability of this event?

- Probability space: all outcomes

11	12	13	14	15	16
21	22	23	24	25	26
31	32	33	34	35	36
41	42	43	44	45	46
51	52	53	54	55	56
61	62	63	64	65	66

0 Computing probabilities...

| 2

Experiment: Throw two dice.

Event: Bigger number on red than blue.

Quiz: What is the probability of this event?

- Probability space: all outcomes
- Event: some favourable outcomes

11	12	13	14	15	16
21	22	23	24	25	26
31	32	33	34	35	36
41	42	43	44	45	46
51	52	53	54	55	56
61	62	63	64	65	66

0 Computing probabilities...

| 2

Experiment: Throw two dice.

Event: Bigger number on red than blue.

Quiz: What is the probability of this event?

11	12	13	14	15	16
21	22	23	24	25	26
31	32	33	34	35	36
41	42	43	44	45	46
51	52	53	54	55	56
61	62	63	64	65	66

- Probability space: all outcomes
- Event: some favourable outcomes
- Probability: (#favorable)/36

11	12	13	14	15	16
21	22	23	24	25	26
31	32	33	34	35	36
41	42	43	44	45	46
51	52	53	54	55	56
61	62	63	64	65	66

0 Computing probabilities...

| 2

Experiment: Throw two dice.

Event: Bigger number on red than blue.

Quiz: What is the probability of this event?

11	12	13	14	15	16
21	22	23	24	25	26
31	32	33	34	35	36
41	42	43	44	45	46
51	52	53	54	55	56
61	62	63	64	65	66

- Probability space: all outcomes
- Event: some favourable outcomes
- Probability: (#favorable)/36

11	12	13	14	15	16
21	22	23	24	25	26
31	32	33	34	35	36
41	42	43	44	45	46
51	52	53	54	55	56
61	62	63	64	65	66

Hence, probability, $p = \frac{15}{36}$

Quiz

- Experiment: Toss a coin n times

Quiz

- Experiment: Toss a coin n times
- Outcome: sequence of n bits

Quiz



- Experiment: Toss a coin n times
- Outcome: sequence of n bits
- Total number of outcomes: 2^n outcomes

Quiz

- **Experiment:** Toss a coin n times
- **Outcome:** sequence of n bits
- **Total number of outcomes:** 2^n outcomes
- **Assumption:** equiprobable


Quiz

- Experiment: Toss a coin n times
- Outcome: sequence of n bits
- Total number of outcomes: 2^n outcomes
- Assumption: equiprobable
- Probability of all heads: $1/2^n$

{H H H}

Quiz

- Experiment: Toss a coin n times
- Outcome: sequence of n bits
- Total number of outcomes: 2^n outcomes
- Assumption: equiprobable
- Probability of all heads: $1/2^n$
- Consider an event of "first bit" = "last bit"



Quiz

- **Experiment:** Toss a coin n times
- **Outcome:** sequence of n bits
- **Total number of outcomes:** 2^n outcomes
- **Assumption:** equiprobable
- **Probability of all heads:** $1/2^n$
- Consider an event of “first bit” = “last bit”
- What is the **probability** of the above event?

0 Answer to Quiz...


Answer to Quiz:

0 Answer to Quiz...

| 4

Answer to Quiz:

- Number of possible cases:



Answer to Quiz:

- Number of possible cases:

 $H * \dots * H$ $T * \dots * H$ $H * \dots * T$ $T * \dots * T$

- Observation: In half of the cases, first bit does not equal last bit

Answer to Quiz:

- Number of possible cases:

 $H * \dots * H$ $T * \dots * H$ $H * \dots * T$ $T * \dots * T$

- Observation: In half of the cases, first bit does not equal last bit
- Hence, probability of the event is 1/2

0 Remark on Counting. Does probability boils down to counting always?

0 Remark on Counting. Does probability boils down to counting always?

| 5

- It is true that in many cases probability boils down to counting

0 Remark on Counting. Does probability boils down to counting always?

| 5

- It is true that in many cases probability boils down to counting
- However, there are **non-uniform distributions** 

0 Remark on Counting. Does probability boils down to counting always?

| 5

- It is true that in many cases probability boils down to counting
- However, there are **non-uniform** distributions
- We also study **continuous** distributions

0 Remark on Counting. Does probability boils down to counting always?


| 5

- It is true that in many cases probability boils down to counting
- However, there are **non-uniform** distributions
- We also study **continuous** distributions
- In the continuous case, we need to consider some other measure

0 Remark on Counting. Does probability boils down to counting always?

| 5

- It is true that in many cases probability boils down to counting
- However, there are **non-uniform** distributions
- We also study **continuous distributions**
- In the continuous case, we need to consider some other measure
- What is the probability that a number 1.5 is picked in $(1, 2]$?



0 Remark on Counting. Does probability boils down to counting always?

| 5

- It is true that in many cases probability boils down to counting
- However, there are **non-uniform** distributions
- We also study **continuous** distributions
- In the continuous case, we need to consider some other measure
- What is the probability that a number 1.5 is picked in $[1, 2]$? Does this make sense?


0 Probability, Galton Board, and Pascal Triangle

| 6

Movie of Galton board here!


0 Probability, Galton Board, and Pascal Triangle

| 7



0 Probability, Galton Board, and Pascal Triangle


| 7



- This is called galton board, Sir Francis Galton (1822-1911)

0 Probability, Galton Board, and Pascal Triangle


| 7



- This is called galton board, Sir Francis Galton (1822-1911)
- The set of red marbles are allowed to fall

0 Probability, Galton Board, and Pascal Triangle


| 7



- This is called galton board, Sir Francis Galton (1822-1911)
- The set of red marbles are allowed to fall
- We observe the distribution of the marbles after the fall

0 Probability, Galton Board, and Pascal Triangle


| 7



- This is called galton board, Sir Francis Galton (1822-1911)
- The set of red marbles are allowed to fall
- We observe the distribution of the marbles after the fall
- Quiz: Why do we have more marbles in the middle?

0 Probability, Galton Board, and Pascal Triangle


| 7



- This is called galton board, Sir Francis Galton (1822-1911)
- The set of red marbles are allowed to fall
- We observe the distribution of the marbles after the fall
- Quiz: Why do we have more marbles in the middle?
- Does probability have any role in this distribution?

0 Probability, Galton Board, and Pascal Triangle


| 7



- This is called galton board, Sir Francis Galton (1822-1911)
- The set of red marbles are allowed to fall
- We observe the distribution of the marbles after the fall
- Quiz: Why do we have more marbles in the middle?
- Does probability have any role in this distribution?
- Can we prove that the distribution will look like this mathematically?

0 Probability, Galton Board, and Pascal Triangle


| 7



- This is called galton board, Sir Francis Galton (1822-1911)
- The set of red marbles are allowed to fall
- We observe the distribution of the marbles after the fall
- Quiz: Why do we have more marbles in the middle?
- Does probability have any role in this distribution?
- Can we prove that the distribution will look like this mathematically?
- Does Pascal triangle has anything to do with this?

0 Probability, Galton Board, and Pascal Triangle


| 7



- This is called galton board, Sir Francis Galton (1822-1911)
- The set of red marbles are allowed to fall
- We observe the distribution of the marbles after the fall
- Quiz: Why do we have more marbles in the middle?
- Does probability have any role in this distribution?
- Can we prove that the distribution will look like this mathematically?
- Does Pascal triangle has anything to do with this?
- Let us analyze this in detail ...

0 Probability, Galton Board, and Pascal Triangle


| 8




0 Probability, Galton Board, and Pascal Triangle


| 8

- Assume that at each level the beams are splitted evenly





- Assume that at each level the beams are splitted evenly
- Probability that ball hits current beam is sum of probability of this ball hitting left and right beam above






- Assume that at each level the beams are splitted evenly
- Probability that ball hits current beam is sum of probability of this ball hitting left and right beam above
- If x is the top left probability, and y is the top right probability, then probability of hitting the current bin is $(x + y)/2$

0 Probability, Galton Board, and Pascal Triangle

| 8

- Assume that at each level the beams are splitted evenly
- Probability that ball hits current beam is sum of probability of this ball hitting left and right beam above
- If x is the top left probability, and y is the top right probability, then probability of hitting the current bin is $(x + y)/2$





- Assume that at each level the beams are splitted evenly
- Probability that ball hits current beam is sum of probability of this ball hitting left and right beam above
- If x is the top left probability, and y is the top right probability, then probability of hitting the current bin is $(x + y)/2$

		1		
		$1/2$	$1/2$	
	$1/4$	$2/4$	$1/4$	
$1/8$	$3/8$	$3/8$	$1/8$	

- The probability of current bin = $\binom{n}{k} / 2^n$

$$\binom{n}{k} / 2^n$$

0 Non equiprobable outcomes!

| 9

What is wrong with these dice?



0 Non equiprobable outcomes!

| 9

What is wrong with these dice?



- So far we assumed **equiprobable** outcomes

0 Non equiprobable outcomes!

| 9

What is wrong with these dice?



- So far we assumed **equiprobable** outcomes
- On the left, we have **skewed dice**, i.e., unfair dice

0 Non equiprobable outcomes!

| 9

What is wrong with these dice?



- So far we assumed **equiprobable** outcomes
- On the left, we have **skewed dice**, i.e., **unfair dice**
- Will our previous definition of calculating probability work?

0 Non equiprobable outcomes!

| 9

What is wrong with these dice?



$$p(s) = \underline{\underline{1}}$$

- So far we assumed **equiprobable** outcomes
- On the left, we have **skewed dice**, i.e., **unfair dice**
- Will our previous definition of calculating probability work?
- Unfortunately not, how to deal with skewed cases?

0 Non equiprobable outcomes!

| 10



0 Non equiprobable outcomes!

| 10



- Since the dice is skewed, frequency of outcomes are different

0 Non equiprobable outcomes!

| 10



- Since the dice is skewed, frequency of outcomes are different
 - That is, for example, number 4 may appear more frequently

0 Non equiprobable outcomes!

| 10




- Since the dice is skewed, frequency of outcomes are **different**
 - That is, for example, number 4 may appear more frequently
- Let p_i denote the probability of i th number

0 Non equiprobable outcomes!

| 10



- Since the dice is skewed, frequency of outcomes are **different**
 - That is, for example, number 4 may appear more frequently
- Let p_i denote the probability of i th number
- If we throw the dice long enough, the frequencies stabilise



0 Non equiprobable outcomes!

| 10



- Since the dice is skewed, frequency of outcomes are **different**
 - That is, for example, number 4 may appear more frequently
- Let p_i denote the probability of i th number
- If we throw the dice long enough, the frequencies **stabilise**
- For example, the probability of getting an even number =
 p_2 + p_4 + p_6

0 Non equiprobable outcomes!

| 10



- Since the dice is skewed, frequency of outcomes are **different**
 - That is, for example, number 4 may appear more frequently
- Let p_i denote the probability of i th number
- If we throw the dice long enough, the frequencies **stabilise**
- For example, the probability of getting an even number = $p_2 + p_4 + p_6$
- Also, we know that the sum of **all the probabilities sum to 1, even though the dice is skewed!**
 - That is: $p_1 + p_2 + \dots + p_6 = 1$

0 Example: Dice Throw and Venn Diagram

| 11

Experiment: Throw a fair dice once

0 Example: Dice Throw and Venn Diagram

| 11

Experiment: Throw a fair dice once


Sample Space: $S = \{1, 2, 3, 4, 5, 6\}$

0 Example: Dice Throw and Venn Diagram

| 11

Experiment: Throw a fair dice once

Sample Space: $S = \{1, 2, 3, 4, 5, 6\}$




- Consider event $\underline{A} = \{1, 2, 3\}$

0 Example: Dice Throw and Venn Diagram

| 11

Experiment: Throw a fair dice once

Sample Space: $S = \{1, 2, 3, 4, 5, 6\}$




- Consider event $A = \{1, 2, 3\}$
- Consider event $B = \{5, 6\}$

0 Example: Dice Throw and Venn Diagram

Experiment: Throw a fair dice once

Sample Space: $S = \{1, 2, 3, 4, 5, 6\}$




- Since the dice is fair, we have $p_1 + p_2 + \cdots + p_6 = 1$

- Consider event $A = \{1, 2, 3\}$
- Consider event $B = \{5, 6\}$

0 Example: Dice Throw and Venn Diagram

Experiment: Throw a fair dice once

Sample Space: $S = \{1, 2, 3, 4, 5, 6\}$



- Since the dice is fair, we have $p_1 + p_2 + \dots + p_6 = 1$
- $Pr(A) = p_1 + p_2 + p_3$


- Consider event $A = \{1, 2, 3\}$
- Consider event $B = \{5, 6\}$

0 Example: Dice Throw and Venn Diagram

| 11

Experiment: Throw a fair dice once

Sample Space: $S = \{1, 2, 3, 4, 5, 6\}$



- Since the dice is fair, we have $p_1 + p_2 + \cdots + p_6 = 1$
- $Pr(A) = p_1 + p_2 + p_3$
- $Pr(B) = p_5 + p_6$


- Consider event $A = \{1, 2, 3\}$
- Consider event $B = \{5, 6\}$

0 Example: Dice Throw and Venn Diagram

| 11

Experiment: Throw a fair dice once

Sample Space: $S = \{1, 2, 3, 4, 5, 6\}$



- Consider event $A = \{1, 2, 3\}$
- Consider event $B = \{5, 6\}$


- Since the dice is fair, we have $p_1 + p_2 + \dots + p_6 = 1$
- $Pr(A) = p_1 + p_2 + p_3$
- $Pr(B) = p_5 + p_6$
- Consider the event $\underbrace{A \text{ or } B}_{\text{event}} = \{1, 2, 3, 5, 6\}$

0 Example: Dice Throw and Venn Diagram

| 11

Experiment: Throw a fair dice once

Sample Space: $S = \{1, 2, 3, 4, 5, 6\}$



Axiom 3

- Since the dice is fair, we have $p_1 + p_2 + \dots + p_6 = 1$
- $Pr(A) = p_1 + p_2 + p_3$
- $Pr(B) = p_5 + p_6$
- Consider the event A or $B = \{1, 2, 3, 5, 6\}$
- $Pr(A \text{ or } B) =$

$$\Pr(A) + \Pr(B) = p_1 + p_2 + p_3 + p_5 + p_6$$

Why $\Pr(A \text{ or } B) = \Pr(A) + \Pr(B)$?


- Consider event $A = \{1, 2, 3\}$
- Consider event $B = \{5, 6\}$

0 Example: Dice Throw and Venn Diagram

| 11

Experiment: Throw a fair dice once

Sample Space: $S = \{1, 2, 3, 4, 5, 6\}$



- Consider event $A = \{1, 2, 3\}$
- Consider event $B = \{5, 6\}$

- Since the dice is fair, we have $p_1 + p_2 + \dots + p_6 = 1$
- $Pr(A) = p_1 + p_2 + p_3$
- $Pr(B) = p_5 + p_6$
- Consider the event A or $B = \{1, 2, 3, 5, 6\}$
- $Pr(A \text{ or } B) =$

$$Pr(A) + Pr(B) = p_1 + p_2 + p_3 + p_5 + p_6$$

Why $Pr(A \text{ or } B) = Pr(A) + Pr(B)$?

When are we allowed to add probabilities?

Mutually Exclusive Events

Two events are said to be **mutually exclusive** if they have disjoint set of outcomes.

Mutually Exclusive Events

Two events are said to be **mutually exclusive** if they have disjoint set of outcomes. Let A and B be two **mutually exclusive events**, then we have

$$Pr(A \cup B) = Pr(A) + Pr(B)$$

Mutually Exclusive Events

Two events are said to be **mutually exclusive** if they have disjoint set of outcomes. Let A and B be two **mutually exclusive events**, then we have

$$Pr(A \cup B) = Pr(A) + Pr(B)$$

Probability of a complement

Let A be any event, and A^c be its **complement**.

Mutually Exclusive Events

Two events are said to be **mutually exclusive** if they have disjoint set of outcomes. Let A and B be two **mutually exclusive events**, then we have

$$Pr(A \cup B) = Pr(A) + Pr(B)$$

Probability of a complement

Let A be any event, and A^c be its **complement**. Then clearly $A \cup A^c$ is the **complete set of outcomes**, i.e., it is a **sample space**.

Mutually Exclusive Events

Two events are said to be **mutually exclusive** if they have disjoint set of outcomes. Let A and B be two **mutually exclusive events**, then we have

$$Pr(A \cup B) = Pr(A) + Pr(B)$$

Probability of a complement

Let A be any event, and A^c be its **complement**. Then clearly $A \cup A^c$ is the **complete set of outcomes**, i.e., it is a **sample space**. Hence, we have

$$Pr(A) + Pr(A^c) = 1 \implies \underline{Pr(A^c) = 1 - Pr(A)}$$

0 Mutually Exclusive Events..

Mutually disjoint events

A set of n events denoted by A_1, A_2, \dots, A_n are called mutually disjoint events if the following holds:

Mutually disjoint events

A set of n events denoted by A_1, A_2, \dots, A_n are called **mutually disjoint events** if the following holds:

$$P(A_1 \cup \dots \cup A_n) = P(A_1) + \dots + P(A_n).$$

Mutually disjoint events

A set of n events denoted by A_1, A_2, \dots, A_n are called **mutually disjoint events** if the following holds:

$$P(A_1 \cup \dots \cup A_n) = P(A_1) + \dots + P(A_n).$$

From **Axiom-3**, it holds for **countably infinite unions**.

0 Quiz: Use concept of disjoint events

| 14

Quiz

In a cricket tournament with four teams denoted by $\{A, B, C, D\}$, team A has 20% chance of winning, while team B has a 40% chance of winning. What is the probability that A or B win the tournament?

$$P(A \text{ wins}) = 20\%$$

$$P(B \text{ wins}) = 40\%$$

$$\begin{aligned} P(A \text{ or } B \text{ wins}) &= P(A \text{ wins}) + P(B \text{ wins}) \\ &\quad \text{Disjoint events} \\ &= 60\% \end{aligned}$$

Quiz

Suppose we know the following:

- there is a 50% chance that it will be hot Today
- there is a 30% chance that it will be hot Tomorrow
- there is 10% chance that it wont be hot Today or Tomorrow

Quiz

Suppose we know the following:

- there is a 50% chance that it will be hot Today
- there is a 30% chance that it will be hot Tomorrow
- there is 10% chance that it wont be hot Today or Tomorrow

Answer the following:

0 Quiz-Use probability axioms

$$P(A \cap B) = \frac{0.9}{0.1} = 0.8$$

Quiz

Suppose we know the following:

- there is a 50% chance that it will be hot Today
- there is a 30% chance that it will be hot Tomorrow
- there is 10% chance that it ~~wont~~^{neither} ~~nor~~^{and} be hot Today or Tomorrow

Answer the following:

- probability that it will be hot today or tomorrow
- probability that it will be hot today and tomorrow
- probability that it will be hot today but not tomorrow
- probability that it either will be hot today or tomorrow, but not both

$$\begin{aligned} P(A \cup B) &= P(A) + P(B) - P(A \cap B) \\ &= 0.5 + 0.3 - 0.9 = -0.1 \end{aligned}$$

$\{A : \text{hot Today}$

$B : \text{hot Tomorrow}$

$P(B^c) = 0.7$

$| 15$

$$\begin{aligned} P(A) &= 50\% = 0.5 \\ P(B) &= 30\% = 0.3 \\ P(A^c \cup B^c) &= 10\% = 0.1 \end{aligned}$$

$$P(A \cup B)$$

$$P(A \cap B)$$

$$P(A \cap B^c)$$

$$P((A \cup B) \setminus (A \cap B))$$

$$\begin{aligned} P(A^c \cap B^c) &= 10\% = 0.1 \\ P((A \cup B)^c) &= 0.1 \\ P(A^c \cup B^c) &= 0.9 \\ P(A \cup B) &= 0.1 \end{aligned}$$

0 Non-Mutually Exclusive Events

| 16

- What if the two events A and B are **not** mutually exclusive?

0 Non-Mutually Exclusive Events

| 16

- What if the two events A and B are **not mutually exclusive**?
- What relation we have

$$Pr(A \cup B) \quad ? \quad Pr(A) + Pr(B) \quad (1)$$


0 Non-Mutually Exclusive Events

| 16

- What if the two events A and B are **not mutually exclusive**?
- What relation we have

$$Pr(A \cup B) \quad ? \quad Pr(A) + Pr(B) \quad (1)$$

- Let us consider an example below with two events A and B shaded




0 Non-Mutually Exclusive Events

| 16

- What if the two events A and B are **not mutually exclusive**?
- What relation we have


$$Pr(A \cup B) \quad ? \quad Pr(A) + Pr(B) \quad (1)$$

- Let us consider an example below with two events A and B shaded




- For the events A and B , which relation (1) holds?

- Consider the example as before




- Consider the example as before



- Let us calculate the probabilities of the two events A and B

$$Pr(A) = p_1 + p_2 + p_3, \quad Pr(B) = p_3 + p_4$$

- Consider the example as before




- Let us calculate the probabilities of the two events A and B

$$Pr(A) = p_1 + p_2 + p_3, \quad Pr(B) = p_3 + p_4$$

- Now we calculate probability of $A \cup B$

- Consider the example as before




- Let us calculate the probabilities of the two events A and B

$$Pr(A) = p_1 + p_2 + p_3, \quad Pr(B) = p_3 + p_4$$

- Now we calculate probability of $A \cup B$

$$Pr(A \cup B) = Pr(\{1, 2, 3, 4\}) = p_1 + p_2 + p_3 + p_4$$

- Consider the example as before



- Let us calculate the probabilities of the two events A and B


$$Pr(A) = p_1 + p_2 + p_3, \quad Pr(B) = p_3 + p_4$$

- Now we calculate probability of $A \cup B$

$$Pr(A \cup B) = Pr(\{1, 2, 3, 4\}) = p_1 + p_2 + p_3 + p_4$$

- Using above: $Pr(A \cup B) = Pr(A) + Pr(B) - Pr(A \cap B)$ ↗

- Consider the example as before



- Let us calculate the probabilities of the two events A and B

$$Pr(A) = p_1 + p_2 + p_3, \quad Pr(B) = p_3 + p_4$$

- Now we calculate probability of $A \cup B$

$$Pr(A \cup B) = Pr(\{1, 2, 3, 4\}) = p_1 + p_2 + p_3 + p_4$$

- Using above: $Pr(A \cup B) = Pr(A) + Pr(B) - Pr(A \cap B)$
- Recall discrete mathematics: inclusion-exclusion principle. (Proof?)

0 Generalized Inclusion-Exclusion Principle

Inclusion-Exclusion Principle

- $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

Inclusion-Exclusion Principle

- $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
- $P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(A \cap C) + P(A \cap B \cap C)$

Inclusion-Exclusion Principle

- $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
- $P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(A \cap C) + P(A \cap B \cap C)$
- In general for n events, we have

Inclusion-Exclusion Principle

- $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
- $P(A \cup B \cup C) = \underbrace{P(A) + P(B) + P(C)}_{+} - P(A \cap B) - P(B \cap C) - P(A \cap C) + P(A \cap B \cap C)$
- In general for n events, we have

$$P(\bigcup_{i=1}^n A_i) = \sum_{i=1}^n P(A_i) - \sum_{i < j} P(A_i \cap A_j) + \sum_{i < j < k} P(A_i \cap A_j \cap A_k) - \cdots (-1)^{n-1} P(\bigcap_{i=1}^n A_i)$$

- Sample space can be of two types: discrete or continuous

- Sample space can be of two types: discrete or continuous
- in discrete probability models we can compute the probability of events by adding all the corresponding outcomes

- Sample space can be of two types: discrete or continuous
- in discrete probability models we can compute the probability of events by adding all the corresponding outcomes
- in continuous probability models we need to use integration instead of summation

- Sample space can be of two types: discrete or continuous
- in discrete probability models we can compute the probability of events by adding all the corresponding outcomes
- in continuous probability models we need to use integration instead of summation

Definition of discrete probability model

If the sample space S is a countable set, then it refers to discrete probability model. Since S is countable:
$$S = \{s_1, s_2, \dots\}$$

- Sample space can be of two types: discrete or continuous
- in discrete probability models we can compute the probability of events by adding all the corresponding outcomes
- in continuous probability models we need to use integration instead of summation

Definition of discrete probability model

If the sample space S is a countable set, then it refers to discrete probability model. Since S is countable:

$$S = \{s_1, s_2, \dots\}$$


- For an event $A \subset S$, by 3rd axiom

$$P(A) = \sum_{s_j \in A} P(s_j)$$


- Consider 6 balls labelled 1,2,3,4,5,6

- Consider 6 balls labelled 1,2,3,4,5,6
- The balls are kept in two boxes. In the first box, we have 1,2 balls, and in the second box, we have 3,4,5,6 balls

- Consider 6 balls labelled 1,2,3,4,5,6
- The balls are kept in two boxes. In the first box, we have 1,2 balls, and in the second box, we have 3,4,5,6 balls




- Consider 6 balls labelled 1,2,3,4,5,6
- The balls are kept in two boxes. In the first box, we have 1,2 balls, and in the second box, we have 3,4,5,6 balls




- Assume the following:

- Consider 6 balls labelled 1,2,3,4,5,6
- The balls are kept in two boxes. In the first box, we have 1,2 balls, and in the second box, we have 3,4,5,6 balls




- Assume the following:
 - The two boxes are **equiprobable**

- Consider 6 balls labelled 1,2,3,4,5,6
- The balls are kept in two boxes. In the first box, we have 1,2 balls, and in the second box, we have 3,4,5,6 balls




- Assume the following:
 - The two boxes are **equiprobable**
 - All the balls in each box is **equiprobable**

- Consider 6 balls labelled 1,2,3,4,5,6
 - The balls are kept in two boxes. In the first box, we have 1,2 balls, and in the second box, we have 3,4,5,6 balls



- Assume the following:
 - The two boxes are **equiprobable**
 - All the balls in each box is **equiprobable**
 - **Question:** What are the probabilities p_1, p_2, \dots, p_6 ?






Figure: Consider a choice tree for the problem

- We have the following

$$p_1 + p_2 = 1/2, \quad p_3 + p_4 + p_5 + p_6 = 1/2$$




Figure: Consider a choice tree for the problem

- We have the following

$$p_1 + p_2 = 1/2, \quad p_3 + p_4 + p_5 + p_6 = 1/2$$

- Balls in box-1 are **equiprobable**: $p_1 = p_2$





Figure: Consider a choice tree for the problem

- We have the following

$$p_1 + p_2 = 1/2, \quad p_3 + p_4 + p_5 + p_6 = 1/2$$




- Balls in box-1 are **equiprobable**: $p_1 = p_2$
- Balls in box-2 are **equiprobable**: $p_3 = p_4 = p_5 = p_6$

Figure: Consider a choice tree for the problem

- We have the following

$$p_1 + p_2 = 1/2, \quad p_3 + p_4 + p_5 + p_6 = 1/2$$



- Balls in box-1 are **equiprobable**: $p_1 = p_2$
- Balls in box-2 are **equiprobable**: $p_3 = p_4 = p_5 = p_6$
- From above, we have

$$p_1 = p_2 = 1/4, \quad p_3 = p_4 = p_5 = p_6 = 1/8$$

Figure: Consider a choice tree for the problem

Shrewd Prisoner Problem

Long ago a prisoner was to be executed.

Shrewd Prisoner Problem

Long ago a prisoner was to be executed. In response to his supplications, he was promised that he would be released if he drew a white ball from one of two similar urns.

Shrewd Prisoner Problem

Long ago a prisoner was to be executed. In response to his supplications, he was promised that he would be released if he drew a white ball from one of two similar urns. The provisions were that he had to distribute 50 white and 50 black balls between the two urns,

Shrewd Prisoner Problem


Long ago a prisoner was to be executed. In response to his supplications, he was promised that he would be released if he drew a white ball from one of two similar urns. The provisions were that he had to distribute 50 white and 50 black balls between the two urns, in any way he liked,

0 Shrewd Prisoner and Maximizing Chance!

| 22

Shrewd Prisoner Problem

Long ago a prisoner was to be executed. In response to his supplications, he was promised that he would be released if he drew a white ball from one of two similar urns. The provisions were that he had to distribute 50 white and 50 black balls between the two urns, in any way he liked, after which he had to draw a ball at random from one of these urns.



Shrewd Prisoner Problem

Long ago a prisoner was to be executed. In response to his supplications, he was promised that he would be released if he drew a white ball from one of two similar urns. The provisions were that he had to distribute 50 white and 50 black balls between the two urns, in any way he liked, after which he had to draw a ball at random from one of these urns.

Question: How should the prisoner put the balls such that the probability of his release is maximized?

0 Solution to Shrewd Prisoner Problem

| 23

- How did the prisoner arrange to make his chance of success as great as possible?

0 Solution to Shrewd Prisoner Problem

| 23

- How did the prisoner arrange to make his chance of success as great as possible?
- What happens if he puts equal number of white and black balls in two boxes?

0 Solution to Shrewd Prisoner Problem

| 23

- How did the prisoner arrange to make his chance of success as great as possible?
- What happens if he puts equal number of white and black balls in two boxes?
 - The probability of choosing a white ball (from either boxes) is $1/2$

0 Solution to Shrewd Prisoner Problem

| 23

- How did the prisoner arrange to make his chance of success as great as possible?
- What happens if he puts equal number of white and black balls in two boxes?
 - The probability of choosing a white ball (from either boxes) is $1/2$
 - Can he do better?

0 Solution to Shrewd Prisoner Problem

| 23

- How did the prisoner arrange to make his chance of success as great as possible?
- What happens if he puts equal number of white and black balls in two boxes?
 - The probability of choosing a white ball (from either boxes) is $1/2$
 - Can he do better?
- What if he puts all white in one, and all black in other urn?

0 Solution to Shrewd Prisoner Problem

| 23

- How did the prisoner arrange to make his chance of success as great as possible?
- What happens if he puts equal number of white and black balls in two boxes?
 - The probability of choosing a white ball (from either boxes) is $1/2$
 - Can he do better?
- What if he puts all white in one, and all black in other urn?
 - probability of choosing white from urn-1 is $1/2$, but probability of choosing white from urn-2 is 0. Can he do better?

0 Solution to Shrewd Prisoner Problem

| 23

- How did the prisoner arrange to make his chance of success as great as possible?
- What happens if he puts equal number of white and black balls in two boxes?
 - The probability of choosing a white ball (from either boxes) is $1/2$
 - Can he do better?
- What if he puts all white in one, and all black in other urn?
 - probability of choosing white from urn-1 is $1/2$, but probability of choosing white from urn-2 is 0. Can he do better?
- Can he do better than this? If yes, then how?

0 Solution to Shrewd Prisoner Problem

| 23

- How did the prisoner arrange to make his chance of success as great as possible?
- What happens if he puts equal number of white and black balls in two boxes?
 - The probability of choosing a white ball (from either boxes) is $1/2$
 - Can he do better?
- What if he puts all white in one, and all black in other urn?
 - probability of choosing white from urn-1 is $1/2$, but probability of choosing white from urn-2 is 0. Can he do better?
- Can he do better than this? If yes, then how?
- Put 1 white and 0 black in urn-1, and put 49 white and 100 black in urn-2

(W, 0B)

(49 W, 50 B)

$\frac{1}{2} \cdot 1 + \frac{1}{2} \cdot 0$

49
B

0 Solution to Shrewd Prisoner Problem

| 23

- How did the prisoner arrange to make his chance of success as great as possible?
- What happens if he puts equal number of white and black balls in two boxes?
 - The probability of choosing a white ball (from either boxes) is 1/2
 - Can he do better?
- What if he puts all white in one, and all black in other urn?
 - probability of choosing white from urn-1 is 1/2, but probability of choosing white from urn-2 is 0. Can he do better?
- Can he do better than this? If yes, then how?
- Put 1 white and 0 black in urn-1, and put 49 white and 100 black in urn-2
- The probability of success is $\frac{1}{2} \times 1 + \frac{1}{2} \times \frac{49}{99} \approx 3/4$

$$\frac{1}{2} \times 1 + \frac{1}{2} \times \frac{49}{99} \approx 3/4$$


0 Extension of the problem

| 24

100 W
100 B

Quiz-1

What is the chance of success of the prisoner if he had been allowed to distribute 100 balls among 4 urns?



Quiz-1

What is the chance of success of the prisoner if he had been allowed to distribute 100 balls among 4 urns?

Quiz-2

What if the number of balls is increased?

- ① Digress: Game Theory
- ② Random Walks
- ③ Conditional Probability, Bayes Theorem
- ④ Define Conditional Probability and Chain Rule
- ⑤ The Monty Hall Problem
- ⑥ Independence
- ⑦ Conditional Independence

1 A related Problem in Game Theory: Prisoner's dilemma

| 26



1 A related Problem in Game Theory: Prisoner's dilemma

| 26



- If both of them do not confess, they serve 2 years each

1 A related Problem in Game Theory: Prisoner's dilemma

| 26



- If both of them do not confess, they serve 2 years each
- If they confess to crime, partner does not, then person confessing will be granted immunity. However, in this case the partner not confessing will serve 10 years

1 A related Problem in Game Theory: Prisoner's dilemma

| 26



- If both of them do not confess, they serve 2 years each
- If they confess to crime, partner does not, then person confessing will be granted immunity. However, in this case the partner not confessing will serve 10 years
- If both confess, then both end up spending 5 years each in jail

1 A related Problem in Game Theory: Prisoner's dilemma

| 26



- If both of them do not confess, they serve 2 years each
- If they confess to crime, partner does not, then person confessing will be granted immunity. However, in this case the partner not confessing will serve 10 years
- If both confess, then both end up spending 5 years each in jail
- Given that Wanda and Fred have no reason to trust, what is the good option?

 Wanda	 Fred	DON'T CONFESS ...	CONFESS!
DON'T CONFESS ...		2/2	10/0
CONFESS!		0/10	5/5

 Wanda	 Fred	DON'T CONFESS ...	CONFESS
DON'T CONFESS ...		2/2	10/0
CONFESS		0/10	5/5

- Indeed, if both don't confess,

 Wanda	 Fred	DON'T CONFESS ...	CONFESS !
DON'T CONFESS ...		2/2	10/0
CONFESS !		0/10	5/5

- Indeed, if both don't confess, then it is **best** for them;

 Wanda Fred	DON'T CONFESS 	 CONFESS
DON'T CONFESS 	2/2	10/0
 CONFESS	0/10	5/5

- Indeed, if both don't confess, then it is **best** for them; 2 years each!

		DON'T CONFESS ...	! CONFESS
DON'T CONFESS ...		2/2	10/0
! CONFESS		0/10	5/5

- Indeed, if both don't confess, then it is **best** for them; 2 years each!
- But another best option is when they both confess. Because in other cases, one of them has to spend 10 years in jail

 Wanda	 Fred	DON'T CONFESS 	 CONFESS
DON'T CONFESS 	$2/2$	$10/0$	
 CONFESS	$0/10$	$5/5$	



- Indeed, if both don't confess, then it is **best** for them; 2 years each!
- But another best option is when they both confess. Because in other cases, one of them has to spend 10 years in jail
- This is part of **co-operative games**, and 5/5 is called **Nash equilibrium**
- These topics are part of topic known as **game theory** (John Nash!)

1 About John Nash

| 28



1 About John Nash

| 28



- John Forbes Nash Jr. (June 13, 1928 – May 23, 2015) was an American mathematician who made fundamental contributions to game theory, differential geometry, and the study of partial differential equations

1 About John Nash

| 28



- John Forbes Nash Jr. (June 13, 1928 – May 23, 2015) was an American mathematician who made fundamental contributions to game theory, differential geometry, and the study of partial differential equations
- Awards
 - Nobel Prize in 1994

1 About John Nash

| 28



- John Forbes Nash Jr. (June 13, 1928 – May 23, 2015) was an American mathematician who made fundamental contributions to game theory, differential geometry, and the study of partial differential equations
- Awards
 - Nobel Prize in 1994
 - Abel Prize in 2015



1 About John Nash

| 28



- John Forbes Nash Jr. (June 13, 1928 – May 23, 2015) was an American mathematician who made fundamental contributions to game theory, differential geometry, and the study of partial differential equations
- Awards
 - Nobel Prize in 1994
 - Abel Prize in 2015
- Movie on him: “A Beautiful Mind”

1 About John Nash

| 28



- John Forbes Nash Jr. (June 13, 1928 – May 23, 2015) was an American mathematician who made fundamental contributions to game theory, differential geometry, and the study of partial differential equations
- Awards
 - Nobel Prize in 1994
 - Abel Prize in 2015
- Movie on him: “*A Beautiful Mind*”
- Bar Scene (Game theory part...):
<https://www.youtube.com/watch?v=LJS7Igvk6ZM>

1 About John Nash

| 28



- John Forbes Nash Jr. (June 13, 1928 – May 23, 2015) was an American mathematician who made fundamental contributions to game theory, differential geometry, and the study of partial differential equations
- Awards
 - Nobel Prize in 1994
 - Abel Prize in 2015
- Movie on him: “A Beautiful Mind”
- Bar Scene (Game theory part...):
<https://www.youtube.com/watch?v=LJS7Igvk6ZM>

More on this in Topics in Applied Optimization Elective!