



# Probability and Statistics

UG2, Core course, IIIT,H

Pawan Kumar

IIIT, Hyderabad

September 28, 2021

- ① Conditional Probability, Bayes Theorem
- ② Conditional Independence

- ③ Random Variables
- ④ Expectation
- ⑤ Saint Petersburg Paradox

## Outline

- ① Conditional Probability, Bayes Theorem
- ② Conditional Independence
- ③ Random Variables
- ④ Expectation
- ⑤ Saint Petersburg Paradox

## Examples involving independent trials...

### Problem: coin toss

Suppose we flip a coin  $n$  times. Each coin flip is an independent trial with probability  $p$  of coming up heads. Write an expression for the following:

## Examples involving independent trials...

### Problem: coin toss

Suppose we flip a coin  $n$  times. Each coin flip is an independent trial with probability  $p$  of coming up heads. Write an expression for the following:

- $P(n \text{ heads on } n \text{ coin flips})$

$$P(\underbrace{H \ H \ \dots \ H}_{n \text{ times}})$$

## Examples involving independent trials...

### Problem: coin toss

Suppose we flip a coin  $n$  times. Each coin flip is an independent trial with probability  $p$  of coming up heads. Write an expression for the following:

- $P(n \text{ heads on } n \text{ coin flips})$

- $P(n \text{ tails on } n \text{ coin flips})$

$P(TT \dots T)$

## Examples involving independent trials...

### Problem: coin toss

Suppose we flip a coin  $n$  times. Each coin flip is an independent trial with probability  $p$  of coming up heads. Write an expression for the following:

- $P(n \text{ heads on } n \text{ coin flips})$
- $P(n \text{ tails on } n \text{ coin flips})$
- $P(\text{first } k \text{ heads, then } n - k \text{ tails})$

$$P(\underbrace{H \cdots H}_k \underbrace{T \cdots T}_{n-k})$$

## Examples involving independent trials...

### Problem: coin toss

Suppose we flip a coin  $n$  times. Each coin flip is an independent trial with probability  $p$  of coming up heads. Write an expression for the following:

- $P(n \text{ heads on } n \text{ coin flips})$
- $P(n \text{ tails on } n \text{ coin flips})$
- $P(\text{first } k \text{ heads, then } n - k \text{ tails})$
- $P(\text{exactly } k \text{ heads on } n \text{ coin flips})$

$$\begin{aligned} &= P(H_1 \cap H_2 \cap H_3 \dots \cap H_n) = P(H_1)P(H_2) \dots P(H_n) \\ &\quad \text{because } H_i \text{'s are independ.} \\ &= \underbrace{p \cdot p \dots p}_{n \text{ times}} = p^n \end{aligned}$$

$$\binom{n}{k} p^k (1-p)^{n-k}$$

$$P(T_1 \cap T_2 \cap \dots \cap T_n) = P(T_1)P(T_2) \dots P(T_n)$$

$$= P(H_1^c)P(H_2^c) \dots P(H_n^c)$$

$$= (1-P(H_1))(1-P(H_2)) \dots (1-P(H_n))$$

$$= (1-p)^n$$

$$\begin{aligned} &P(H_1 \cap \dots \cap H_k \cap T_{k+1} \cap \dots \cap T_n) \\ &= P(H_1) \dots P(H_k)P(T_{k+1}) \dots P(T_n) \\ &= p^k (1-p)^{n-k} \end{aligned}$$



**Biased Coin, Independence, Infinite Sample Space, Total Probability, Bayes Theorem...**

## Biased Coin, Independence, Infinite Sample Space, Total Probability, Bayes Theorem...

2, 4, 6, 8, ...

### Problem

A biased coin (with probability of obtaining a Head equal to  $p > 0$ ) is tossed repeatedly and independently until the first head is observed. Compute the probability that the first head appears at an even numbered toss.

Solution Consider the partition of  $E$  into  $E_1, E_2, \dots, E_k, \dots$  where  $E_k$  = event that 1st head occurs on the  $2k^{\text{th}}$  toss even!

$E = \bigcup_{k=1}^{\infty} E_k$ ,  $E_k$ 's are mutually exclusive!

$$P(E) = \sum_{k=1}^{\infty} P(E_k), \quad P(E_k) = \underline{(1-p)^{2k-1}} \cdot \underline{p}$$

$$P(E) = \sum_{k=1}^{\infty} (1-p)^{2k-1} \cdot p = \frac{1-p}{2-p} //$$

## Biased Coin, Independence, Infinite Sample Space, Total Probability, Bayes Theorem...

### Problem

A **biased** coin (with probability of obtaining a Head equal to  $p > 0$ ) is tossed **repeatedly and independently** until the first head is observed. Compute the probability that the first head appears at an even numbered toss.

Solution to the problem...

## Biased Coin, Independence, Infinite Sample Space, Total Probability, Bayes Theorem...

### Problem

A **biased** coin (with probability of obtaining a Head equal to  $p > 0$ ) is tossed **repeatedly and independently** until the first head is observed. Compute the probability that the first head appears at an even numbered toss.

### Solution to the problem...

What are sample space and events in this problem?

## Biased Coin, Independence, Infinite Sample Space, Total Probability, Bayes Theorem...

### Problem

A **biased** coin (with probability of obtaining a Head equal to  $p > 0$ ) is tossed **repeatedly and independently** until the first head is observed. Compute the probability that the first head appears at an even numbered toss.

### Solution to the problem...

What are sample space and events in this problem?

- 1 Sample space,  $S$  = all possible infinite binary sequences of coin toss.

## Biased Coin, Independence, Infinite Sample Space, Total Probability, Bayes Theorem...

### Problem

A **biased** coin (with probability of obtaining a Head equal to  $p > 0$ ) is tossed **repeatedly and independently** until the first head is observed. Compute the probability that the first head appears at an even numbered toss.

### Solution to the problem...

What are sample space and events in this problem?

- 1 Sample space,  $S$  = all possible infinite binary sequences of coin toss.
- 2 Consider event  $H_1$  : head on first toss.

## Biased Coin, Independence, Infinite Sample Space, Total Probability, Bayes Theorem...

### Problem

A biased coin (with probability of obtaining a Head equal to  $p > 0$ ) is tossed repeatedly and independently until the first head is observed. Compute the probability that the first head appears at an even numbered toss.

### Solution to the problem...

What are sample space and events in this problem?

- 1 Sample space,  $S$  = all possible infinite binary sequences of coin toss.
- 2 Consider event  $H_1$  : head on first toss.
- 3 Consider event  $E$  : first head on even numbered toss.

We want to compute  $P(E)$ .

## Biased Coin, Independence, Infinite Sample Space, Total Probability, Bayes Theorem...

### Problem

A biased coin (with probability of obtaining a Head equal to  $p > 0$ ) is tossed repeatedly and independently until the first head is observed. Compute the probability that the first head appears at an even numbered toss.

### Solution to the problem...

What are sample space and events in this problem?

- 1 Sample space,  $S$  = all possible infinite binary sequences of coin toss.
- 2 Consider event  $H_1$  : head on first toss.
- 3 Consider event  $E$  : first head on even numbered toss.

We want to compute  $P(E)$ .

How do we solve problems like this?



**Solution...**

**Solution...**

## Coin Toss Example...

## Coin Toss Example...


→ TT | HTHT ... TT ← experiment ends!  
→ TT | HTHT ... TT  
→ TT | HTHT ... TT  
n-1 n

### Problem

A coin for which  $P(\text{Heads}) = p$  is tossed until two successive Tails are obtained. Find the probability that the experiment is completed on the  $n$ th toss.

$E_1$ : 1st toss is H  
 $E_2$ : first two tosses are TH  
 $E_3$ : first two tosses are TT  
 $F_n$ : experiment completes on  $n$ th toss.  
partition  $\Omega$  (sample sp.)  
 $E_1, E_2, E_3$

For  $n=2$ :  $p_2 = P(F_2) = (1-p)(1-p) = \underline{\underline{(1-p)^2}}$



Must have  
 $n$  at least 2

## Coin Toss Example...

### Problem

A coin for which  $P(\text{Heads}) = p$  is tossed until two successive Tails are obtained. Find the probability that the experiment is completed on the  $n$ th toss.

### Solution

What are sample space and events?

## Coin Toss Example...

### Problem

A coin for which  $P(\text{Heads}) = p$  is tossed until two successive Tails are obtained. Find the probability that the experiment is completed on the  $n$ th toss.

### Solution

What are sample space and events?

- 1 Sample space,  $S$ : all possible infinite sequences of tosses

## Coin Toss Example...

### Problem

A coin for which  $P(\text{Heads}) = p$  is tossed until two successive Tails are obtained. Find the probability that the experiment is completed on the  $n$ th toss.

### Solution

What are sample space and events?

- 1 Sample space,  $S$ : all possible infinite sequences of tosses
- 2 event  $E_1$  : first toss is  $H$

## Coin Toss Example...

### Problem

A coin for which  $P(\text{Heads}) = p$  is tossed until two successive Tails are obtained. Find the probability that the experiment is completed on the  $n$ th toss.

### Solution

What are sample space and events?

- 1 Sample space,  $S$ : all possible infinite sequences of tosses
- 2 event  $E_1$  : first toss is  $H$
- 3 event  $E_2$  : first two tosses are  $TH$



## Coin Toss Example...

### Problem

A coin for which  $P(\text{Heads}) = p$  is tossed until two successive Tails are obtained. Find the probability that the experiment is completed on the  $n$ th toss.

### Solution

What are sample space and events?

- 1 Sample space,  $S$ : all possible infinite sequences of tosses
- 2 event  $E_1$  : first toss is  $H$
- 3 event  $E_2$  : first two tosses are  $TH$
- 4 event  $E_3$  : first two tosses are  $TT$

## Coin Toss Example...

### Problem

A coin for which  $P(\text{Heads}) = p$  is tossed until two successive Tails are obtained. Find the probability that the experiment is completed on the  $n$ th toss.

### Solution

What are sample space and events?

- 1 Sample space,  $S$ : all possible infinite sequences of tosses
- 2 event  $E_1$  : first toss is  $H$
- 3 event  $E_2$  : first two tosses are  $TH$
- 4 event  $E_3$  : first two tosses are  $TT$
- 5 event  $F_n$  : experiment completed on the  $n$ th toss.

Solution to problem in previous slide...part-1

For  $n > 2$ :  $P(F_n | E_1) = P(F_{n-1})$

$$P(F_n | E_2) = P(F_{n-2})$$

$$P(F_n | E_3) = 0$$

Using total probab.  $n > 2$ , Notation:  $P(F_n) = p_n$

$$P(F_n) = P(F_n | E_1)P(E_1) + P(F_n | E_2)P(E_2) + P(F_n | E_3)P(E_3)$$

$$p_n = p_{n-1} \cdot p + p_{n-2} \cdot (1-p)p + 0$$

$$p_n = p p_{n-1} + p(1-p) p_{n-2}$$

$$\begin{aligned} p_1 &= 0 \\ p_2 &= (1-p)^2 \end{aligned}$$

Handwritten notes and diagrams:  
- A sequence of outcomes:  $H \dots TT$  with  $H$  and  $TT$  circled in red.  
- An arrow points from the text "as if exp. started from  $n=2$ " to the circled  $TT$ .  
- Another sequence:  $TH \dots TT$  with  $TH$  circled in red and  $TT$  circled in green.  
- A green circle contains the text  $TT$ .

Solution to problem in previous slide...part-2

Solution to problem in previous slide...part-3

## Properties of conditional probabilities...

## Properties of conditional probabilities...

### Properties

For any events  $A, B$ , and  $E$  we have the following:

## Properties of conditional probabilities...

### Properties

For any events  $A, B$ , and  $E$  we have the following:

- $0 \leq P(\underbrace{A \cap E}) \leq 1$



## Properties of conditional probabilities...

### Properties

For any events  $A, B$ , and  $E$  we have the following:

- $0 \leq P(A \cap E) \leq 1$
- $P(\underline{A} | E) = 1 - P(A^c | E)$

$$P(A|E) = \frac{P(A \cap E)}{P(E)}$$

$$P(A^c|E) = \frac{P(A^c \cap E)}{P(E)}$$

$$P(A|E) + P(A^c|E) = \frac{1}{P(E)} [P(A \cap E) + P(A^c \cap E)]$$


$$E = \underbrace{(A \cap E)}_{\text{mutually excl.}} \cup \underbrace{(A^c \cap E)}_{\text{mutually excl.}}$$

$$P(E) = P(A \cap E) + P(A^c \cap E)$$

## Properties of conditional probabilities...

### Properties

For any events  $A, B$ , and  $E$  we have the following:

- $0 \leq P(A \cap E) \leq 1$
- $P(A \mid E) = 1 - P(A^c \mid E)$
- $P(A \cap B \mid E) = P(B \cap A \mid E)$  

## Properties of conditional probabilities...

### Properties

For any events  $A, B$ , and  $E$  we have the following:

- $0 \leq P(A \cap E) \leq 1$
- $P(A \mid E) = 1 - P(A^c \mid E)$
- $P(A \cap B \mid E) = P(B \cap A \mid E)$
- $P(A \cap B \mid E) = P(B \mid E)P(A \mid B \cap E)$

## Properties of conditional probabilities...

### Properties

For any events  $A, B$ , and  $E$  we have the following:

- $0 \leq P(A \cap E) \leq 1$
- $P(A | E) = 1 - P(A^c | E)$
- $P(A \cap B | E) = P(B \cap A | E)$  ←
- $P(A \cap B | E) = P(B | E)P(A | B \cap E)$  ←
- $P(A | B \cap E) = \frac{P(B | A \cap E)P(A | E)}{P(B | E)}$

## Scratch Space for Proving Conditional Probabilities...

## Scratch Space for Proving Conditional Probabilities...

## Outline

- ① Conditional Probability, Bayes Theorem
- ② Conditional Independence
- ③ Random Variables
- ④ Expectation
- ⑤ Saint Petersburg Paradox

## Conditional Independence...



## Conditional Independence...

### Definition of conditional independence

Two events  $A$  and  $B$  are conditionally independent given  $E$  if

$$P(A \cap B | E) = P(A | E)P(B | E)$$

## Conditional Independence...

### Definition of conditional independence

Two events  $A$  and  $B$  are **conditionally independent** given  $E$  if

$$P(A \cap B | E) = P(A | E)P(B | E)$$

### Fact on Conditional Independence

$A$  and  $B$  independent **does not mean** that  $A$  and  $B$  are independent given  $E$ . That is,

$$\underbrace{P(A \cap B) = P(A)P(B)} \not\Rightarrow P(A \cap B | E) = P(A | E)P(B | E)$$

Quiz on independence...

## Quiz on independence...

### Quiz-1

Two events  $E$  and  $F$  are independent if

- 1 Knowing that  $F$  happens means that  $E$  can't happen

## Quiz on independence...

### Quiz-1

Two events  $E$  and  $F$  are independent if

- 1 Knowing that  $F$  happens means that  $E$  can't happen
- ✓ 2 Knowing that  $F$  happens doesn't change probability that  $E$  happened.

## Quiz on independence...

### Quiz-1

Two events  $E$  and  $F$  are independent if

- 1 Knowing that  $F$  happens means that  $E$  can't happen
- 2 Knowing that  $F$  happens doesn't change probability that  $E$  happened.

What is your answer?

## Quiz on independence...

### Quiz-1



Two events  $E$  and  $F$  are **independent** if

- 1 Knowing that  $F$  happens means that  $E$  can't happen
- 2 Knowing that  $F$  happens doesn't change probability that  $E$  happened.

What is your answer?

### Quiz-1

Are  $E$  and  $F$  independent in the following pictures (not to scale)?



$E, F$  ind.  
 $P(E \cap F) = P(E)P(F) \Rightarrow E, F$  not ind.  
 $= \frac{1}{4} \cdot \frac{1}{4} \neq 0$

$P(E)P(F) = \frac{2}{9} \cdot \frac{1}{3} = \frac{2}{9}$   
 $\Rightarrow E, F$  are ind.  
 $P(E \cap F) = \frac{2}{9}$   
 $P(E) = \frac{6}{9}, P(F) = \frac{1}{3}$

## Mutually Exclusive and Independent Events...



## Mutually Exclusive and Independent Events...



### Quiz

When are two events both mutually exclusive and independent?

## More Problems on Independent Trials...

### Problem: String-part 1

There are  $m$  strings that are hashed unequally into a hash table with  $n$  buckets. Each string hashed is an independent trial with probability  $p_i$  of getting hashed into bucket  $i$ . What is  $P(E)$  if

- $E =$  bucket 1 has  $\geq 1$  string hashed into it?

## More Problems on Independent Trials...

### Problem: String-part 2

There are  $m$  strings that are hashed unequally into a hash table with  $n$  buckets. Each string hashed is an independent trial with probability  $p_i$  of getting hashed into bucket  $i$ . What is  $P(E)$  if

- $E =$  at least 1 of buckets 1 to  $k$  has  $\geq 1$  string hashed into it?

## Outline

- ① Conditional Probability, Bayes Theorem
- ② Conditional Independence
- ③ Random Variables
- ④ Expectation
- ⑤ Saint Petersburg Paradox

## Random Variables

## Random Variables

Python ← JIT

### Examples of typed variables in C

In some languages, such as, C/C++. ~~we~~ we have the concept of a **typed variable**:

## Random Variables

### Examples of typed variables in C

In some languages, such as, C/C++. we have the concept of a **typed variable**:

- `int i = 4;`
- `float x = 10;`
- `char y = 'x';`

## Random Variables


### Examples of typed variables in C


In some languages, such as, C/C++. we have the concept of a **typed variable**:

- `int i = 4;`
- `float x = 10;`
- `char y = 'x';`

### Examples of random variable

Let  $X$  denote the outputs after we roll a die, then


$$X = 3$$


$$P(\{X=3\})$$

means that after rolling a die, we obtain 3 as output.



## Random Variables

### Examples of typed variables in C

In some languages, such as, C/C++. we have the concept of a **typed variable**:

- `int i = 4;`
- `float x = 10;`
- `char y = 'x';`

### Examples of random variable

Let  $X$  denote the outputs after we roll a die, then

$$X = 3$$

means that after rolling a die, we obtain 3 as output.

Since the number that is going to be assigned to **variable**  $X$  is going to be **random**, it is called **random variable**.

## Define Random Variable

## Define Random Variable

$$X: \{0, 1, 2, \dots\} \rightarrow \mathbb{R} \quad \begin{aligned} X=1 &\in \\ X=2 &\in \end{aligned} \quad P\{\underline{X=-2}\} \in [0, 1]$$

### Definition of Random Variable

A **random variable**  $X$  is a function from the sample space to the real numbers.

$$X: \underline{S} \rightarrow \mathbb{R} \quad P$$

## Define Random Variable

### Definition of Random Variable

A **random variable**  $X$  is a function from the sample space to the real numbers.

$$X : S \rightarrow \mathbb{R}$$

- We usually denote random variables by **capital** letters:  $X$ ,  $Y$ , ...

## Define Random Variable

### Definition of Random Variable

A **random variable**  $X$  is a function from the sample space to the real numbers.

$$X : S \rightarrow \mathbb{R}$$

- We usually denote random variables by **capital** letters:  $X, Y, \dots$
- Random variable is a function with **domain**  $S$  and **co-domain**  $\mathbb{R}$

## Define Random Variable

### Definition of Random Variable

A **random variable**  $X$  is a function from the sample space to the real numbers.

$$X : S \rightarrow \mathbb{R}$$

- We usually denote random variables by **capital** letters:  $X, Y, \dots$
- Random variable is a function with **domain**  $S$  and **co-domain**  $\mathbb{R}$
- The range of a random variable is the set of possible values of  $X$

## Define Random Variable

### Definition of Random Variable

A **random variable**  $X$  is a function from the sample space to the real numbers.

$$X : S \rightarrow \mathbb{R}$$

- We usually denote random variables by **capital** letters:  $X, Y, \dots$
- Random variable is a function with **domain**  $S$  and **co-domain**  $\mathbb{R}$
- The range of a random variable is the set of possible values of  $X$

### Examples of Random Variables...

Find the range of the following random variables:

- I toss a coin 10 times. Let  $X$  be the number of heads I observe

$$X = \{0, 1, 2, 3, \dots, 10\}$$

## Define Random Variable

### Definition of Random Variable

A **random variable**  $X$  is a function from the sample space to the real numbers.


$$X : S \rightarrow \mathbb{R}$$

- We usually denote random variables by **capital** letters:  $X, Y, \dots$
- Random variable is a function with **domain**  $S$  and **co-domain**  $\mathbb{R}$
- The range of a random variable is the set of possible values of  $X$

### Examples of Random Variables...

Find the range of the following random variables:

- I toss a coin 10 times. Let  $X$  be the number of heads I observe
- I toss a coin until the first tail appears. Let  $Y$  be the total number of coin tosses





## Quiz on Random Variable...

## Quiz on Random Variable...

### Quiz on Random Variable

Consider and Experiment: 3 coins are flipped. Let  $X$  be the number of tails. Answer the following:

## Quiz on Random Variable...

### Quiz on Random Variable

Consider and Experiment: 3 coins are flipped. Let  $X$  be the number of tails. Answer the following:

- What is the value of  $X$  for the outcomes?

## Quiz on Random Variable...

### Quiz on Random Variable

Consider and Experiment: 3 coins are flipped. Let  $X$  be the number of tails. Answer the following:

- What is the value of  $X$  for the outcomes?

- $(H, H, H) \leftarrow 0$
- $(T, T, H) \leftarrow 2$

## Quiz on Random Variable...

### Quiz on Random Variable

Consider and Experiment: 3 coins are flipped. Let  $X$  be the number of tails. Answer the following:

- What is the value of  $X$  for the outcomes?
  - $(H, H, H)$
  - $(T, T, H)$
- What is the event when  $X = 2$ ?

## Quiz on Random Variable...

### Quiz on Random Variable

Consider and Experiment: 3 coins are flipped. Let  $X$  be the number of tails. Answer the following:

- What is the value of  $X$  for the outcomes?

- $(H, H, H)$
- $(T, T, H)$

- What is the event when  $X = 2$ ?

- What is  $P(X = 2)$ ?

= event that we get 2 tails in when 3 coins are tossed.

$$\{X=2\} = \{(TTH), (HTT), (THT)\}$$

$\uparrow \quad \uparrow \quad \uparrow$   
 $2 \quad 2 \quad 2 = 6$

$$P(X=2) = \frac{3}{8}$$

**Random Variables are Not Events!**

## Random Variables are Not Events!

### Remarks on Random variables

- random variables are **not** events!



## Random Variables are Not Events!



### Remarks on Random variables

- random variables are **not** events!
- when a random variable is assigned a value, then it becomes event

$X = x$	Set of Outcomes	$P(X = k)$
$X = 0$	$\{(T, T, T)\}$	$1/8$
$X = 1$	$\{(H, T, T), (T, H, T), (T, T, H)\}$	$3/8$
$X = 2$	$\{(H, H, T), (H, T, H), (T, H, T)\}$	$3/8$
$X = 3$	$\{(H, H, H)\}$	$1/8$
$X \geq 4$	$\{\}$	$0$

Table: Consider an experiment where 3 coins are flipped, and  $X$  denotes number of heads

## Discrete Random Variables...

## Discrete Random Variables...



Recall: countable sets

A set  $A$  is **countable** if either it is a **finite** set, or it can be put in **1-1 correspondence** with set of natural numbers.

## Discrete Random Variables...

### Recall: countable sets

A set  $A$  is **countable** if either it is a **finite** set, or it can be put in **1-1 correspondence** with set of natural numbers.

### Discrete Random Variables

A random variable  $X$  is called discrete random variable, if its range is countable.

## Discrete Random Variables...

### Recall: countable sets

A set  $A$  is **countable** if either it is a **finite** set, or it can be put in **1-1 correspondence** with set of natural numbers.

### Discrete Random Variables

A random variable  $X$  is called **discrete random variable**, if its range is **countable**.

### Types of Random Variables...

## Discrete Random Variables...

### Recall: countable sets

A set  $A$  is **countable** if either it is a **finite** set, or it can be put in **1-1 correspondence** with set of natural numbers.

### Discrete Random Variables

A random variable  $X$  is called **discrete random variable**, if its range is **countable**.

### Types of Random Variables...

There are three types of random variables:

## Discrete Random Variables...

### Recall: countable sets

A set  $A$  is **countable** if either it is a **finite** set, or it can be put in **1-1 correspondence** with set of natural numbers.

### Discrete Random Variables

A random variable  $X$  is called **discrete random variable**, if its range is **countable**.

### Types of Random Variables...

There are three types of random variables:

- 1 discrete random variables

## Discrete Random Variables...

### Recall: countable sets

A set  $A$  is **countable** if either it is a **finite** set, or it can be put in **1-1 correspondence** with set of natural numbers.

### Discrete Random Variables

A random variable  $X$  is called **discrete random variable**, if its range is **countable**.

### Types of Random Variables...

There are three types of random variables:

- 1 discrete random variables
- 2 continuous random variables



## Discrete Random Variables...

### Recall: countable sets

A set  $A$  is **countable** if either it is a **finite** set, or it can be put in **1-1 correspondence** with set of natural numbers.

### Discrete Random Variables

A random variable  $X$  is called **discrete random variable**, if its range is **countable**.

### Types of Random Variables...

There are three types of random variables:

- 1 discrete random variables
- 2 continuous random variables
- 3 mixed random variables