



# Probability and Statistics

UG2, Core course, IIIT,H

Pawan Kumar

IIIT, Hyderabad

October 12, 2021

- ① Random Variables
- ② Special Distributions
  - Uniform Distribution
  - Bernoulli Distribution
  - Geometric Distribution
  - Binomial Distribution

## Poisson Distribution

- ③ Examples of Distributions
- ④ Expectations of Some Distributions
- ⑤ Variance and Standard Deviation: Understand Variability in Data
- ⑥ Higher Order Moments and Moment Generating Function

## Outline

- ① Random Variables
- ② Special Distributions
- ③ Examples of Distributions
- ④ Expectations of Some Distributions
- ⑤ Variance and Standard Deviation: Understand Variability in Data
- ⑥ Higher Order Moments and Moment Generating Function

## Outline

### ① Random Variables

### ② Special Distributions

Uniform Distribution

Bernoulli Distribution

Geometric Distribution

Binomial Distribution

Poisson Distribution

### ③ Examples of Distributions

### ④ Expectations of Some Distributions

### ⑤ Variance and Standard Deviation: Understand Variability in Data

### ⑥ Higher Order Moments and Moment Generating Function

## Motivation for Uniform Distribution: Distribution of a Die Roll...

## Motivation for Uniform Distribution: Distribution of a Die Roll...

### Example: Motivation for Uniform Distribution

Consider rolling a fair die. The possible outcomes are  $\{1, 2, 3, 4, 5, 6\}$ . Then the PMF is given by

$$p(x) = \begin{cases} \frac{1}{6}, & x \in \{1, 2, 3, 4, 5, 6\} \\ 0, & \text{otherwise} \end{cases}$$

We note here that  $\sum_{x \in \mathbb{Z}} p(x) = 1$ . We note here that PMF takes uniform values for all values of  $X = x$ .

## Uniform Distribution...

## Uniform Distribution...

### Definition: Uniform Distribution

Motivated from the previous example, we now define **uniform distribution** on  $\{1, 2, \dots, n\}$  by

$$p(x) = \begin{cases} \frac{1}{n}, & x \in \{1, 2, \dots, n\} \\ 0, & \text{otherwise} \end{cases}$$

We verify here that  $\sum_{x \in \mathbb{Z}} p(x) = 1$ .



## Bernoulli Distribution

### Bernoulli distribution

A random variable  $X$  is called a **Bernoulli random variable** with parameter  $p$ , denoted by  $X \sim \text{Bernoulli}(p)$ , if its **PMF** is given by

$$P_X(x) = \begin{cases} p & \text{for } x = 1, \\ 1 - p & \text{for } x = 0, \\ 0 & \text{otherwise,} \end{cases}$$

where  $0 < p < 1$ .

## Bernoulli Distribution

### Bernoulli distribution

A random variable  $X$  is called a **Bernoulli random variable** with parameter  $p$ , denoted by  $X \sim \text{Bernoulli}(p)$ , if its **PMF** is given by

$$P_X(x) = \begin{cases} p & \text{for } x = 1, \\ 1 - p & \text{for } x = 0, \\ 0 & \text{otherwise,} \end{cases}$$

where  $0 < p < 1$ .

- This models random experiments that have **two** possible outcomes

## Bernoulli Distribution

### Bernoulli distribution

A random variable  $X$  is called a **Bernoulli random variable** with parameter  $p$ , denoted by  $X \sim \text{Bernoulli}(p)$ , if its **PMF** is given by

$$P_X(x) = \begin{cases} p & \text{for } x = 1, \\ 1 - p & \text{for } x = 0, \\ 0 & \text{otherwise,} \end{cases}$$

where  $0 < p < 1$ .

- This models random experiments that have **two** possible outcomes
- **Example:** You take a pass-fail exam. You either **pass or fail**

## Bernoulli Distribution

### Bernoulli distribution

A random variable  $X$  is called a **Bernoulli random variable** with parameter  $p$ , denoted by  $X \sim \text{Bernoulli}(p)$ , if its **PMF** is given by

$$P_X(x) = \begin{cases} p & \text{for } x = 1, \\ 1 - p & \text{for } x = 0, \\ 0 & \text{otherwise,} \end{cases}$$

where  $0 < p < 1$ .

- This models random experiments that have **two** possible outcomes
- **Example:** You take a pass-fail exam. You either **pass or fail**
- **Example:** A coin is tossed, the outcome is either **heads or tails**

## Geometric Distribution

## Geometric Distribution

### Definition of Geometric Distribution

A random variable  $X$  is called **geometric random variable** with parameter  $p$ , denoted by  $X \sim \text{Geometric}(p)$ , if its **PMF** is given by

$$P_X(k) = \begin{cases} p(1-p)^{k-1}, & \text{for } k = 1, 2, 3, \dots \\ 0, & \text{otherwise,} \end{cases}$$

where  $0 < p < 1$ .

## Geometric Distribution

### Definition of Geometric Distribution

A random variable  $X$  is called **geometric random variable** with parameter  $p$ , denoted by  $X \sim \text{Geometric}(p)$ , if its **PMF** is given by

$$P_X(k) = \begin{cases} p(1-p)^{k-1}, & \text{for } k = 1, 2, 3, \dots \\ 0, & \text{otherwise,} \end{cases}$$

where  $0 < p < 1$ .

- **Example:** Suppose we have an unfair coin with  $P(H) = p$ .

## Geometric Distribution

### Definition of Geometric Distribution

A random variable  $X$  is called **geometric random variable** with parameter  $p$ , denoted by  $X \sim \text{Geometric}(p)$ , if its **PMF** is given by

$$P_X(k) = \begin{cases} p(1-p)^{k-1}, & \text{for } k = 1, 2, 3, \dots \\ 0, & \text{otherwise,} \end{cases}$$

where  $0 < p < 1$ .

- **Example:** Suppose we have an unfair coin with  $P(H) = p$ . We **toss the coin until we obtain first heads**.



## Geometric Distribution

### Definition of Geometric Distribution

A random variable  $X$  is called **geometric random variable** with parameter  $p$ , denoted by  $X \sim \text{Geometric}(p)$ , if its **PMF** is given by

$$P_X(k) = \begin{cases} p(1-p)^{k-1}, & \text{for } k = 1, 2, 3, \dots \\ 0, & \text{otherwise,} \end{cases}$$

where  $0 < p < 1$ .

- **Example:** Suppose we have an unfair coin with  $P(H) = p$ . We **toss the coin until we obtain first heads**. Let RV  $X$  be the total number of tosses.

## Geometric Distribution

### Definition of Geometric Distribution

A random variable  $X$  is called **geometric random variable** with parameter  $p$ , denoted by  $X \sim \text{Geometric}(p)$ , if its **PMF** is given by

$$P_X(k) = \begin{cases} p(1-p)^{k-1}, & \text{for } k = 1, 2, 3, \dots \\ 0, & \text{otherwise,} \end{cases}$$

where  $0 < p < 1$ .

- **Example:** Suppose we have an unfair coin with  $P(H) = p$ . We **toss the coin until we obtain first heads**. Let RV  $X$  be the total number of tosses. Then  $X$  have **geometric** distribution.

## Geometric Distribution

### Definition of Geometric Distribution

A random variable  $X$  is called **geometric random variable** with parameter  $p$ , denoted by  $X \sim \text{Geometric}(p)$ , if its **PMF** is given by

$$P_X(k) = \begin{cases} p(1-p)^{k-1}, & \text{for } k = 1, 2, 3, \dots \\ 0, & \text{otherwise,} \end{cases}$$

where  $0 < p < 1$ .

- **Example:** Suppose we have an unfair coin with  $P(H) = p$ . We **toss the coin until we obtain first heads**. Let RV  $X$  be the total number of tosses. Then  $X$  have **geometric** distribution.
- **Caution:** Some books define geometric random variable  $X$  as total number of failures before observing first success. Then

## Geometric Distribution

### Definition of Geometric Distribution

A random variable  $X$  is called **geometric random variable** with parameter  $p$ , denoted by  $X \sim \text{Geometric}(p)$ , if its **PMF** is given by

$$P_X(k) = \begin{cases} p(1-p)^{k-1}, & \text{for } k = 1, 2, 3, \dots \\ 0, & \text{otherwise,} \end{cases}$$

where  $0 < p < 1$ .

- **Example:** Suppose we have an unfair coin with  $P(H) = p$ . We **toss the coin until we obtain first heads**. Let RV  $X$  be the total number of tosses. Then  $X$  have **geometric** distribution.
- **Caution:** Some books define geometric random variable  $X$  as total number of failures before observing first success. Then

$$P_X(k) = \begin{cases} p(1-p)^k, & \text{for } k = 1, 2, 3, \dots \\ 0, & \text{otherwise,} \end{cases}$$

## Binomial Distribution

## Binomial Distribution

### Definition of Binomial Distribution

A random variable  $X$  is called **Binomial random variable** with parameters  $n$  and  $p$ , denoted by  $X \sim \text{Binomial}(n, p)$ , if the **PMF** is given by

## Binomial Distribution

### Definition of Binomial Distribution

A random variable  $X$  is called **Binomial random variable** with parameters  $n$  and  $p$ , denoted by  $X \sim \text{Binomial}(n, p)$ , if the **PMF** is given by

$$P_X(k) = \begin{cases} \binom{n}{k} p^k (1-p)^{n-k} & \text{for } k = 0, 1, 2, \dots, n \\ 0 & \text{otherwise,} \end{cases}$$

where  $0 < p < 1$ .

## Binomial Distribution

### Definition of Binomial Distribution

A random variable  $X$  is called **Binomial random variable** with parameters  $n$  and  $p$ , denoted by  $X \sim \text{Binomial}(n, p)$ , if the **PMF** is given by

$$P_X(k) = \begin{cases} \binom{n}{k} p^k (1-p)^{n-k} & \text{for } k = 0, 1, 2, \dots, n \\ 0 & \text{otherwise,} \end{cases}$$

where  $0 < p < 1$ .

- **Example:** Consider an unfair coin with  $P(H) = p$ . Consider tossing the coin  $n$  times and let  $X$  be the total number of heads we observe. Then  $X$  is a **Binomial** with parameters  $n$  and  $p$ .



## Binomial Distribution

### Definition of Binomial Distribution

A random variable  $X$  is called **Binomial random variable** with parameters  $n$  and  $p$ , denoted by  $X \sim \text{Binomial}(n, p)$ , if the **PMF** is given by

$$P_X(k) = \begin{cases} \binom{n}{k} p^k (1-p)^{n-k} & \text{for } k = 0, 1, 2, \dots, n \\ 0 & \text{otherwise,} \end{cases}$$

where  $0 < p < 1$ .

- **Example:** Consider an unfair coin with  $P(H) = p$ . Consider tossing the coin  $n$  times and let  $X$  be the total number of heads we observe. Then  $X$  is a **Binomial** with parameters  $n$  and  $p$ .
- $\text{Binomial}(n, p)$  is a sum of  $n$  independent **Bernoulli**( $p$ ) random variables.

## Binomial Distribution

### Definition of Binomial Distribution

A random variable  $X$  is called **Binomial random variable** with parameters  $n$  and  $p$ , denoted by  $X \sim \text{Binomial}(n, p)$ , if the **PMF** is given by

$$P_X(k) = \begin{cases} \binom{n}{k} p^k (1-p)^{n-k} & \text{for } k = 0, 1, 2, \dots, n \\ 0 & \text{otherwise,} \end{cases}$$

where  $0 < p < 1$ .

- **Example:** Consider an unfair coin with  $P(H) = p$ . Consider tossing the coin  $n$  times and let  $X$  be the total number of heads we observe. Then  $X$  is a **Binomial** with parameters  $n$  and  $p$ .
- $\text{Binomial}(n, p)$  is a sum of  $n$  independent **Bernoulli**( $p$ ) random variables.
- If  $X_1, X_2, \dots, X_n$  are independent **Bernoulli**( $p$ ) random variables, then  $X = X_1 + \dots + X_n$  has **Bernoulli**( $n, p$ ) distribution.

## Binomial Distribution

### Definition of Binomial Distribution

A random variable  $X$  is called **Binomial random variable** with parameters  $n$  and  $p$ , denoted by  $X \sim \text{Binomial}(n, p)$ , if the **PMF** is given by

$$P_X(k) = \begin{cases} \binom{n}{k} p^k (1-p)^{n-k} & \text{for } k = 0, 1, 2, \dots, n \\ 0 & \text{otherwise,} \end{cases}$$

where  $0 < p < 1$ .

- **Example:** Consider an unfair coin with  $P(H) = p$ . Consider tossing the coin  $n$  times and let  $X$  be the total number of heads we observe. Then  $X$  is a **Binomial** with parameters  $n$  and  $p$ .
- $\text{Binomial}(n, p)$  is a sum of  $n$  independent **Bernoulli**( $p$ ) random variables.
- If  $X_1, X_2, \dots, X_n$  are independent **Bernoulli**( $p$ ) random variables, then  $X = X_1 + \dots + X_n$  has **Bernoulli**( $n, p$ ) distribution.
- We verify that  $\sum_{x \in \mathbb{Z}} P_X(x) = \sum_{x=0}^n \binom{n}{x} p^x (1-p)^{n-x} = 1$

## Example

### Example

Let  $X \sim \text{Binomial}(n, p)$  and  $Y \sim \text{Binomial}(m, p)$  be two independent random variables. We define a random variable  $Z = X + Y$ . What is the PMF of  $Z$ ?

Scratch Space...

Scratch Space...

## Motivation for Poisson Distribution...

## Motivation for Poisson Distribution...

- Imagine you are an Uber driver




## Motivation for Poisson Distribution...

- Imagine you are an Uber driver
- To maximize profit, you want to remain close to an area with more requests
- How can you use probability to predict requests for car ride?


## Motivation for Poisson Distribution...

- Imagine you are an Uber driver
- To maximize profit, you want to remain close to an area with more requests
- How can you use probability to predict requests for car ride?



## Motivation for Poisson Distribution...

- Imagine you are an Uber driver
- To maximize profit, you want to remain close to an area with more requests
- How can you use probability to predict requests for car ride?



### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

## Introduce Poisson Using an Example...

## Introduce Poisson Using an Example...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- What do we know for this problem?

## Introduce Poisson Using an Example...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- What do we know for this problem? There are requests for car ride!

## Introduce Poisson Using an Example...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- What do we know for this problem? There are requests for car ride!
- There are two possibilities: request (1) or no request (0):

## Introduce Poisson Using an Example...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- What do we know for this problem? There are requests for car ride!
- There are two possibilities: request (1) or no request (0): binary choices



## Introduce Poisson Using an Example...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- What do we know for this problem? There are requests for car ride!
- There are two possibilities: request (1) or no request (0): binary choices
- Sounds familiar? Yes!

## Introduce Poisson Using an Example...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- What do we know for this problem? There are requests for car ride!
- There are two possibilities: request (1) or no request (0): binary choices
- Sounds familiar? **Yes!** Recall: How many heads in  $n$  trials?

## Introduce Poisson Using an Example...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- What do we know for this problem? There are requests for car ride!
- There are two possibilities: request (1) or no request (0): binary choices
- Sounds familiar? **Yes!** Recall: How many heads in  $n$  trials?
- The problem here is: we don't have  $n$  trials, instead a time interval! So?

## Introduce Poisson Using an Example...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- What do we know for this problem? There are requests for car ride!
- There are two possibilities: request (1) or no request (0): binary choices
- Sounds familiar? **Yes!** Recall: How many heads in  $n$  trials?
- The problem here is: we don't have  $n$  trials, instead a time interval! So?
- **Key idea:** Want to use discrete distribution, so anyhow convert time to discrete intervals

## Introduce Poisson Using an Example...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- What do we know for this problem? There are requests for car ride!
- There are two possibilities: request (1) or no request (0): binary choices
- Sounds familiar? **Yes!** Recall: How many heads in  $n$  trials?
- The problem here is: we don't have  $n$  trials, instead a time interval! So?
- **Key idea:** Want to use discrete distribution, so anyhow convert time to discrete intervals

0	0	1	0	1	...	...	0	1	0	1	1	0
---	---	---	---	---	-----	-----	---	---	---	---	---	---

## Introduce Poisson Using an Example...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- What do we know for this problem? There are requests for car ride!
- There are two possibilities: request (1) or no request (0): binary choices
- Sounds familiar? **Yes!** Recall: How many heads in  $n$  trials?
- The problem here is: we don't have  $n$  trials, instead a time interval! So?
- **Key idea:** Want to use discrete distribution, so anyhow convert time to discrete intervals

0	0	1	0	1	...	...	0	1	0	1	1	0
---	---	---	---	---	-----	-----	---	---	---	---	---	---

- break a minute into 60 seconds;

## Introduce Poisson Using an Example...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- What do we know for this problem? There are requests for car ride!
- There are two possibilities: request (1) or no request (0): binary choices
- Sounds familiar? **Yes!** Recall: How many heads in  $n$  trials?
- The problem here is: we don't have  $n$  trials, instead a time interval! So?
- **Key idea:** Want to use discrete distribution, so anyhow convert time to discrete intervals

0	0	1	0	1	...	...	0	1	0	1	1	0
---	---	---	---	---	-----	-----	---	---	---	---	---	---

- break a minute into 60 seconds; each second is independent trial with request or no request

## Introduce Poisson Using an Example...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- What do we know for this problem? There are requests for car ride!
- There are two possibilities: request (1) or no request (0): binary choices
- Sounds familiar? **Yes!** Recall: How many heads in  $n$  trials?
- The problem here is: we don't have  $n$  trials, instead a time interval! So?
- **Key idea:** Want to use discrete distribution, so anyhow convert time to discrete intervals

0	0	1	0	1	...	...	0	1	0	1	1	0
---	---	---	---	---	-----	-----	---	---	---	---	---	---

- break a minute into 60 seconds; each second is independent trial with request or no request
- Let  $X$  = number of requests in a minute



## Introduce Poisson Using an Example...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- What do we know for this problem? There are requests for car ride!
- There are two possibilities: request (1) or no request (0): binary choices
- Sounds familiar? **Yes!** Recall: How many heads in  $n$  trials?
- The problem here is: we don't have  $n$  trials, instead a time interval! So?
- **Key idea:** Want to use discrete distribution, so anyhow convert time to discrete intervals

0	0	1	0	1	...	...	0	1	0	1	1	0
---	---	---	---	---	-----	-----	---	---	---	---	---	---

- break a minute into 60 seconds; each second is independent trial with request or no request
- Let  $X$  = number of requests in a minute
- $E[X] = \lambda = 5 = np$ , where  $p$  is the probability of request

## Introduce Poisson Using an Example...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- What do we know for this problem? There are requests for car ride!
- There are two possibilities: request (1) or no request (0): binary choices
- Sounds familiar? **Yes!** Recall: How many heads in  $n$  trials?
- The problem here is: we don't have  $n$  trials, instead a time interval! So?
- **Key idea:** Want to use discrete distribution, so anyhow convert time to discrete intervals

0	0	1	0	1	...	...	0	1	0	1	1	0
---	---	---	---	---	-----	-----	---	---	---	---	---	---

- break a minute into 60 seconds; each second is independent trial with request or no request
- Let  $X$  = number of requests in a minute
- $E[X] = \lambda = 5 = np$ , where  $p$  is the probability of request
- We now identify this problem as  $X \sim \text{Binomial}(n = 60, p = 5/60)$

## Introduce Poisson Using an Example...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- What do we know for this problem? There are requests for car ride!
- There are two possibilities: request (1) or no request (0): binary choices
- Sounds familiar? **Yes!** Recall: How many heads in  $n$  trials?
- The problem here is: we don't have  $n$  trials, instead a time interval! So?
- **Key idea:** Want to use discrete distribution, so anyhow convert time to discrete intervals

0	0	1	0	1	...	...	0	1	0	1	1	0
---	---	---	---	---	-----	-----	---	---	---	---	---	---

- break a minute into 60 seconds; each second is independent trial with request or no request
- Let  $X$  = number of requests in a minute
- $E[X] = \lambda = 5 = np$ , where  $p$  is the probability of request
- We now identify this problem as  $X \sim \text{Binomial}(n = 60, p = 5/60)$
- $P(X = k) = \binom{60}{k} \left(\frac{5}{60}\right)^k \left(1 - \frac{5}{60}\right)^{n-k}$

## Motivation for Poisson Distribution...

## Motivation for Poisson Distribution...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- We identified this problem as  $X \sim \text{Binomial}(n = 60, p = 5/60)$

## Motivation for Poisson Distribution...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- We identified this problem as  $X \sim \text{Binomial}(n = 60, p = 5/60)$
- $P(X = k) = \binom{60}{k} \left(\frac{5}{60}\right)^k \left(1 - \frac{5}{60}\right)^{n-k}$
- There is a problem!

## Motivation for Poisson Distribution...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- We identified this problem as  $X \sim \text{Binomial}(n = 60, p = 5/60)$
- $P(X = k) = \binom{60}{k} \left(\frac{5}{60}\right)^k \left(1 - \frac{5}{60}\right)^{n-k}$
- There is a problem! What is that?

## Motivation for Poisson Distribution...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- We identified this problem as  $X \sim \text{Binomial}(n = 60, p = 5/60)$
- $P(X = k) = \binom{60}{k} \left(\frac{5}{60}\right)^k \left(1 - \frac{5}{60}\right)^{n-k}$
- There is a problem! What is that? Time is continuous!



## Motivation for Poisson Distribution...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- We identified this problem as  $X \sim \text{Binomial}(n = 60, p = 5/60)$
- $P(X = k) = \binom{60}{k} \left(\frac{5}{60}\right)^k \left(1 - \frac{5}{60}\right)^{n-k}$
- There is a problem! What is that? Time is continuous!
- Since time is continuous, no guarantee that there will be only one request per second

## Motivation for Poisson Distribution...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- We identified this problem as  $X \sim \text{Binomial}(n = 60, p = 5/60)$
- $P(X = k) = \binom{60}{k} \left(\frac{5}{60}\right)^k \left(1 - \frac{5}{60}\right)^{n-k}$
- There is a problem! What is that? Time is continuous!
- Since time is continuous, no guarantee that there will be only one request per second
- What do we do now?

## Motivation for Poisson Distribution...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- We identified this problem as  $X \sim \text{Binomial}(n = 60, p = 5/60)$
- $P(X = k) = \binom{60}{k} \left(\frac{5}{60}\right)^k \left(1 - \frac{5}{60}\right)^{n-k}$
- There is a problem! What is that? Time is continuous!
- Since time is continuous, no guarantee that there will be only one request per second
- What do we do now? We can make the interval smaller, take time intervals as milliseconds?

## Motivation for Poisson Distribution...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- We identified this problem as  $X \sim \text{Binomial}(n = 60, p = 5/60)$
- $P(X = k) = \binom{60}{k} \left(\frac{5}{60}\right)^k \left(1 - \frac{5}{60}\right)^{n-k}$
- There is a problem! What is that? Time is continuous!
- Since time is continuous, no guarantee that there will be only one request per second
- What do we do now? We can make the interval smaller, take time intervals as milliseconds?
- But there is still no guarantee that there will be only one request per millisecond

## Motivation for Poisson Distribution...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- We identified this problem as  $X \sim \text{Binomial}(n = 60, p = 5/60)$
- $P(X = k) = \binom{60}{k} \left(\frac{5}{60}\right)^k \left(1 - \frac{5}{60}\right)^{n-k}$
- There is a problem! What is that? Time is continuous!
- Since time is continuous, no guarantee that there will be only one request per second
- What do we do now? We can make the interval smaller, take time intervals as milliseconds?
- But there is still no guarantee that there will be only one request per millisecond
- What should we do next?

## Motivation for Poisson Distribution...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- We identified this problem as  $X \sim \text{Binomial}(n = 60, p = 5/60)$
- $P(X = k) = \binom{60}{k} \left(\frac{5}{60}\right)^k \left(1 - \frac{5}{60}\right)^{n-k}$
- There is a problem! What is that? Time is continuous!
- Since time is continuous, no guarantee that there will be only one request per second
- What do we do now? We can make the interval smaller, take time intervals as milliseconds?
- But there is still no guarantee that there will be only one request per millisecond
- What should we do next? Here comes calculus...

## Motivation for Poisson Distribution...

### Example

Given that there are on average 5 requests for car ride per minute from certain area. What is the probability of  $k$  requests from this area in next 1 minute?

- We identified this problem as  $X \sim \text{Binomial}(n = 60, p = 5/60)$
- $P(X = k) = \binom{60}{k} \left(\frac{5}{60}\right)^k \left(1 - \frac{5}{60}\right)^{n-k}$
- There is a problem! What is that? Time is continuous!
- Since time is continuous, no guarantee that there will be only one request per second
- What do we do now? We can make the interval smaller, take time intervals as milliseconds?
- But there is still no guarantee that there will be only one request per millisecond
- What should we do next? Here comes calculus... to go from discrete to continuous...

**Binomial in the Limit is Poisson Distribution...**



## Binomial in the Limit is Poisson Distribution...

Derivation:

## Definition of Poisson Distribution...

## Definition of Poisson Distribution...

### Definition of Poisson

A random variable  $X$  is said to be a Poisson random variable with parameter  $\lambda$ , shown as  $X \sim \text{Poisson}(\lambda)$ , if its range is  $R_X = \{0, 1, 2, \dots, \}$ , and its PMF is given by

$$P_X(k) = \begin{cases} \frac{e^{-\lambda} \lambda^k}{k!}, & k \in R_X \\ 0 & \text{otherwise} \end{cases}$$



- Simeon-Denis Poisson, was a French mathematician (1781-1840)
- He published his first paper at 18, became professor at 21
- He published over 300 papers

## Outline

- ① Random Variables
- ② Special Distributions
- ③ Examples of Distributions
- ④ Expectations of Some Distributions
- ⑤ Variance and Standard Deviation: Understand Variability in Data
- ⑥ Higher Order Moments and Moment Generating Function

## Example of Poisson Distribution...

## Example of Poisson Distribution...

### Example

We want to send a bit string of length  $n = 10^4$ .

## Example of Poisson Distribution...

### Example

We want to send a bit string of length  $n = 10^4$ . The probability that a bit can be corrupted is  $p = 10^{-6}$ .

## Example of Poisson Distribution...

### Example

We want to send a bit string of length  $n = 10^4$ . The probability that a bit can be corrupted is  $p = 10^{-6}$ . What is the probability that the message will arrive **uncorrupted**?



## Example of Poisson Distribution...

### Example

We want to send a bit string of length  $n = 10^4$ . The probability that a bit can be corrupted is  $p = 10^{-6}$ . What is the probability that the message will arrive **uncorrupted**?

- We can solve this using a Poisson with  $\lambda = np = 10^4 10^{-6}$

## Example of Poisson Distribution...

### Example

We want to send a bit string of length  $n = 10^4$ . The probability that a bit can be corrupted is  $p = 10^{-6}$ . What is the probability that the message will arrive **uncorrupted**?

- We can solve this using a Poisson with  $\lambda = np = 10^4 10^{-6}$
- With  $X \sim \text{Poisson}(0.01)$  be the number of corrupted bits

## Example of Poisson Distribution...

### Example

We want to send a bit string of length  $n = 10^4$ . The probability that a bit can be corrupted is  $p = 10^{-6}$ . What is the probability that the message will arrive **uncorrupted**?

- We can solve this using a Poisson with  $\lambda = np = 10^4 10^{-6}$
- With  $X \sim \text{Poisson}(0.01)$  be the number of corrupted bits
- Using the PMF for Poisson

## Example of Poisson Distribution...

### Example

We want to send a bit string of length  $n = 10^4$ . The probability that a bit can be corrupted is  $p = 10^{-6}$ . What is the probability that the message will arrive **uncorrupted**?

- We can solve this using a Poisson with  $\lambda = np = 10^4 10^{-6}$
- With  $X \sim \text{Poisson}(0.01)$  be the number of corrupted bits
- Using the PMF for Poisson

$$\begin{aligned} P(X = 0) &= \frac{\lambda^i}{i!} e^{-\lambda} \\ &= \frac{0.01^0}{0!} e^{-0.01} \\ &\approx 0.9900498 \end{aligned}$$

- We could have solved this using Binomial  $X \sim \text{Binomial}(10^4, 10^{-6})$ . This may have been harder to compute!

## Example of Poisson Distribution...

### Example

We want to send a bit string of length  $n = 10^4$ . The probability that a bit can be corrupted is  $p = 10^{-6}$ . What is the probability that the message will arrive **uncorrupted**?

- We can solve this using a Poisson with  $\lambda = np = 10^4 10^{-6}$
- With  $X \sim \text{Poisson}(0.01)$  be the number of corrupted bits
- Using the PMF for Poisson

$$\begin{aligned} P(X = 0) &= \frac{\lambda^i}{i!} e^{-\lambda} \\ &= \frac{0.01^0}{0!} e^{-0.01} \\ &\approx 0.9900498 \end{aligned}$$

- We could have solved this using Binomial  $X \sim \text{Binomial}(10^4, 10^{-6})$ . This may have been harder to compute!
- When  $n$  large, and  $p$  small: can use Poisson!

## Another Example of Poisson Distribution...

## Another Example of Poisson Distribution...

### Example of Poisson Distribution

The Poisson distribution is often used to model the number of events that occur independently at any time in an interval of time or space, with a **constant** average rate.

## Another Example of Poisson Distribution...

### Example of Poisson Distribution

The Poisson distribution is often used to model the number of events that occur independently at any time in an interval of time or space, with a **constant** average rate. Earthquakes are a good example of this.



## Another Example of Poisson Distribution...

### Example of Poisson Distribution

The Poisson distribution is often used to model the number of events that occur independently at any time in an interval of time or space, with a **constant** average rate. Earthquakes are a good example of this. Suppose there are an average of 2.8 major earthquakes in the world each year.

## Another Example of Poisson Distribution...

### Example of Poisson Distribution

The Poisson distribution is often used to model the number of events that occur independently at any time in an interval of time or space, with a **constant** average rate. Earthquakes are a good example of this. Suppose there are an average of 2.8 major earthquakes in the world each year. What is the probability of getting more than one major earthquake next year?

## Another Example of Poisson Distribution...

### Example of Poisson Distribution

The Poisson distribution is often used to model the number of events that occur independently at any time in an interval of time or space, with a **constant** average rate. Earthquakes are a good example of this. Suppose there are an average of 2.8 major earthquakes in the world each year. What is the probability of getting more than one major earthquake next year?

- Let  $X \sim \text{Poisson}(2.8)$  be the number of major earthquakes next year. We want  $P(X > 1)$ .

## Another Example of Poisson Distribution...

### Example of Poisson Distribution

The Poisson distribution is often used to model the number of events that occur independently at any time in an interval of time or space, with a **constant** average rate. Earthquakes are a good example of this. Suppose there are an average of 2.8 major earthquakes in the world each year. What is the probability of getting more than one major earthquake next year?

- Let  $X \sim \text{Poisson}(2.8)$  be the number of major earthquakes next year. We want  $P(X > 1)$ .
- We can use

$$P(X > 1) = 1 - P(X = 0) - P(X = 1).$$

Using PMF for Poisson

## Another Example of Poisson Distribution...

### Example of Poisson Distribution

The Poisson distribution is often used to model the number of events that occur independently at any time in an interval of time or space, with a **constant** average rate. Earthquakes are a good example of this. Suppose there are an average of 2.8 major earthquakes in the world each year. What is the probability of getting more than one major earthquake next year?

- Let  $X \sim \text{Poisson}(2.8)$  be the number of major earthquakes next year. We want  $P(X > 1)$ .
- We can use

$$P(X > 1) = 1 - P(X = 0) - P(X = 1).$$

Using PMF for Poisson

$$\begin{aligned} P(X > 1) &= 1 - P(X = 0) - P(X = 1) \\ &= 1 - e^{-2.8} \frac{2.8^0}{0!} - e^{-2.8} \frac{2.8^1}{1!} \\ &= 1 - e^{-2.8} - 2.8e^{-2.8} \\ &\approx 1 - 0.06 - 0.17 = 0.77 \end{aligned}$$

## Outline

- ① Random Variables
- ② Special Distributions
- ③ Examples of Distributions
- ④ Expectations of Some Distributions
- ⑤ Variance and Standard Deviation: Understand Variability in Data
- ⑥ Higher Order Moments and Moment Generating Function

## Expectation of Poisson Distribution...

## Expectation of Poisson Distribution...

### Expectation of Poisson

Let  $X \sim \text{Poisson}(\lambda)$ , with PMF given by

$$P_X(k) = \begin{cases} \frac{e^{-\lambda} \lambda^k}{k!}, & k \in R_X \\ 0 & \text{otherwise} \end{cases}$$

Show that the expectation  $E[X] = \lambda$ .



## Expectation of Poisson Distribution...

### Expectation of Poisson

Let  $X \sim \text{Poisson}(\lambda)$ , with PMF given by

$$P_X(k) = \begin{cases} \frac{e^{-\lambda} \lambda^k}{k!}, & k \in R_X \\ 0 & \text{otherwise} \end{cases}$$

Show that the expectation  $E[X] = \lambda$ .

### Proof

$$\begin{aligned} E[X] &= \sum_{x=0}^{\infty} x \frac{e^{-\lambda} \lambda^x}{x!} = \sum_{x=1}^{\infty} x \frac{e^{-\lambda} \lambda^x}{x!} = e^{-\lambda} \sum_{x=1}^{\infty} \frac{\lambda^x}{(x-1)!} \\ &= e^{-\lambda} \sum_{z=0}^{\infty} \frac{\lambda^{z+1}}{z!} = \lambda e^{-\lambda} \sum_{z=0}^{\infty} \frac{\lambda^z}{z!} = \lambda. \end{aligned}$$

## Expectation of Binomial Distribution...

## Expectation of Binomial Distribution...

### Expectation of a Binomial Distribution

Show that  $E[X] = np$ , where  $X \sim \text{Binomial}(n, p)$ ,  $0 < p < 1$  with PMF given as follows

$$P_X(k) = \begin{cases} \binom{n}{k} p^k (1-p)^{n-k} & \text{for } k = 0, 1, 2, \dots, n \\ 0 & \text{otherwise,} \end{cases}$$

## Expectation of Binomial Distribution...

### Expectation of a Binomial Distribution

Show that  $E[X] = np$ , where  $X \sim \text{Binomial}(n, p)$ ,  $0 < p < 1$  with PMF given as follows

$$P_X(k) = \begin{cases} \binom{n}{k} p^k (1-p)^{n-k} & \text{for } k = 0, 1, 2, \dots, n \\ 0 & \text{otherwise,} \end{cases}$$

### Proof

$$E[X] = \sum_{x=0}^n x \binom{n}{x} p^x q^{n-x} = \sum_{x=1}^n x \binom{n}{x} p^x q^{n-x}.$$

$$\text{For } 0 < x \leq n, \quad x \binom{n}{x} = x \frac{n!}{(n-x)!x!} = \frac{n!}{(n-x)!(x-1)!} = n \binom{n-1}{x-1},$$

$$\Rightarrow E[X] = \sum_{x=1}^n n \binom{n-1}{x-1} p^x q^{n-x} = \sum_{z=0}^{n-1} n \binom{n-1}{z} p^{z+1} q^{n-1-z} = np.$$

## Outline

- ① Random Variables
- ② Special Distributions
- ③ Examples of Distributions
- ④ Expectations of Some Distributions
- ⑤ Variance and Standard Deviation: Understand Variability in Data
- ⑥ Higher Order Moments and Moment Generating Function

## Variance

### Motivation for Variance

- The expectation value  $E(X)$  (also called the mean) of a discrete random variable is a rather coarse measure of how  $X$  is distributed. For example, consider the following three situations:

## Variance

### Motivation for Variance

- The expectation value  $E(X)$  (also called the mean) of a discrete random variable is a rather coarse measure of how  $X$  is distributed. For example, consider the following three situations:
  - You get Rs 1000

## Variance

### Motivation for Variance

- The expectation value  $E(X)$  (also called the mean) of a discrete random variable is a rather coarse measure of how  $X$  is distributed. For example, consider the following three situations:
  - You get Rs 1000
  - A fair coin is tossed, and if it is head you get Rs 2000



## Variance

### Motivation for Variance

- The expectation value  $E(X)$  (also called the mean) of a discrete random variable is a rather coarse measure of how  $X$  is distributed. For example, consider the following three situations:
  - You get Rs 1000
  - A fair coin is tossed, and if it is head you get Rs 2000
  - A number is chosen from 1 to 1000 and if you can guess this, you get Rs 1 Million

## Variance

### Motivation for Variance

- The expectation value  $E(X)$  (also called the mean) of a discrete random variable is a rather coarse measure of how  $X$  is distributed. For example, consider the following three situations:
  - You get Rs 1000
  - A fair coin is tossed, and if it is head you get Rs 2000
  - A number is chosen from 1 to 1000 and if you can guess this, you get Rs 1 Million
- Let  $X$  be the discrete random variable corresponding to your winnings. In all three cases,  $E(X) = \text{Rs } 1000$ , but you will agree that your chances of actually getting any money are quite different in all three cases.

## Variance

### Motivation for Variance

- The expectation value  $E(X)$  (also called the mean) of a discrete random variable is a rather coarse measure of how  $X$  is distributed. For example, consider the following three situations:
  - You get Rs 1000
  - A fair coin is tossed, and if it is head you get Rs 2000
  - A number is chosen from 1 to 1000 and if you can guess this, you get Rs 1 Million
- Let  $X$  be the discrete random variable corresponding to your winnings. In all three cases,  $E(X) = \text{Rs } 1000$ , but you will agree that your chances of actually getting any money are quite different in all three cases.
- The three cases differ in the spread of PMF. This can be measured by Variance.

## Variance

### Motivation for Variance

- The expectation value  $E(X)$  (also called the mean) of a discrete random variable is a rather coarse measure of how  $X$  is distributed. For example, consider the following three situations:
  - You get Rs 1000
  - A fair coin is tossed, and if it is head you get Rs 2000
  - A number is chosen from 1 to 1000 and if you can guess this, you get Rs 1 Million
- Let  $X$  be the discrete random variable corresponding to your winnings. In all three cases,  $E(X) = \text{Rs } 1000$ , but you will agree that **your chances of actually getting any money are quite different in all three cases.**
- The three cases differ in the spread of PMF. This can be measured by **Variance.**
- Recall Saint Petersburg Paradox! High Risk High Reward!

## Variance and Standard Deviation...

## Variance and Standard Deviation...

### Variance

Let  $\mu = E[X]$ . The variance  $\text{Var}(X)$  of  $X$  is defined by

## Variance and Standard Deviation...

### Variance

Let  $\mu = E[X]$ . The **variance**  $\text{Var}(X)$  of  $X$  is defined by

$$\text{Var}(X) = E((X - \mu)^2) = \sum_x (x - \mu)^2 f_X(x)$$

## Variance and Standard Deviation...

### Variance

Let  $\mu = E[X]$ . The **variance**  $\text{Var}(X)$  of  $X$  is defined by

$$\text{Var}(X) = E((X - \mu)^2) = \sum_x (x - \mu)^2 f_X(x)$$

The **positive** square root is called the **standard deviation** and it is usually denoted by  $\sigma$ , hence



## Variance and Standard Deviation...

### Variance

Let  $\mu = E[X]$ . The **variance**  $\text{Var}(X)$  of  $X$  is defined by

$$\text{Var}(X) = E((X - \mu)^2) = \sum_x (x - \mu)^2 f_X(x)$$

The **positive** square root is called the **standard deviation** and it is usually denoted by  $\sigma$ , hence

$$\sigma(X) = \sqrt{\sum_x (x - \mu)^2 f_X(x)} = \sqrt{\text{Var}(X)}.$$

## Example of Computing Variance...

## Example of Computing Variance...

### Example

Let  $X$  be the value on one roll of a 6-sided die. Recall that  $E[X] = 7/2$ . What is  $\text{Var}(X)$ ?

## Variance

### Motivation for Variance

- The expectation value  $E(X)$  (also called the mean) of a discrete random variable is a rather coarse measure of how  $X$  is distributed. For example, consider the following three situations:

## Variance

### Motivation for Variance

- The **expectation value**  $E(X)$  (also called the mean) of a discrete random variable is a rather **coarse** measure of how  $X$  is distributed. For example, consider the following three situations:
  - You get Rs 1000

## Variance

### Motivation for Variance

- The **expectation value**  $E(X)$  (also called the mean) of a discrete random variable is a rather **coarse** measure of how  $X$  is distributed. For example, consider the following three situations:
  - You get Rs 1000
  - A fair coin is tossed, and if it is head you get Rs 2000

## Variance

### Motivation for Variance

- The **expectation value**  $E(X)$  (also called the mean) of a discrete random variable is a rather **coarse** measure of how  $X$  is distributed. For example, consider the following three situations:
  - You get Rs 1000
  - A fair coin is tossed, and if it is head you get Rs 2000
  - A number is chosen from 1 to 1000 and if you can guess this, you get Rs 1 Million

## Variance

### Motivation for Variance

- The **expectation value**  $E(X)$  (also called the mean) of a discrete random variable is a rather **coarse** measure of how  $X$  is distributed. For example, consider the following three situations:
  - You get Rs 1000
  - A fair coin is tossed, and if it is head you get Rs 2000
  - A number is chosen from 1 to 1000 and if you can guess this, you get Rs 1 Million



## Variance

### Motivation for Variance

- The **expectation value**  $E(X)$  (also called the mean) of a discrete random variable is a rather **coarse** measure of how  $X$  is distributed. For example, consider the following three situations:
  - You get Rs 1000
  - A fair coin is tossed, and if it is head you get Rs 2000
  - A number is chosen from 1 to 1000 and if you can guess this, you get Rs 1 Million

### Calculate Variance

- **First case:** There is only one outcome and it is the mean, variance is 0

## Variance

### Motivation for Variance

- The **expectation value**  $E(X)$  (also called the mean) of a discrete random variable is a rather **coarse** measure of how  $X$  is distributed. For example, consider the following three situations:
  - You get Rs 1000
  - A fair coin is tossed, and if it is head you get Rs 2000
  - A number is chosen from 1 to 1000 and if you can guess this, you get Rs 1 Million

### Calculate Variance

- **First case:** There is only one outcome and it is the mean, variance is 0
- **Second case:**  $\text{Var}(X) = \frac{1}{2}(2000 - 1000)^2 + \frac{1}{2}(0 - 1000)^2 = 10^6$

## Variance

### Motivation for Variance

- The **expectation value**  $E(X)$  (also called the mean) of a discrete random variable is a rather **coarse** measure of how  $X$  is distributed. For example, consider the following three situations:
  - You get Rs 1000
  - A fair coin is tossed, and if it is head you get Rs 2000
  - A number is chosen from 1 to 1000 and if you can guess this, you get Rs 1 Million

### Calculate Variance

- **First case:** There is only one outcome and it is the mean, variance is 0
- **Second case:**  $\text{Var}(X) = \frac{1}{2}(2000 - 1000)^2 + \frac{1}{2}(0 - 1000)^2 = 10^6$
- **Third case:**  $\text{Var}(X) = 10^{-3}(10^6 - 10^3)^2 + 999 \times 10^{-3}(0 - 10^3)^2 \approx 10^9$

## Remarks on Expectation, Variance, Standard Deviation...

## Remarks on Expectation, Variance, Standard Deviation...

### Example

If you are told that the average starting salary for someone working at Company Statistix is Rs 70,000, you may think, "Wow! That's great."

## Remarks on Expectation, Variance, Standard Deviation...

### Example

If you are told that the average starting salary for someone working at Company Statistix is Rs 70,000, you may think, "Wow! That's great." But if the standard deviation for starting salaries at Company Statistix is Rs 20,000, that's a lot of variation in terms of how much money you can make,

## Remarks on Expectation, Variance, Standard Deviation...

### Example

If you are told that the average starting salary for someone working at Company Statistix is Rs 70,000, you may think, "Wow! That's great." But if the standard deviation for starting salaries at Company Statistix is Rs 20,000, that's a lot of variation in terms of how much money you can make, so the average starting salary of Rs 70,000 isn't as informative in the end, is it?

## Remarks on Expectation, Variance, Standard Deviation...

### Example

If you are told that the average starting salary for someone working at Company Statistix is Rs 70,000, you may think, "Wow! That's great." But if the standard deviation for starting salaries at Company Statistix is Rs 20,000, that's a lot of variation in terms of how much money you can make, so the average starting salary of Rs 70,000 isn't as informative in the end, is it?

- If the standard deviation was only Rs 5,000, you would have a much better idea of what to expect for a starting salary at that company



## Remarks on Expectation, Variance, Standard Deviation...

### Example

If you are told that the average starting salary for someone working at Company Statistix is Rs 70,000, you may think, "Wow! That's great." But if the standard deviation for starting salaries at Company Statistix is Rs 20,000, that's a lot of variation in terms of how much money you can make, so the average starting salary of Rs 70,000 isn't as informative in the end, is it?

- If the standard deviation was only Rs 5,000, you would have a much better idea of what to expect for a starting salary at that company
- Without calculating standard deviation, you can't get a handle on whether the data are close to the average

## Remarks on Expectation, Variance, Standard Deviation...

### Example

If you are told that the average starting salary for someone working at Company Statistix is Rs 70,000, you may think, "Wow! That's great." But if the standard deviation for starting salaries at Company Statistix is Rs 20,000, that's a lot of variation in terms of how much money you can make, so the average starting salary of Rs 70,000 isn't as informative in the end, is it?

- If the standard deviation was only Rs 5,000, you would have a much better idea of what to expect for a starting salary at that company
- Without calculating standard deviation, you can't get a handle on whether the data are close to the average

### Same Standard Deviation Does Not Imply Same Dataset

For example, the data sets 199, 200, 201 and 0, 200, 400 both have the same average (200) yet they have very different standard deviations.

## Remarks on Expectation, Variance, Standard Deviation...

### Example

If you are told that the average starting salary for someone working at Company Statistix is Rs 70,000, you may think, "Wow! That's great." But if the standard deviation for starting salaries at Company Statistix is Rs 20,000, that's a lot of variation in terms of how much money you can make, so the average starting salary of Rs 70,000 isn't as informative in the end, is it?

- If the standard deviation was only Rs 5,000, you would have a much better idea of what to expect for a starting salary at that company
- Without calculating standard deviation, you can't get a handle on whether the data are close to the average

### Same Standard Deviation Does Not Imply Same Dataset

For example, the data sets 199, 200, 201 and 0, 200, 400 both have the same average (200) yet they have very different standard deviations. The first data set has a very small standard deviation ( $s=1$ ) compared to the second data set ( $s=200$ ).

## Another expression for the variance

### Theorem (Another Expression for Variance)

If  $X$  is a discrete random variable with mean  $\mu$ , then

$$\text{Var}(X) = E[X^2] - \mu^2$$

## Another expression for the variance

### Theorem (Another Expression for Variance)

If  $X$  is a discrete random variable with mean  $\mu$ , then

$$\text{Var}(X) = E[X^2] - \mu^2$$

### Proof

$$\begin{aligned}\text{Var}(X) &= \sum_x (x - \mu)^2 p_X(x) = \sum_x (x^2 - 2\mu x + \mu^2) p_X(x) \\ &= \sum_x x^2 p_X(x) - 2\mu \sum_x x p_X(x) + \mu^2 \sum_x p_X(x) \\ &= E[X^2] - 2\mu E[X] + \mu^2 = E[X^2] - \mu^2\end{aligned}$$

## Properties of Variance...

## Properties of Variance...

### Theorem

Let  $X$  be a discrete random variable and  $\alpha$  a constant. Then

$$\text{Var}(\alpha X) = \alpha^2 \text{Var}(X) \quad \text{and} \quad \text{Var}(X + \alpha) = \text{Var}(X)$$

## Computing Variance: Binomial

### Variance of Binomial Distribution

Let  $X \sim \text{Binomial}(n, p)$ . Then the variance  $\text{Var}(X) = np(1 - p)$ .



## Variance of Binomial Distribution...

## Variance of Binomial Distribution...

## Outline

- ① Random Variables
- ② Special Distributions
- ③ Examples of Distributions
- ④ Expectations of Some Distributions
- ⑤ Variance and Standard Deviation: Understand Variability in Data
- ⑥ Higher Order Moments and Moment Generating Function

## Higher Order Moments...

### Define $n$ th moment

The  $n$ th moment about the mean or  $n$ th central moment of a real valued random variable  $X$  is defined as follows

$$\mu_n = E[(X - E[X])^n],$$

where  $E$  is the expectation operator.

## Higher Order Moments...

### Define $n$ th moment

The  $n$ th moment about the mean or  $n$ th central moment of a real valued random variable  $X$  is defined as follows

$$\mu_n = E[(X - E[X])^n],$$

where  $E$  is the expectation operator.

- The zeroth central moment  $\mu_0$  is 1

## Higher Order Moments...

### Define $n$ th moment

The  $n$ th moment about the mean or  $n$ th central moment of a real valued random variable  $X$  is defined as follows

$$\mu_n = E[(X - E[X])^n],$$

where  $E$  is the expectation operator.

- The zeroth central moment  $\mu_0$  is 1
- The first central moment  $\mu_1$  is 0 (not the same as expected value!)

## Higher Order Moments...

### Define $n$ th moment

The  $n$ th moment about the mean or  $n$ th central moment of a real valued random variable  $X$  is defined as follows

$$\mu_n = E[(X - E[X])^n],$$

where  $E$  is the expectation operator.

- The zeroth central moment  $\mu_0$  is 1
- The first central moment  $\mu_1$  is 0 (not the same as expected value!)
  - First central moment about origin is indeed expectation!

## Higher Order Moments...

### Define $n$ th moment

The  $n$ th moment about the mean or  $n$ th central moment of a real valued random variable  $X$  is defined as follows

$$\mu_n = E[(X - E[X])^n],$$

where  $E$  is the expectation operator.

- The zeroth central moment  $\mu_0$  is 1
- The first central moment  $\mu_1$  is 0 (not the same as expected value!)
  - First central moment about origin is indeed expectation!
- The second central moment  $\mu_2$  is called variance, denoted by  $\sigma^2$



## Higher Order Moments...

### Define $n$ th moment

The  $n$ th moment about the mean or  $n$ th central moment of a real valued random variable  $X$  is defined as follows

$$\mu_n = E[(X - E[X])^n],$$

where  $E$  is the expectation operator.

- The zeroth central moment  $\mu_0$  is 1
- The first central moment  $\mu_1$  is 0 (not the same as expected value!)
  - First central moment about origin is indeed expectation!
- The second central moment  $\mu_2$  is called variance, denoted by  $\sigma^2$

### Generating Moments...

Is there a quick way to generate moments?

## Moment Generating Function...

## Moment Generating Function...

### Moment Generating Function

The moment generating function  $M_X(t)$  is the expectation value

$$M_X(t) = E[e^{tX}] = \sum_x e^{tx} p_X(x)$$

### Lemma

- $M_X(0) = 1$
- $E[X] = M'_X(0)$ , where ' is the derivative w.r.t.  $t$

## Moment Generating Function for Binomial Distribution...

## Moment Generating Function for Binomial Distribution...

- Let  $X$  be a discrete random variable whose PMF is given by **Binomial** distribution with parameters  $n$  and  $p$

## Moment Generating Function for Binomial Distribution...

- Let  $X$  be a discrete random variable whose PMF is given by **Binomial** distribution with parameters  $n$  and  $p$
- The moment generating function is given by

## Moment Generating Function for Binomial Distribution...

- Let  $X$  be a discrete random variable whose PMF is given by **Binomial** distribution with parameters  $n$  and  $p$
- The moment generating function is given by

$$\begin{aligned}M_X(t) &= \sum_{x=0}^n \binom{n}{x} p^x (1-p)^{n-x} e^{tx} \\&= \sum_{x=0}^n \binom{n}{x} (e^t p)^x (1-p)^{n-x} \\&= (e^t p + 1 - p).\end{aligned}$$

## Moment Generating Function for Binomial Distribution...

- Let  $X$  be a discrete random variable whose PMF is given by **Binomial** distribution with parameters  $n$  and  $p$
- The moment generating function is given by

$$\begin{aligned}M_X(t) &= \sum_{x=0}^n \binom{n}{x} p^x (1-p)^{n-x} e^{tx} \\&= \sum_{x=0}^n \binom{n}{x} (e^t p)^x (1-p)^{n-x} \\&= (e^t p + 1 - p).\end{aligned}$$

Differentiating w.r.t.  $t$ , we have



## Moment Generating Function for Binomial Distribution...

- Let  $X$  be a discrete random variable whose PMF is given by **Binomial** distribution with parameters  $n$  and  $p$
- The moment generating function is given by

$$\begin{aligned}M_X(t) &= \sum_{x=0}^n \binom{n}{x} p^x (1-p)^{n-x} e^{tx} \\&= \sum_{x=0}^n \binom{n}{x} (e^t p)^x (1-p)^{n-x} \\&= (e^t p + 1 - p).\end{aligned}$$

Differentiating w.r.t.  $t$ , we have

$$M'_X(t) = n(e^t p + 1 - p)^{n-1} p e^t.$$

## Moment Generating Function for Binomial Distribution...

- Let  $X$  be a discrete random variable whose PMF is given by **Binomial** distribution with parameters  $n$  and  $p$
- The moment generating function is given by

$$\begin{aligned}M_X(t) &= \sum_{x=0}^n \binom{n}{x} p^x (1-p)^{n-x} e^{tx} \\&= \sum_{x=0}^n \binom{n}{x} (e^t p)^x (1-p)^{n-x} \\&= (e^t p + 1 - p).\end{aligned}$$

Differentiating w.r.t.  $t$ , we have

$$M'_X(t) = n(e^t p + 1 - p)^{n-1} p e^t.$$

- Setting  $t = 0$ ,  $M'_X(0) = np = E[X]$

## Moment Generating Function for Poisson Distribution...

## Moment Generating Function for Poisson Distribution...

- Let  $X$  be the random variable whose PMF is a Poisson distribution with parameter  $\lambda$ .

## Moment Generating Function for Poisson Distribution...

- Let  $X$  be the random variable whose PMF is a Poisson distribution with parameter  $\lambda$ .
- The moment generating function is

## Moment Generating Function for Poisson Distribution...

- Let  $X$  be the random variable whose PMF is a Poisson distribution with parameter  $\lambda$ .
- The moment generating function is

$$\begin{aligned}M_X(t) &= \sum_{x=0}^{\infty} e^{-\lambda} \frac{\lambda^x}{x!} e^{tx} \\&= \sum_{x=0}^{\infty} e^{-\lambda} \frac{(\lambda e^t)^x}{x!} \\&= e^{\lambda(e^t - 1)}\end{aligned}$$

## Moment Generating Function for Poisson Distribution...

- Let  $X$  be the random variable whose PMF is a **Poisson** distribution with parameter  $\lambda$ .
- The moment generating function is

$$\begin{aligned}M_X(t) &= \sum_{x=0}^{\infty} e^{-\lambda} \frac{\lambda^x}{x!} e^{tx} \\&= \sum_{x=0}^{\infty} e^{-\lambda} \frac{(\lambda e^t)^x}{x!} \\&= e^{\lambda(e^t - 1)}\end{aligned}$$

- Differentiating w.r.t. to  $t$ ,

## Moment Generating Function for Poisson Distribution...

- Let  $X$  be the random variable whose PMF is a Poisson distribution with parameter  $\lambda$ .
- The moment generating function is

$$\begin{aligned}M_X(t) &= \sum_{x=0}^{\infty} e^{-\lambda} \frac{\lambda^x}{x!} e^{tx} \\&= \sum_{x=0}^{\infty} e^{-\lambda} \frac{(\lambda e^t)^x}{x!} \\&= e^{\lambda(e^t-1)}\end{aligned}$$

- Differentiating w.r.t. to  $t$ ,

$$M'_X(t) = e^{\lambda(e^t-1)} \lambda e^t$$



## Moment Generating Function for Poisson Distribution...

- Let  $X$  be the random variable whose PMF is a Poisson distribution with parameter  $\lambda$ .
- The moment generating function is

$$\begin{aligned}M_X(t) &= \sum_{x=0}^{\infty} e^{-\lambda} \frac{\lambda^x}{x!} e^{tx} \\&= \sum_{x=0}^{\infty} e^{-\lambda} \frac{(\lambda e^t)^x}{x!} \\&= e^{\lambda(e^t-1)}\end{aligned}$$

- Differentiating w.r.t. to  $t$ ,

$$M'_X(t) = e^{\lambda(e^t-1)} \lambda e^t$$

- Setting  $t = 0$ ,  $M'_X(0) = \lambda$

## Variance Using Moment Generating Function...

## Variance Using Moment Generating Function...

### Variance Using Moment Generating Function

$$\text{Var}(X) = M_X''(0) - M_X'(0)^2$$

## Computing Variance Using Moment Generating Function...

## Computing Variance Using Moment Generating Function...

### Computing Variance using moment generating function

Let  $X$  be a discrete random variable whose PMF is a binomial distribution with parameters  $n$  and  $p$ . It has mean  $\mu = np$  and the moment generating function is

$$M_X(t) = (e^t p + 1 - p)^n$$

## Computing Variance Using Moment Generating Function...

### Computing Variance using moment generating function

Let  $X$  be a discrete random variable whose PMF is a binomial distribution with parameters  $n$  and  $p$ . It has mean  $\mu = np$  and the moment generating function is

$$M_X(t) = (e^t p + 1 - p)^n$$

Find the Variance using  $M_X(t)$ .

## Computing Variance Using Moment Generating Function...

## Computing Variance Using Moment Generating Function...

### Computing Variance using moment generating function

Let  $X$  be a discrete random variable whose PMF is a Poisson distribution with mean  $\lambda$ . The moment generating function is

$$M_X(t) = e^{\lambda(e^t - 1)}$$



## Computing Variance Using Moment Generating Function...

### Computing Variance using moment generating function

Let  $X$  be a discrete random variable whose PMF is a Poisson distribution with mean  $\lambda$ . The moment generating function is

$$M_X(t) = e^{\lambda(e^t - 1)}$$

Find the Variance using  $M_X(t)$ .