

# **Probability and Statistics**

UG2, Core course, IIIT,H

Pawan Kumar

IIIT, Hyderabad

November 19, 2021

- ① Joint Distributions: Two Random Variables
- ② Joint Continuous Random Variables
  - Marginal Continuous PDFs
  - Joint Cumulative Distribution
  - Conditional PDF, Conditional CDF, and Conditional Expectation
  - Conditional Expectation and Variance
  - Conditional PDF, Conditional Probability, and Conditional CDF

- Conditional Expectation and Variance Functions of Two Continuous Random Variables
- Computing CDF of Function of Two RVs
- Method of Transformation for Function of Two Variables
- Covariance and Correlation
- Solved Problems

## Outline

- ① Joint Distributions: Two Random Variables
- ② Joint Continuous Random Variables

**Fact...**

## Fact...

### Fact

Let  $X, Y$  be two RVs and  $g, h$  be two functions of  $X$  and  $Y$  respectively. Show that

$$E[g(X)h(Y) | X] = g(X)E[h(Y) | X]$$

### Solution

$$E[g(x)h(y) \Big| x=x] = g(x)E[h(y) \Big| x=x]$$

becomes a const  
instantiated with  $x$

**Iterated Expectations...**

## Iterated Expectations...

L.O.I.E

Law of Iterated Expectations

Let  $X, Y$  be two RVs, then we have

$$E[X] = E[E[X | Y]]$$

Proof

Recall  $g(Y) = E[X | Y]$  [Saw in last class]

Applying Law of Total Probability:

$$\underline{E[X]} = \sum_{Y_j \in \mathcal{Y}} E[X | Y=Y_j] P(Y=Y_j)$$

L.O.T.P

$$= \sum g(Y_j) P(Y=Y_j) = E[g(Y)] = \underline{\underline{E[E[X | Y]]}}$$

## Solved Example on Application of Iterated Expectation...

## Solved Example on Application of Iterated Expectation...

### Solved Example 2

Number of customers  $N$  visiting a fast food restaurant follows Poisson distribution  $N \sim \text{Poisson}(\lambda)$ .

## Solved Example on Application of Iterated Expectation...

### Solved Example 2

Number of customers  $N$  visiting a fast food restaurant follows Poisson distribution  $N \sim \text{Poisson}(\lambda)$ . Each customer arriving in this restaurant purchases a drink with probability  $p$ ,

## Solved Example on Application of Iterated Expectation...

### Solved Example 2

Number of customers  $N$  visiting a fast food restaurant follows Poisson distribution  $N \sim \text{Poisson}(\lambda)$ . Each customer arriving in this restaurant purchases a drink with probability  $p$ , which is independent from other customers.

## Solved Example on Application of Iterated Expectation...

L.O.I.E

### Solved Example 2

Number of customers  $N$  visiting a fast food restaurant follows Poisson distribution  $N \sim \text{Poisson}(\lambda)$ . Each customer arriving in this restaurant purchases a drink with probability  $p$ , which is independent from other customers. What is the average number of customers who purchase drinks?

Solution

Recall:  $X = \# \text{customers who purchase drinks with prob } p$   
 $N = \# \text{customers visiting the restaurant}$   $\sim \text{Poisson}(\lambda)$

Recall:  $X | N \sim \text{Binomial}(n, p)$

$$E[X] \stackrel{\text{L.O.I.E}}{=} E[E[X|N]] = E[Np] = pE[N] = \underline{\underline{p\lambda}}$$

## Expectation for Independent RV...

## Expectation for Independent RV...

### Expectation for Independent RVs

Let  $X, Y$  be two independent RVs. Then we have the following

- 1  $E[X | Y] = E[X]$  ←
- 2  $E[g(X) | Y] = E[g(X)]$  ←
- 3  $E[XY] = E[X]E[Y]$  ←
- 4  $E[g(X)h(Y)] = E[g(X)]E[h(Y)]$  ←

$$\textcircled{1} \quad E[X | Y=y] = \sum_{x \in R_X} P_{X|Y}(x|y) \rightarrow \star$$

$$\text{Since } X, Y \text{ ind.} \Rightarrow P_{X|Y}(x|y) = P_X(x)$$

$$\text{From } \textcircled{1}: \quad \sum_x P_X(x) \stackrel{\text{def}}{=} E[X] //$$

Answer to previous problem...



Answer to previous problem...



## Conditional Variance...

## Conditional Variance...

### Definition of Conditional Variance

Let  $X, Y$  be two RVs.

- $X|Y \sim \text{RN}$
- $P_{X|Y}, f_{X|Y}$
- $E[X|Y]$

## Conditional Variance...

### Definition of Conditional Variance

Let  $X, Y$  be two RVs. By  $\text{Var}(X | Y = y)$  the conditional variance of  $X$  given  $Y = y$ .

## Conditional Variance...

### Definition of Conditional Variance

Let  $X, Y$  be two RVs. By  $\text{Var}(X | Y = y)$  the conditional variance of  $X$  given  $Y = y$ . Let  $\mu_{X|Y}(y) = E[X | Y = y]$ . Then

## Conditional Variance...

$$\text{Var}(X) = \underline{E[X^2] - E[X]^2}$$

### Definition of Conditional Variance

Let  $X, Y$  be two RVs. By  $\text{Var}(X | Y = y)$  the conditional variance of  $X$  given  $Y = y$ . Let  $\mu_{X|Y}(y) = E[X | Y = y]$ . Then

$$\underbrace{\text{Var}(X | Y = y)}_{\text{Definition}} = E[X^2 | Y = y] - \mu_{X|Y}(y)^2$$

(<sup>17</sup> probabilitycourse.com)

Proof

Assume

## Solved Example ...

## Solved Example ...

$$X|Y = 1$$

### Solved Example

Let  $X, Y$  be RV with joint PMF given as follows

	$Y = 0$	$Y = 1$
$X = 0$	$\frac{1}{5}$	$\frac{2}{5}$
$X = 1$	$\frac{2}{5}$	0

Let  $Z = E[X | Y]$  and  $V = \text{Var}(X | Y)$ .

- 1 Find the PMF of  $V$
- 2 Find  $E[V]$
- 3 Verify that  $\text{Var}(X) = E[V] + \text{Var}(Z)$

Recall:  $X, Y \sim \text{Bernoulli}(\frac{2}{5})$

and

$X|Y=0 \sim \text{Bernoulli}(\frac{2}{3})$

$\text{Var}(X) = p(1-p)$   
if  $X$  is Bernoulli( $p$ )

Note that  $V$  is a fn of  $Y$ .

$$V = \text{Var}(X|Y) = \begin{cases} \text{Var}(X|Y=0) & Y=0 \\ \text{Var}(X|Y=1) & Y=1 \end{cases}$$

$$\begin{aligned} & \stackrel{?}{=} \left\{ \begin{array}{l} \text{Var}(X|Y=0), \text{ with prob } \frac{3}{5} \\ \text{Var}(X|Y=1), \text{ with prob } \frac{2}{5} \end{array} \right\} \quad \text{Since } Y \sim \text{Bernoulli}(\frac{2}{5}) \\ & \text{Var}(X|Y=0) = \frac{2}{3} \cdot \frac{1}{3} = \frac{2}{9} \\ & \boxed{\text{Var}(X|Y=1) = 0} \quad [\text{check}] \end{aligned}$$

Answer to previous problem...

$$V = \text{Var}(X|Y) = \begin{cases} 2/5 & \text{with prob } 3/5 \\ 0 & \text{with prob } 2/5 \end{cases}$$

(c)  $\text{Var}(X) = \frac{2}{5} \cdot \frac{3}{5}$

a)  $P_V(v)$  =

$$\begin{cases} 3/5 & \text{if } v = 2/5 \\ 2/5 & \text{if } v = 0 \\ 0 & \text{otherwise} \end{cases}$$

$E(V) = 1/45 = \frac{2}{15}$

$\text{Var}(Z) = 8/75$  [check]

(b)  $E(V) = \frac{2}{5} \cdot \frac{3}{5} + 0 \cdot \frac{4}{5}$

$$= \frac{6}{25}$$

(Check this in Probability course via TA)

$$\frac{6}{25} = \frac{\frac{2}{15} + \frac{8}{75}}{?}$$
$$\frac{6}{25} = \frac{10 + 8}{75}$$

Answer to previous problem...



Answer to previous problem...



## Law of Total Variance...

## Law of Total Variance...

### Law of Total Variance

Let  $X, Y$  be two RVs.

## Law of Total Variance...

### Law of Total Variance

Let  $X, Y$  be two RVs. The law of total variance says that

## Law of Total Variance...

### Law of Total Variance

Let  $X, Y$  be two RVs. The law of total variance says that

$$\text{Var}(X) = E[\text{Var}(X | Y)] + \text{Var}(E[X | Y])$$

Proof:

See this video

## Solved Problem 1

### Solved Problem 1

Let  $X, Y$  be two independent RVs with the same CDFs  $F_X$  and  $F_Y$ . Let

$$Z = \max(X, Y)$$

$$W = \min(X, Y)$$

Find the CDFs of  $Z$  and  $W$ .

Answer to previous problem...



**Answer to previous problem...**

## Outline

### ① Joint Distributions: Two Random Variables

### ② Joint Continuous Random Variables

Marginal Continuous PDFs

Joint Cumulative Distribution

Conditional PDF, Conditional CDF, and Conditional Expectation

Conditional Expectation and Variance

Conditional PDF, Conditional Probability, and Conditional CDF

Conditional Expectation and Variance

Functions of Two Continuous Random Variables

Computing CDF of Function of Two RVs

Method of Transformation for Function of Two Variables

Covariance and Correlation

Solved Problems

## Joint Probability Density Function...

## Joint Probability Density Function...

### Joint Continuous Density Functions

Two RVs  $X, Y$  are jointly continuous if there exists a nonnegative function  $f_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ , such that, for any set  $A \in \mathbb{R}^2$ , we have

$$P(\underbrace{(X, Y) \in A}_{}) = \int \int_A f_{XY}(x, y) dx dy$$

## Joint Probability Density Function...

### Joint Continuous Density Functions

Two RVs  $X, Y$  are **jointly continuous** if there exists a nonnegative function  $f_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ , such that, for any set  $A \in \mathbb{R}^2$ , we have

$$P((X, Y) \in A) = \int \int_A f_{XY}(x, y) \, dxdy$$

The function  $\underline{f_{XY}(x, y)}$  is called the **joint probability density function**, PDF, of  $X$  and  $Y$ .

## Joint Probability Density Function...

### Joint Continuous Density Functions

Two RVs  $X, Y$  are **jointly continuous** if there exists a nonnegative function  $f_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ , such that, for any set  $A \in \mathbb{R}^2$ , we have

$$P((X, Y) \in A) = \int \int_A f_{XY}(x, y) dx dy$$

The function  $f_{XY}(x, y)$  is called the **joint probability density function**, PDF, of  $X$  and  $Y$ .

- 1 Here domain of  $f_{XY}(x, y)$  is  $\mathbb{R}^2$ , but we may redefine

## Joint Probability Density Function...

### Joint Continuous Density Functions

Two RVs  $X, Y$  are **jointly continuous** if there exists a nonnegative function  $f_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ , such that, for any set  $A \in \mathbb{R}^2$ , we have

$$P((X, Y) \in A) = \int \int_A f_{XY}(x, y) dx dy$$

The function  $f_{XY}(x, y)$  is called the **joint probability density function**, PDF, of  $X$  and  $Y$ .

- 1 Here domain of  $f_{XY}(x, y)$  is  $\mathbb{R}^2$ , but we may redefine

$$\underline{R}_{XY} = \{(x, y) \mid \underline{f_{X,Y}}(x, y) > 0\}$$

## Joint Probability Density Function...

### Joint Continuous Density Functions

Two RVs  $X, Y$  are **jointly continuous** if there exists a nonnegative function  $f_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ , such that, for any set  $A \in \mathbb{R}^2$ , we have

$$P((X, Y) \in A) = \int \int_A f_{XY}(x, y) dx dy$$

The function  $f_{XY}(x, y)$  is called the **joint probability density function**, PDF, of  $X$  and  $Y$ .

- 1 Here domain of  $f_{XY}(x, y)$  is  $\mathbb{R}^2$ , but we may redefine

$$R_{XY} = \{(x, y) \mid f_{X,Y}(x, y) > 0\}$$

- 2 If we indeed choose  $A = \mathbb{R}^2$ , then we must have

## Joint Probability Density Function...

### Joint Continuous Density Functions

Two RVs  $X, Y$  are **jointly continuous** if there exists a nonnegative function  $f_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ , such that, for any set  $A \in \mathbb{R}^2$ , we have

$$P((X, Y) \in A) = \int \int_A f_{XY}(x, y) dx dy$$

The function  $f_{XY}(x, y)$  is called the **joint probability density function**, PDF, of  $X$  and  $Y$ .

- 1 Here domain of  $f_{XY}(x, y)$  is  $\mathbb{R}^2$ , but we may redefine

$$R_{XY} = \{(x, y) \mid f_{X,Y}(x, y) > 0\}$$

- 2 If we indeed choose  $A = \mathbb{R}^2$ , then we must have

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x, y) dx dy = 1$$

## Solved Example...

## Solved Example...

### Solved Example

Let  $X, Y$  be two jointly continuous RVs with joint PDF

## Solved Example...

### Solved Example

Let  $X, Y$  be two jointly continuous RVs with joint PDF

$$f_{XY}(x, y) = \begin{cases} x + cy^2 & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

## Solved Example...

$$\iint f_{XY}(x,y) = 1$$

### Solved Example

Let  $X, Y$  be two jointly continuous RVs with joint PDF

$$f_{XY}(x,y) = \begin{cases} x + cy^2 & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- Find the constant  $c$

We have


$$\iint_0^1 (x + cy^2) dx dy = 1$$
$$\text{L.H.S} = \left[ \frac{x^2}{2} + cy^2 x \right]_0^1 = \left[ \frac{1}{2} + cy^2 \right]_0^1 = \frac{1}{2} + \frac{c}{2} = 1$$
$$\Rightarrow c = \frac{3}{2}$$

## Solved Example...

### Solved Example

Let  $X, Y$  be two jointly continuous RVs with joint PDF

$$f_{XY}(x, y) = \begin{cases} x + cy^2 & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$



1 Find the constant  $c$

2 Find  $P(0 \leq X \leq \frac{1}{2}, 0 \leq Y \leq \frac{1}{2})$

$$P(0 \leq X \leq y_2, 0 \leq Y \leq y_2) =$$

$$\int_0^{y_2} \left[ \int_0^{x_2} \left( x + \frac{3}{2}y^2 \right) dx \right] dy$$

Answer to previous problem...



Answer to previous problem...



Marginal Continuous PDF...

## Marginal Continuous PDF...

$$P_X(x) = \sum_y P_{XY}(x,y)$$

### Marginal Continuous PDF

Let  $X, Y$  be two random variable.

## Marginal Continuous PDF...

### Marginal Continuous PDF

Let  $X, Y$  be two random variable. Then the marginal PDFs of  $X$  and  $Y$  can be computed from the joint PDF  $f_{XY}$  as follows

## Marginal Continuous PDF...

### Marginal Continuous PDF

Let  $X, Y$  be two random variable. Then the marginal PDFs of  $X$  and  $Y$  can be computed from the joint PDF  $f_{XY}$  as follows

$$\underline{f_X(x)} = \int_{-\infty}^{\infty} f_{XY}(x, y) \underline{dy}, \quad \text{for all } x,$$

## Marginal Continuous PDF...

### Marginal Continuous PDF

Let  $X, Y$  be two random variable. Then the marginal PDFs of  $X$  and  $Y$  can be computed from the joint PDF  $f_{XY}$  as follows

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy, \quad \text{for all } x,$$

$$\underline{f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx}, \quad \text{for all } y.$$

## Solved Example...

## Solved Example...

### Solved Example

Let  $X, Y$  be two jointly continuous RVs with joint PDF

## Solved Example...

### Solved Example

Let  $X, Y$  be two jointly continuous RVs with joint PDF

$$f_{XY}(x, y) = \begin{cases} x + cy^2 & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

## Solved Example...

### Solved Example

Let  $X, Y$  be two jointly continuous RVs with joint PDF

$$f_{XY}(x, y) = \begin{cases} x + cy^2 & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- Find the PDFs  $f_X(x)$  and  $f_Y(y)$

$$\underline{f_X(x)} = \int_{-\infty}^{\infty} \left( x + \frac{3}{2}y^2 \right) dy = \int_0^1 \left( x + \frac{3}{2}y^2 \right) dy$$

Try

Answer to previous problem...



## Solved Example...

## Solved Example...

### Solved Example

Let  $X, Y$  be two jointly continuous RVs with joint PDFs

## Solved Example...

### Solved Example

Let  $X, Y$  be two jointly continuous RVs with joint PDFs

$$f_{XY}(x, y) = \begin{cases} cx^2y & 0 \leq y \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

## Solved Example...

### Solved Example

Let  $X, Y$  be two jointly continuous RVs with joint PDFs

$$f_{XY}(x, y) = \begin{cases} cx^2y & 0 \leq y \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- 1 Find  $R_{XY}$  and plot it

## Solved Example...

### Solved Example

Let  $X, Y$  be two jointly continuous RVs with joint PDFs

$$f_{XY}(x, y) = \begin{cases} cx^2y & 0 \leq y \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- 1 Find  $R_{XY}$  and plot it
- 2 Find the constant  $c$

## Solved Example...

### Solved Example

Let  $X, Y$  be two jointly continuous RVs with joint PDFs

$$f_{XY}(x, y) = \begin{cases} cx^2y & 0 \leq y \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- 1 Find  $R_{XY}$  and plot it
- 2 Find the constant  $c$
- 3 Find the marginal PDFs  $f_X(x)$  and  $f_Y(y)$

## Solved Example...

### Solved Example

Let  $X, Y$  be two jointly continuous RVs with joint PDFs

$$f_{XY}(x, y) = \begin{cases} cx^2y & 0 \leq y \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- 1 Find  $R_{XY}$  and plot it
- 2 Find the constant  $c$
- 3 Find the marginal PDFs  $f_X(x)$  and  $f_Y(y)$
- 4 Find  $P\left(Y \leq \frac{X}{2}\right)$

## Solved Example...

### Solved Example

Let  $X, Y$  be two jointly continuous RVs with joint PDFs

$$f_{XY}(x, y) = \begin{cases} cx^2y & 0 \leq y \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- 1 Find  $R_{XY}$  and plot it →
- 2 Find the constant  $c$
- 3 Find the marginal PDFs  $f_X(x)$  and  $f_Y(y)$
- 4 Find  $P\left(Y \leq \frac{X}{2}\right)$
- 5 Find  $P\left(Y \leq \frac{X}{2} \mid Y \leq \frac{X}{2}\right)$

**Solution to previous problem...**

## Solution to previous problem...

Looking at the joint PDF, we have

## Solution to previous problem...

Looking at the joint PDF, we have

$$R_{XY} = \{(x, y) \in \mathbb{R}^2 \mid 0 \leq y \leq x \leq 1\}$$

## Solution to previous problem...

Looking at the joint PDF, we have

$$R_{XY} = \{(x, y) \in \mathbb{R}^2 \mid 0 \leq y \leq x \leq 1\}$$

The plot is the following

## Solution to previous problem...

Looking at the joint PDF, we have

$$R_{XY} = \{(x, y) \in \mathbb{R}^2 \mid 0 \leq y \leq x \leq 1\}$$

The plot is the following





Figure: Figure showing  $R_{XY}$  and integration region  $P(Y \leq \frac{X}{2})$

Answer to previous problem...

$$\textcircled{5} \quad \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x,y) dx dy = 1$$

$$\Rightarrow \frac{C}{2} \left[ \frac{x^5}{5} \right]_0^1 = 1$$

$$\Rightarrow \int_0^1 \int_0^x cx^2 y dy dx = 1$$

$$\Rightarrow \frac{C}{10} = 1$$

$$\Rightarrow \int_0^1 \left[ cx^2 y \right]_0^x dx = 1$$

$$\Rightarrow C = 10$$

$$\Rightarrow \frac{C}{2} \left[ x^4 \right]_0^1 = 1$$

Answer to previous problem...

c) To find the marginal

$$R_x \cap R_y = [0, 1]$$

For  $0 \leq x \leq 1$

$$\begin{aligned} f_x(x) &= \int_{-\infty}^{\infty} f_{xy}(y|x) dy \\ &= \int_0^x 10x^2 y dy = \left[ 10x^2 \frac{y^2}{2} \right]_0^x \\ &= 5x^4 // \end{aligned}$$

$$\begin{aligned} d) P(Y \leq x/2) &= \int_0^{\infty} \int_0^{x/2} f_{xy}(y|x) dy dx \\ &= \int_0^{\infty} \int_0^{x/2} 10xy dy dx \\ &= \text{easy } \underline{\text{to}} \end{aligned}$$

## Joint Cumulative Distribution...

## Joint Cumulative Distribution...

### Joint cumulative distribution

Let  $X, Y$  be two continuous RVs with joint CDF  $F_{XY}(x, y)$  as follows

## Joint Cumulative Distribution...

### Joint cumulative distribution

Let  $X, Y$  be two continuous RVs with joint CDF  $F_{XY}(x, y)$  as follows

$$\underline{F_{XY}(x, y) = P(X \leq x, Y \leq y)}.$$

## Joint Cumulative Distribution...

### Joint cumulative distribution

Let  $X, Y$  be two continuous RVs with joint CDF  $F_{XY}(x, y)$  as follows

$$F_{XY}(x, y) = P(X \leq x, Y \leq y).$$

The joint CDF satisfies the following properties:

## Joint Cumulative Distribution...

### Joint cumulative distribution

Let  $X, Y$  be two continuous RVs with joint CDF  $F_{XY}(x, y)$  as follows

$$F_{XY}(x, y) = P(X \leq x, Y \leq y).$$

The joint CDF satisfies the following properties:

- 1  $F_X(x) = F_{XY}(x, \infty)$  for any  $x$  (marginal CDF of  $X$ )

## Joint Cumulative Distribution...

### Joint cumulative distribution

Let  $X, Y$  be two continuous RVs with joint CDF  $F_{XY}(x, y)$  as follows

$$F_{XY}(x, y) = P(X \leq x, Y \leq y).$$

The joint CDF satisfies the following properties:

- 1  $F_X(x) = F_{XY}(x, \infty)$  for any  $x$  (marginal CDF of  $X$ )
- 2  $F_Y(y) = F_{XY}(\infty, y)$  for any  $y$  (marginal CDF of  $Y$ )

## Joint Cumulative Distribution...

### Joint cumulative distribution

Let  $X, Y$  be two continuous RVs with joint CDF  $F_{XY}(x, y)$  as follows

$$F_{XY}(x, y) = P(X \leq x, Y \leq y).$$

The joint CDF satisfies the following properties:

- 1  $F_X(x) = F_{XY}(x, \infty)$  for any  $x$  (marginal CDF of  $X$ )
- 2  $F_Y(y) = F_{XY}(\infty, y)$  for any  $y$  (marginal CDF of  $Y$ )
- 3  $F_{XY}(\infty, \infty) = 1$

## Joint Cumulative Distribution...

### Joint cumulative distribution

Let  $X, Y$  be two continuous RVs with joint CDF  $F_{XY}(x, y)$  as follows

$$F_{XY}(x, y) = P(X \leq x, Y \leq y).$$

The joint CDF satisfies the following properties:

- 1  $F_X(x) = F_{XY}(x, \infty)$  for any  $x$  (marginal CDF of  $X$ )
- 2  $F_Y(y) = F_{XY}(\infty, y)$  for any  $y$  (marginal CDF of  $Y$ )
- 3  $F_{XY}(\infty, \infty) = 1$
- 4  $\underline{F_{XY}(-\infty, y)} = F_{XY}(x, \underline{-\infty}) = 0$

## Joint Cumulative Distribution...


### Joint cumulative distribution

Let  $X, Y$  be two continuous RVs with joint CDF  $F_{XY}(x, y)$  as follows

$$F_{XY}(x, y) = P(X \leq x, Y \leq y).$$

The joint CDF satisfies the following properties:

- 1  $F_X(x) = F_{XY}(x, \infty)$  for any  $x$  (marginal CDF of  $X$ )
- 2  $F_Y(y) = F_{XY}(\infty, y)$  for any  $y$  (marginal CDF of  $Y$ )
- 3  $F_{XY}(\infty, \infty) = 1$
- 4  $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0$
- 5  $P(x_1 < X \leq x_2, y_1 < Y \leq y_2) = F_{XY}(x_2, y_2) - F_{XY}(x_2, y_1) - F_{XY}(x_1, y_2) + F_{XY}(x_1, y_1)$



## Joint Cumulative Distribution...

### Joint cumulative distribution

Let  $X, Y$  be two continuous RVs with joint CDF  $F_{XY}(x, y)$  as follows

$$F_{XY}(x, y) = P(X \leq x, Y \leq y).$$

The joint CDF satisfies the following properties:

- 1  $F_X(x) = F_{XY}(x, \infty)$  for any  $x$  (marginal CDF of  $X$ )
- 2  $F_Y(y) = F_{XY}(\infty, y)$  for any  $y$  (marginal CDF of  $Y$ )
- 3  $F_{XY}(\infty, \infty) = 1$
- 4  $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0$
- 5  $P(x_1 < X \leq x_2, y_1 < Y \leq y_2) = F_{XY}(x_2, y_2) - F_{XY}(x_2, y_1) - F_{XY}(x_1, y_2) + F_{XY}(x_1, y_1)$
- 6 If  $X, Y$  are independent, then  $F_{XY}$  =  $F_X(x)F_Y(y)$

## Solved Example

### Solved Example Independent

Let  $X, Y$  be two random variables with Uniform(0,1) distribution. Find  $F_{XY}(x, y)$ .

Since  $X, Y$  are uniform(0,1) we have

$$F_X(x) = \begin{cases} 0 & \text{for } x < 0 \\ x & \text{for } 0 \leq x \leq 1 \\ 1 & \text{for } x > 1 \end{cases}, F_Y(y) = \begin{cases} 0 & \text{for } y < 0 \\ y & \text{for } 0 \leq y \leq 1 \\ 1 & \text{for } y > 1 \end{cases}$$

Since  $X, Y$  are independent  $\Rightarrow F_{XY}(x, y) = \underline{F_X(x)} \underline{F_Y(y)}$ .

$$\underline{F_{XY}(x, y)} = \begin{cases} 0 & \text{for } x < 0, y < 0 \\ xy & \text{for } 0 \leq x \leq 1, 0 \leq y \leq 1 \\ y & \text{for } x > 1, 0 \leq y \leq 1 \\ x & \text{for } y > 1, 0 \leq x \leq 1 \\ 1 & \text{for } y > 1, x > 1 \end{cases}$$

Answer to previous problem...



**Figure for Solved Example...**

## Figure for Solved Example...




Figure: Plot of joint CDF

## Relationship Between CDF and PDF...

## Relationship Between CDF and PDF...

### Relationship between CDF and PDF

Recall that for single RV we have

## Relationship Between CDF and PDF...

### Relationship between CDF and PDF

Recall that for single RV we have

$$F_x(x) = \int_{-\infty}^x f_x(u) du$$

$$\underline{f_x(x)} = \frac{dF_x(x)}{dx}$$

## Relationship Between CDF and PDF...

### Relationship between CDF and PDF

Recall that for single RV we have

$$F_X(x) = \int_{-\infty}^x f_X(u) du$$
$$f_X(x) = \frac{dF_X(x)}{dx}$$

Similarly, for two RVs we have

## Relationship Between CDF and PDF...

### Relationship between CDF and PDF

Recall that for single RV we have

$$F_X(x) = \int_{-\infty}^x f_X(u) du$$
$$f_X(x) = \frac{dF_X(x)}{dx}$$

Similarly, for two RVs we have

$$\underline{F_{XY}(x, y)} = \int_{-\infty}^y \int_{-\infty}^x \underline{f_{XY}(u, v)} du dv$$
$$\underline{\underline{f_{XY}}} = \frac{\partial^2}{\partial x \partial y} F_{XY}(x, y)$$

## Example of Joint CDF...

## Example of Joint CDF...

### Solved Example

Let  $X, Y$  be two jointly continuous RVs with joint PDF

## Example of Joint CDF...

### Solved Example

Let  $X, Y$  be two jointly continuous RVs with joint PDF

$$f_{XY}(x, y) = \begin{cases} x + cy^2 & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

## Example of Joint CDF...

### Solved Example

Let  $X, Y$  be two jointly continuous RVs with joint PDF

$$f_{XY}(x, y) = \begin{cases} x + cy^2 & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

$\frac{3}{2}$

- Find the joint CDF of  $X$  and  $Y$

$$\begin{aligned} F_{XY}(x, y) &= \int_{-\infty}^x \int_{-\infty}^y f_{XY}(u, v) du dv = \int_{-\infty}^x \int_{-\infty}^y \left(u + \frac{3}{2}v^2\right) du dv \\ &= \int_{-\infty}^y \left[ \frac{u^2}{2} + \frac{3}{2}v^2 u \right]_0^x dv = \int_{-\infty}^y \left[ \frac{x^2}{2} + \frac{3}{2}v^2 x \right] dv = \left[ \frac{x}{2}v + \frac{3}{2}\frac{v^3}{3}x \right]_0^y \\ &= \frac{x^2 y}{2} + \frac{y^3 x}{2} \end{aligned}$$

**Answer to previous problem...**

Answer to previous problem...



## Definition of Conditional PDF and Conditional CDF

Let  $X$  be a continuous RV and  $A$  be an event that  $a < X < b$  (where possibly  $b = \infty$  or  $a = -\infty$ ), then

$$\rightarrow f_{X|A}(x) = \begin{cases} 1 & x > b \\ \frac{F_X(x) - F_X(a)}{F_X(b) - F_X(a)} & a \leq x < b \\ 0 & x < a \end{cases}$$

$$\rightarrow f_{X|A}(x) = \begin{cases} \frac{f_X(x)}{P(A)} & a \leq x < b \\ 0 & \text{otherwise} \end{cases}$$

Answer to previous problem...



Answer to previous problem...



## Conditional Expectation and Variance...

## Conditional Expectation and Variance...

### Definition of Conditional Expectation and Variance

For a random variable  $X$  and event  $A$ , we have

## Conditional Expectation and Variance...

### Definition of Conditional Expectation and Variance

For a random variable  $X$  and event  $A$ , we have

$$\left. \begin{aligned} E[X | A] &= \int_{-\infty}^{\infty} xf_{X|A}(x) dx \\ E[g(X) | A] &= \int_{-\infty}^{\infty} g(x)f_{X|A}(x) dx \\ \text{Var}(X | A) &= E[X^2 | A] - (E[X | A])^2 \end{aligned} \right\}$$

## Solved Example: Conditional PDF and CDF...

## Solved Example: Conditional PDF and CDF...

### Solved Example

Let  $X \sim \text{Exponential}(1)$ .

## Solved Example: Conditional PDF and CDF...

### Solved Example

Let  $X \sim \text{Exponential}(1)$ . Answer the following.

## Solved Example: Conditional PDF and CDF...

### Solved Example

Let  $X \sim \text{Exponential}(1)$ . Answer the following.

- 1 Find the conditional PDF and CDF of  $X$  given  $X > 1$

## Solved Example: Conditional PDF and CDF...

### Solved Example

Let  $X \sim \text{Exponential}(1)$ . Answer the following.

- 1 Find the conditional PDF and CDF of  $X$  given  $X > 1$
- 2 Find  $E[X | X > 1]$

## Solved Example: Conditional PDF and CDF...

### Solved Example

Let  $X \sim \text{Exponential}(1)$ . Answer the following.

- 1 Find the conditional PDF and CDF of  $X$  given  $X > 1$
- 2 Find  $E[X | X > 1]$
- 3 Find  $\text{Var}(X | X > 1)$

④ Let  $A$  be the event that  $X > 1$

$$\underline{\underline{P(A)}} = \int_1^{\infty} e^{-x} dx = 1/e$$

$$\underline{\underline{f_{X|X>1}(x)}} = \frac{P(X=x, X>1)}{P(X>1)} = \frac{e^{-x}}{1/e} = e^{-x}$$

$$F_{X|X>1}(x) = \int_1^x f_{X|A}(u) du$$

$$= \int_1^x e^{-u+1} du = \left[ \frac{e^{-u+1}}{-1} \right]_1^x$$

$$= \frac{e^{-x+1}}{-1} + \frac{e^{-1+1}}{-1} = \underline{\underline{1 - e^{-x+1}}}$$

Answer to previous problem...

$$E[X|X>1] = \int_{-\infty}^{\infty} x f_{X|X>1}(x) dx$$

$$= \int_1^{\infty} x e^{-x+1} dx = \cancel{try}$$