

HEP NTUA Top Angular Report

2/4/2021

George Bakas




Summary Z' Analysis

- Switch to $m_{JJ} > 1000$ GeV cut:
 - No sensitivity for higher Z' masses (> 2.5 TeV)
 - Calculate significance and Brazilian plots for different m_{JJ} cuts
 - m_{JJ} Cuts: [1000, 1200, 1400, 1600, 1800, 2000] GeV
 - $Significance = \frac{Signal}{\sqrt{Signal+Bkg}}$ where signal is the Z' distribution and $Bkg := ttbar + QCD + Subdominant$
- Calculate asymptotic limit to find optimal m_{JJ} cut
 - Compare this to the significance for each m_{JJ} cut
- I was using $ttbar$ as the extracted signal from data: Instead I use the $ttbar$ MC distribution (scaled to the signal strength)
- For QCD I use the QCD MC distribution which is scaled to data (using k-factor)
- Both the Significance and Brazilian plots use these files as input → this is the reason that the Brazilian plots are different than what I showed on the HEP NTUA weekly on 24th of March
- Sliding m_{JJ} Cut




Significance (left) and Asymptotic (right) Graphs (2017)

Significance_M1200_W12




$M_{Z'} = 1200, w = 1\%$

Asymptotic M1200 W12




$M_{Z'} = 1200, w = 1\%$

Significance_M1400_W14



$M_{Z'} = 1400, w = 1\%$

Asymptotic M1400 W14




$M_{Z'} = 1400, w = 1\%$




Significance (left) and Asymptotic (right) Graphs (2017)

Significance_M1600_W16




$M_{Z'} = 1600, w = 1\%$

Asymptotic M1600 W16




$M_{Z'} = 1600, w = 1\%$

Significance_M1800_W18



$M_{Z'} = 1800, w = 1\%$

Asymptotic M1800 W18




$M_{Z'} = 1800, w = 1\%$




Significance (left) and Asymptotic (right) Graphs (2017)


Significance_M2000_W20




Asymptotic M2000 W20



Significance_M2500_W25




Asymptotic M2500 W25




Significance (left) and Asymptotic (right) Graphs (2017)


Significance_M3000_W30




Asymptotic M3000 W30



Significance_M3500_W35



Asymptotic M3500 W35



$M_{Z'} = 3000, w = 1\%$

$M_{Z'} = 3000, w = 1\%$


$M_{Z'} = 3500, w = 1\%$

$M_{Z'} = 3500, w = 1\%$




Significance (left) and Asymptotic (right) Graphs (2017)

Significance_M4000_W40




$M_{Z'} = 4000$, $w = 1\%$

Asymptotic M4000 W40




$M_{Z'} = 4000$, $w = 1\%$

Significance_M4500_W45



$M_{Z'} = 4500$, $w = 1\%$

Asymptotic M4500 W45




$M_{Z'} = 4500$, $w = 1\%$




Brazilian Plots (2017)

$m_{JJ} > 1000 \text{ GeV}$




$m_{JJ} > 1200 \text{ GeV}$




Asymptotic Limits - Brazilian Plots (2017)

$m_{JJ} > 1400$ GeV




$m_{JJ} > 1600$ GeV




Brazilian Plots (2017)

$m_{JJ} > 1800$ GeV



$m_{JJ} > 2000$ GeV




Brazilian Plots (2017 and 2018) with sliding mJJ Cut


Mass Cut Mapping

{"mZ_1200_12":1000, "mZ_1400_14":1200, "mZ_1600_16":1400, "mZ_1800_18":1600, "mZ_2000_20":1600, "mZ_2500_25":2000, "mZ_3000_30":2000, "mZ_3500_35":2000, "mZ_4000_40":2000, "mZ_4500_45":2000}

2017



2018



BACKUP




Angular Distributions (Brazilian Plot using data!!!!)


Asymptotic limits for $M_{Z'}$: 1.2, 1.4, 1.6, 1.8, 2, 2.5, 3, 3.5, 4, 4.5 TeV

Width 1%

2017



2018




Angular Distributions (Brazilian Plot using extracted signal!!) vs B2G-16-015


Asymptotic limits for $M_{Z'}$: 1.2, 1.4, 1.6, 1.8, 2, 2.5, 3, 3.5, 4, 4.5 TeV

Width 1%

2017



B2G-16-015




Angular Distributions (Brazilian Plot using extracted signal!!) vs B2G-16-015


Asymptotic limits for $M_{Z'}$: 1.2, 1.4, 1.6, 1.8, 2, 2.5, 3, 3.5, 4, 4.5 TeV

Width 1%

2018



B2G-16-015




Signal Selection

Variables	Selected Cut
pT (both leading jets)	> 400 GeV
Njets	> 1
N leptons	= 0
eta (both leading jets)	< 2.4
mJJ	> 1000 GeV
jetMassSoftDrop (only for fit)	(50,300) GeV
Top Tagger	> 0.2, 0, 0.1
B tagging (2 btagged jets)	> Medium WP
Signal Trigger	

Control Region Selection

Variables	Selected Cut
pT (both leading jets)	> 400 GeV
Njets	> 1
N leptons	= 0
eta (both leading jets)	< 2.4
mJJ	> 1000 GeV
jetMassSoftDrop (only for fit)	(50,300) GeV
Top Tagger	> 0.2, 0, 0.1
B tagging (0 btagged jets)	< Medium WP
Control Trigger	



Top Angular Distributions

- We employ the dijet angular variable χ from the rapidities of the two leading jets
- Why χ ?
 - The distributions associated with the final states produced via QCD interactions are relatively flat in comparison with the distributions of the BSM models or new particles, which typically peak at low values of χ
- We can measure the variable χ in two ways

1. By measuring the difference of the rapidities of the two leading jets such as the corresponding rapidity in the ZMF is:

$$y^* = \frac{1}{2}(y_1 - y_2)$$

χ is defined as $\chi = e^{|y^*|} = e^{|y_1 - y_2|}$ (1) and can be measured by creating the TLorentzVector, boost it to the ZMF and find the rapidity difference of the two leading jets

2. By measuring the scattering angle θ^* (angle between top quark and z-axis in the Zero Momentum Frame)

We define as $y^* = \frac{1}{2} \ln\left(\frac{1+|\cos\theta^*|}{1-|\cos\theta^*|}\right)$ and from (1) we can find that:

$$\chi = \frac{1 + |\cos\theta^*|}{1 - |\cos\theta^*|}$$

