

HEP NTUA Weekly Report

17/3/2021

George Bakas



Summary

ttX analysis:

- Removed geometrical matching from files
 - Chain of analysis completed
 - Integrated angular variables
- I managed to apply combination of results
 - Comparison with Giannis
 - Particle level also
- Looking into results without matching

Z' analysis

- Switch to UL files for Z' (missing: 2016_postVFP (all widths) and 2016_preVFP, 2017, 2018 10% and 30% width files)
 - Email to B2G MC generation group
- Production of 1200, 1400, 1600 and 1800 GeV Masses
 - mJJ cut is now 1 TeV as in the ttX analysis
 - Calculation of XSEC using [GenXSecAnalyzer](#)
 - Results (work in progress, something is not looking good)

1% width: (dataset name: ZPrimeToTT_M*_TuneCP2_13TeV-madgraph-pythia8)

2016 [B2G-2020Nov18-00005](#) 17 samples Submitted Ongoing

2016APV [B2G-2020Nov18-00006](#) 17 samples Submitted DONE

2017 [B2G-2020Nov18-00007](#) 10 samples Submitted DONE

2018 [B2G-2020Nov18-00008](#) 10 samples Submitted DONE

10% and 30% width: (dataset name: ZPrimeToTT_M*_TuneCP2_13TeV-madgraph-pythia8)

2016 [B2G-2021Feb17-00002](#) 34 samples Submitted Ongoing

2016 APV [B2G-2021Feb17-00004](#) 34 samples Submitted Ongoing

2016 [B2G-2021Feb17-00006](#) 34 samples Submitted Ongoing

2016 [B2G-2021Feb17-00008](#) 34 samples Submitted Ongoing



BACKUP



Summary

Angular Distributions, Z' analysis:

- New Signal Region:
 - $SR_C = SR + m_{JJ} > 1.5\text{TeV}$
- Stack histograms for SR_C
- Asymptotic Limits (Brazilian plots) for 2016, 2017, 2018
 - Total Cross section x BR
 - Total Cross section = $\sum_{i=1}^N S_i$, where S_i is the signal yield in the reconstructed level
- X distributions show a different slope than the B2G-16-015
 - Recreated Brazilian plot using m_{JJ} variable (only for 2016 and Zprime 1% width)
 - Tried to increase mass cut from 1.5 TeV to 2 TeV to improve sensitivity → not enough events coming from signal extraction
 - If I use $t\bar{t}$ MC (χ dists) as input, the shape is the same as with the 1.5 TeV cut
 - Maybe sliding mass cuts? For each Z' use a different m_{JJ} cut



Signal Selection

Variables	Selected Cut
pT (both leading jets)	> 400 GeV
Njets	> 1
N leptons	= 0
eta (both leading jets)	< 2.4
mJJ	> 1000 GeV
jetMassSoftDrop (only for fit)	(50,300) GeV
Top Tagger	> 0.2, 0, 0.1
B tagging (2 btagged jets)	> Medium WP
Signal Trigger	

Control Region Selection

Variables	Selected Cut
pT (both leading jets)	> 400 GeV
Njets	> 1
N leptons	= 0
eta (both leading jets)	< 2.4
mJJ	> 1000 GeV
jetMassSoftDrop (only for fit)	(50,300) GeV
Top Tagger	> 0.2, 0, 0.1
B tagging (0 btagged jets)	< Medium WP
Control Trigger	



Signal Extraction

$$S_{1.5TeV}(x_{reco}) = D_{1.5TeV}(x_{reco}) - QCD_{1.5TeV}(x_{reco}) - Sub_{1.5TeV}(x_{reco}) \rightarrow$$

$$\text{Where } QCD_{1.5TeV}(x_{reco}) = D_{1.5TeV,shape}^{0-btag}(x_{reco}) \times N_{SR(1.5TeV)} \times C_{closure}^{shape SF}$$

$$\text{and } N_{SR(1.5TeV)} = R_{yield}^{1TeV \rightarrow 1.5TeV} \times N_{SR(1TeV)}^{QCD} = R_{yield}^{1TeV \rightarrow 1.5TeV} \times R_{yield}^{SRA \rightarrow SR} \times N_{SRA}^{QCD}$$

- The variable of interest here: $x_{reco} \rightarrow \chi$
- 1.5 TeV refers to the mJJ cut
- We deploy a fit in the Signal Region (2btag) to extract the N_{QCD}^{fit} in SRA (mJJ > 1TeV)

$$D(m^t)^{(i)} = N_{tt}^{(i)} T^{(i)}(m^t, k_{MassScale}, k_{MassResolution}) + N_{bkg}^{(i)} B(m^t)(1 + k_1 x) + N_{sub}^{(i)} O^{(i)}(m^t)$$



Top Angular Distributions

- We employ the dijet angular variable χ from the rapidities of the two leading jets
- Why χ ?
 - The distributions associated with the final states produced via QCD interactions are relatively flat in comparison with the distributions of the BSM models or new particles, which typically peak at low values of χ
- We can measure the variable χ in two ways

1. By measuring the difference of the rapidities of the two leading jets such as the corresponding rapidity in the ZMF is:

$$y^* = \frac{1}{2}(y_1 - y_2)$$

χ is defined as $\chi = e^{|y^*|} = e^{|y_1 - y_2|}$ (1) and can be measured by creating the TLorentzVector, boost it to the ZMF and find the rapidity difference of the two leading jets

2. By measuring the scattering angle θ^* (angle between top quark and z-axis in the Zero Momentum Frame)

We define as $y^* = \frac{1}{2} \ln\left(\frac{1+|\cos\theta^*|}{1-|\cos\theta^*|}\right)$ and from (1) we can find that:

$$\chi = \frac{1 + |\cos\theta^*|}{1 - |\cos\theta^*|}$$

