


Weekly Report NTUA 13/12/2019

George Bakas



Status Report

- TOP-18-013 approval
- Signal Extraction
 - Various methods for extracting number of QCD events in Signal reduced mass region
 - Fixed eb or free eb
- Unfolding
 - Response Matrices where Nbins Reco \sim 2Nbins Parton/Particle
 - Unfolding to parton and particle level
 - For a number of regularization parameters tau do the unfolding and find the average global correlation
 - $\rho_j = \sqrt{1 - [(V_x)_{jj} - (V_x^{-1})_{jj}]^{-1}}$ where V_x is the covariance matrix of x (gen level)
 - We select the **tau that minimizes ρ**
 - The tau spectrum is $\sim [10E-10, 10]$



Signal Extraction

$$S(x_{reco}) = D(x_{reco}) - N_{QCD, reduced}^{(2)} C_{QCD}^{shape}(x_{reco}) Q(x_{reco}) - B(x_{reco})$$

Diagram illustrating the components of the signal extraction equation:

- Fiducial Yield**: $S(x_{reco})$
- ABCD Method**: $D(x_{reco})$ (Measured dist from data)
- QCD shape correction factor**: $N_{QCD, reduced}^{(2)} C_{QCD}^{shape}(x_{reco})$ (QCD shape taken from Data (CR))
- Subdominant bkg shape and contribution (MC)**: $B(x_{reco})$

- Where x_{reco} is the respected variable of interest (ttbar mass, pt, rapidity, leading and subleading jetPt and |jetY|)
- We deploy a simultaneous fit in 3 regions (0,1,2 btag) because we do not have a pure Control Region.
 - Our data CR is ttbar contaminated

$$D(m^t)^{(i)} = N_{tt}^{(i)} T^{(i)}(m^t, k_{MassScale}, k_{MassResolution}) + N_{bkg}^{(i)} B(m^t)(1 + k_1 x) + N_{sub}^{(i)} O^{(i)}(m^t)$$

- We assume that $N_{tt}^{(0)} = (1 - e_b)^2 N_{tt}$, $N_{tt}^{(2)} = e_b^2 N_{tt}$ and $N_{tt}^{(1)} = 2(1 - e_b)e_b N_{tt}$ where e_b is the b tagging efficiency and N_{tt} is the total ttbar yield.



Transfer Factor Calculation

We will use an ABCD method to extract the N_{QCD} in the SR


From shape we have:

$$\frac{N_{\text{QCD, reduced}}^{(2)}}{N_{\text{QCD, reduced}}^{(0)}} = \frac{N_{\text{QCD, extended}}^{(2)}}{N_{\text{QCD, extended}}^{(0)}} \Rightarrow$$

$$N_{\text{QCD, reduced}}^{(2)} = \frac{N_{\text{QCD, extended}}^{(2)}}{N_{\text{QCD, extended}}^{(0)}} N_{\text{QCD, reduced}}^{(0)}$$

↓ Taken from fit using free e_b

↓ $D_{\text{reduced}}^{(0)} - TT_{\text{MC, reduced}}^{(0)}$



NQCD in Reduced (SR):

2016: 846.365


2017: 677.041

2018: 1201.35




Fiducial Measurements (jetPt0) ABCD method and free eb


2016



2017



2018




- Output is not consistent with what we expected
 - We expect $r \sim 0.85$ for 2016, ~ 0.65 for 2017 and ~ 0.75 for 2018
 - After signal extraction $r \sim 0.96$ for 2016, ~ 0.63 for 2017 and ~ 0.89 for 2018
 - Nqcd in SR reduced probably the problem (not enough QCD extracted?)




Fiducial Measurements (m_{JJ}) ABCD method and free $e b$


2016



2017




2018



Fiducial Measurements for all years combined

Fid. Measurements All years combined



Fid. Measurements All years combined

