# HEP NTUA Weekly Report

9/2/2022

George Bakas





## Summary

- ttX analysis:
  - New UL b-tag WPs for 2016preVFP and 2016 postVFP released
    - Switch and implementation
  - $p_T>450$  GeV for leading jet and  $p_T>400$  GeV for second leading jet
  - We are writing the AN:
    - 1. Basic outline along with text
    - 2. Input all images that are needed for the analysis
    - 3. Appendices that include
      - Response matrices, efficiencies, acceptance, purity and stability per year
      - Fiducial Measurements per year
      - Systematic uncertainties breakdown per year
      - Closure tests
    - 4. Issues are handled on gitlab
  - Z' analysis:
    - Production for files that were missing
    - Integration of M1400 W14 for 2016\_preVFP in analysis chain


Contents


Content	•	t	า	ľ	е	t	n	ı	o	C	(	
---------	---	---	---	---	---	---	---	---	---	---	---	--

2	1	Introduction
3	2	Samples
4		2.1 Data
5		2.2 Simulation
6	3	Trigger
7	4	Reconstrunction and Selection
8		4.1 Object Reconstruction
9		4.2 Selection
10		4.3 Multivariate Discriminant
11		4.4 Parton level
12		4.5 Particle level
13	5	Signal Extraction
14		5.1 Inclusive cross section
15		5.2 Differential cross sections
16	6	Data vs Monte Carlo
17	7	Fiducial Measurement
18	8	Systematic Uncertainties
19	9	Combination of different years
20	10	Unfolded Measurement
21		10.1 Parton Level
22		10.2 Particle Level
23	A	appendix/Fiducial





### **Efficiency Parton Level**







### **Efficiency Particle Level**







#### Acceptance Parton Level






#### Acceptance Particle Level






### Mass Fit Results 2016 (pre and postVFP)

#### A RooPlot of "mTop"




#### A RooPlot of "mTop"




### Mass Fit Results 2017, 2018


#### A RooPlot of "mTop"



#### A RooPlot of "mTop"





### Systematics Breakdown Combined Initial Fiducial Result






### Systematics Breakdown Combined Initial Fiducial Result












### Systematics Breakdown Combined Initial Fiducial Result












### Systematics Breakdown Combined Unfolded Result






#### Systematics Breakdown Combined Unfolded Result










## Systematics Breakdown Combined Unfolded Result





























































































































## **BACKUP**



## Summary


- ttX analysis Pipeline Creation
  - 1. We want to be able to handle all Nominal files and their variations in an automated way
  - This requires deciding consistent naming conventions and a efficient planning
  - 3. Handling of:
    - 1. Nominal
    - 2. Parton Shower Weights
    - PDF Variations
    - 4. JES
    - 5. Scale Variations
    - 6. bTagVariations
    - 7. Top quark mass variations
  - 4. Per year For all these we need to
    - 1. Create template files that have 2btag and 0btag in Extended and Reduced jetMassSoftDrop phase space
    - 2. 9 variables (mJJ, pTJJ, yJJ, jetPt[0,1], jetY[0,1], chi, |cosTheta\*|[0,1]
    - 3. Template fit files (bkg qcd, bkg subdominant) and signal templates for all variations
    - 4. Fit on extended signal region for all variations

- 5. Response matrices, Acceptance, Efficiency
- 6. Signal Extraction
- Combine all Fiducial Level results (4 years) into 1 Extracted Signal for all variations
- 6. Unfold the combined result into Parton & Particle levels
- 7. Show systematic variations compared to the Nominal file
- 8. The same procedure must be done using different nominal files
  - 1. Fill in 2btag histograms in our signal region in the parton
  - 2. For each variation and each year
  - 3. Combine all years together
  - 4. Calculate systematics for samples other than the nominal





## Brazilian Plots (2016\_preVFP, 2017 and 2018) with sliding mJJ Cut

2016\_preVFP 2017



2018








#### Combined Datacard for 2016 preVFP, 2017 and 2018

#### Mass Cut Mapping


```
{"mZ_1200_12":1000, "mZ_1400_14":1200, "mZ_1600_16":1400, "mZ_1800_18":1600, "mZ_2000_20":1600, "mZ_2500_25":2000, "mZ_3000_30":2000, "mZ_3500_35":2000, "mZ_4000_40":2000, "mZ_4500_45":2000}
```





## Brazilian Plots (2016\_preVFP, 2017 and 2018) with sliding mJJ Cut wrt 2018

2016\_preVFP 2017



#### Combined Datacard for 2016 preVFP, 2017 and 2018 wrt 2018

#### Mass Cut Mapping

```
{"mZ_1200_12":1000, "mZ_1400_14":1200, "mZ_1600_16":1400, "mZ_1800_18":1600, "mZ_2000_20":1600, "mZ_2500_25":2000, "mZ_3000_30":2000, "mZ_3500_35":2000, "mZ_4000_40":2000, "mZ_4500_45":2000}
```

