

# Measurement of differential production cross section for boosted top quarks in the all hadronic channel

NTUA  
23/9/2020

G. Bakas, K. Kousouris, I. Papakrivopoulos, G. Tsipolitis



---



## Motivation

Top, anti-top production in the fully hadronic final state.

Trying to identify two big jets that contain the products of the top/anti-top decay.



# Overview

- Variables of interest:
  - ttbar mass, pt, rapidity
  - Leading and Subleading jetPt and |jetY|
  - Nominal ttbar MC samples
  - Mtt samples (700-1000, 1000-Inf) (only for 2016)
- Baseline Parton cuts:
  - Jet Matching
  - partonPt[0],[1] > 400
  - |partonEta[0],[1]| < 2.4
  - mTTbarParton > 1000 GeV
- Baseline Reconstructed level cuts:
  - nJets > 1, nLeptons = 0, Dijet mass (mJJ) > 1000
  - Leading and Subleading jet  $p_T > 400$
  - Leading and Subleading absolute jet eta  $|\eta| < 2.4$
- Btagging selection:
  - bTagging (medium WP **deepCSV**) (2016: 0.6321, 2017: 0.4941, 2018: 0.4184)
- Top Tagger WP:
  - New top Tagger:** (2016: 0.2, 2017:0.0, 2018: 0.1)

Region	Requirements
Signal Region (SR)	Baseline + topTagger + $m_{SD}^{jet1,2} \in (120,220)GeV + 2btags$
Control Region (CR)	Baseline + topTagger + $m_{SD}^{jet1,2} \in (120,220)GeV + 0btags$
Extended SR (SR <sub>A</sub> ) (QCD fit region)	Baseline + topTagger + $m_{SD}^{jet1,2} \in (50,300)GeV + 2btags$
Extended CR (CR <sub>A</sub> ) (QCD fit region)	Baseline + topTagger + $m_{SD}^{jet1,2} \in (50,300)GeV + 0btags$

Goal is to Unfold to the Parton And Particle Levels

- Closure Tests with Nominal MC's
- Unfolding and extrapolation with Data

Discovered that our Control Region is contaminated from ttbar and Subdominant bkg:

- Extract this contribution from the Data CR distribution → pure QCD

Differences with TOP-18-013:

- mJJ and mTTbarParton cut at 1000 GeV instead of 800 GeV
- New top tagger, tagging jets and not events. The goal is to have higher efficiency in the far end of the spectrum




**B tagging SF's are applied**




# Efficiency and Acceptance Plots

## Parton Efficiency '16,'17,'18 NominalMC




## Parton Acceptance '16,'17,'18 NominalMC




# Efficiency and Acceptance Plots

Parton Efficiency '16,'17,'18 NominalMC




Parton Acceptance '16,'17,'18 NominalMC



# Response Matrices


(2016)

Response Reco-Parton jetY0 2016 NominalMC




(2017)

Response Reco-Parton jetY0 2017 NominalMC



(2018)

Response Reco-Parton jetY0 2018 NominalMC




Assuming that response matrices are compatible to each other

1. Unfold each year and combine results?
2. Combine Fiducial Measurements and then unfold




# Contamination Plots Medium WP (CR) 2016, 2017, 2018


(2016)



(2017)





(2018)



# Contamination Plots Medium WP (CR) 2016, 2017, 2018

## Comment

Shapes and in general contamination is NOT affected by btagging by Scale factors






# Mass Fit in Extended SR ( $SR_A$ )

$$QCD_0(m^t) = D_0(m^t) - T_0(m^t) - Sub_0(m^t)$$

- Both SR and Control Region use the Medium btag WP.
- Intuition is to remove the  $t\bar{t}$  and subdominant bkg contribution from the data Control Region


A RooPlot of "mTop"



Floating Parameter	FinalValue	+/-	Error
kMassResol	9.2251e-01	+/-	2.73e-02
kMassScale	9.9891e-01	+/-	2.01e-03
kQCD_2b	6.9753e-02	+/-	5.26e-02
nFitBkg_2b	2.4472e+02	+/-	1.47e+02
nFitQCD_2b	2.9890e+03	+/-	1.74e+02
nFitSig2b	5.2763e+03	+/-	1.67e+02

$t\bar{t}$  Signal strength:  $r = 0.686668 \pm 0.0263103$  (new)


A RooPlot of "mTop"



Floating Parameter	FinalValue	+/-	Error
kMassResol	1.0998e+00	+/-	4.02e-02
kMassScale	9.8340e-01	+/-	2.66e-03
kQCD_2b	1.6593e-02	+/-	7.44e-03
nFitBkg_2b	4.9791e+02	+/-	2.68e+02
nFitQCD_2b	2.1662e+03	+/-	3.11e+02
nFitSig2b	4.8059e+03	+/-	1.50e+02

$t\bar{t}$  Signal strength:  $r = 0.644361 \pm 0.023851$  (new)

A RooPlot of "mTop"




Floating Parameter	FinalValue	+/-	Error
kMassResol	1.0071e+00	+/-	2.84e-02
kMassScale	9.9006e-01	+/-	1.92e-03
kQCD_2b	1.2862e-02	+/-	2.71e-03
nFitBkg_2b	1.0737e+02	+/-	2.97e+02
nFitQCD_2b	5.0444e+03	+/-	2.83e+02
nFitSig2b	7.6593e+03	+/-	1.86e+02

$t\bar{t}$  Signal strength:  $r = 0.683382 \pm 0.0200866$



# Signal Strength Results



# Signal Extraction

$$S(x_{reco}) = D(x_{reco}) - C_{bkg}^{yield} N_{QCD}^{fit} C_{QCD}^{shape}(x_{reco}) Q(x_{reco}) - B(x_{reco})$$

Diagram illustrating the components of the signal extraction equation:


- Fiducial Yield**: Points to  $S(x_{reco})$
- Measured dist from data**: Points to  $D(x_{reco})$
- Fitted number of QCD events in  $SR_A$** : Points to  $N_{QCD}^{fit}$
- QCD shape taken from Data (CR)**: Points to  $C_{QCD}^{shape}(x_{reco})$
- Transfer factor from  $SR_A$  to SR**: Points to  $C_{bkg}^{yield}$
- QCD shape correction factor**: Points to  $C_{QCD}^{shape}(x_{reco})$
- Subdominant bkg shape and contribution (MC)**: Points to  $B(x_{reco})$

- Where  $x_{reco}$  is the respected variable of interest (ttbar mass, pt, rapidity, leading and subleading jetPt and |jetY|)
  - We deploy a fit in the Signal Region (2btag) to extract the  $N_{QCD}^{fit}$
- $$D(m^t)^{(i)} = N_{tt}^{(i)} T^{(i)}(m^t, k_{MassScale}, k_{MassResolution}) + N_{bkg}^{(i)} B(m^t)(1 + k_1 x) + N_{sub}^{(i)} O^{(i)}(m^t)$$
- Our data CR is contaminated from ttbar and subdominant bkg which has to be dealt with.




# Fiducial Differential Cross Section


Data vs MC 2016 for jetPt1




Data vs MC 2017 for jetPt1




Data vs MC 2018 for jetPt1




Data vs MC 2016 for jetPt1



Data vs MC 2017 for jetPt1




Data vs MC 2018 for jetPt1



# Parton & Particle levels

Parton

Observable	Requirement
$p_T^{t,\bar{t}}$	$> 400 \text{ GeV}$
$ \eta^{t,\bar{t}} $	$< 2.4$
$m_{t\bar{t}}$	$> 1000 \text{ GeV}$



Particle level Top Candidates

Observable	Requirement
$N_{jets}$	$> 1$
$p_T^{jet1,2}$	$> 400 \text{ GeV}$
$ \eta^{jet1,2} $	$< 2.4$
$m_{SD}^{jet1,2}$	$(120, 220) \text{ GeV}$
$m_{jj}$	$> 1000 \text{ GeV}$

$$\frac{d\sigma_i^{\text{unf}}}{dx} = \frac{1}{\mathcal{L} \cdot \Delta x_i} \cdot \frac{1}{f_{2,i}} \cdot \sum_j \left( R_{ij}^{-1} \cdot f_{1,j} \cdot S_j \right)$$

efficiency of the reco+true selection

migration matrix


reco efficiency of the reco+true selection

Unfolding: simple response matrix inversion w/o regularisation




# Parton Differential Cross Section


Particle Unfolded vs Theory jetPt1 2016



Parton Unfolded vs Theory jetPt1 2017




Parton Unfolded vs Theory jetPt1 2018




**WORK IN PROGRESS**


Parton Unfolded vs Theory jetPt1 2016



Parton Unfolded vs Theory jetPt1 2017




Parton Unfolded vs Theory jetPt1 2018




# Parton Differential Cross Section Comparison


Parton DataOverMC ratio (2016, TOP18013)




Normalised Parton DataOverMC ratio (2016, TOP18013)




Parton DataOverMC ratio (2017, TOP18013)




Normalised Parton DataOverMC ratio (2017, TOP18013)



Parton DataOverMC ratio (2018, TOP18013)




Normalised Parton DataOverMC ratio (2018, TOP18013)




# Parton Differential Cross Section Comparison

**WORK IN PROGRESS**

Parton Unfolded Ratio Comparison ('16,'17,'18)



Normalized Parton Unfolded Ratio Comparison ('16,'17,'18)





# Tag And Probe

- Top Tagger Scale Factors

- Validation method to ensure that no SF's are needed
- From data we subtract QCD and Subdominant bkg (MC) so that the data sample is pure

$$efficiency = \frac{\# (1 \text{ jet pass baseline} + \textit{Tight TopTagger Cut AND 1 jet pass SR})}{\# (1 \text{ jet pass baseline} + \textit{Tight TopTagger Cut AND 1 jet pass only baseline})}$$

- Randomization: Randomly select leading/subleading jet to use as tag or probe to avoid pT bias
- Divide the phase space into pT regions based on the topTagger categories: [400-600] GeV, [600-800] GeV, [800-Inf] GeV



# Tag And Probe Calculations

(2016)

eff data:  $0.781 \pm 0.038$

eff ttbar:  $0.772 \pm 0.014$

-----

Efficiency per Pt region

eff data pT[400-600]:  $0.761 \pm 0.042$

eff ttbar pT[400-600]:  $0.778 \pm 0.016$

-----

eff data pT[600-800]:  $0.851 \pm 0.100$

eff ttbar pT[600-800]:  $0.748 \pm 0.031$

-----

eff data pT[800-Inf]:  $0.886 \pm 0.160$

eff ttbar pT[800-Inf]:  $0.775 \pm 0.063$

(2017)

eff data:  $0.857 \pm 0.040$

eff ttbar:  $0.875 \pm 0.0072$

-----

Efficiency per Pt region

eff data pT[400-600]:  $0.872 \pm 0.047$

eff ttbar pT[400-600]:  $0.874 \pm 0.008$

-----

eff data pT[600-800]:  $0.795 \pm 0.088$

eff ttbar pT[600-800]:  $0.876 \pm 0.018$

-----

eff data pT[800-Inf]:  $0.797 \pm 0.186$

eff ttbar pT[800-Inf]:  $0.899 \pm 0.045$

(2018)

eff data:  $0.798 \pm 0.034$

eff ttbar:  $0.839 \pm 0.005$

-----

Efficiency per Pt region

eff data pT[400-600]:  $0.793 \pm 0.04$

eff ttbar pT[400-600]:  $0.836 \pm 0.006$

-----

eff data pT[600-800]:  $0.829 \pm 0.066$

eff ttbar pT[600-800]:  $0.851 \pm 0.013$

-----

eff data pT[800-Inf]:  $0.752 \pm 0.13$

eff ttbar pT[800-Inf]:  $0.865 \pm 0.032$



# Summary

- Applied the b-tagging SF's
  - Significant effect on the ttbar signal strength on **2017** and **2018**
- We have extracted the first results of the cross section in the Fiducial and the parton/particle levels
- Start investigating ttbar Systematic Uncertainties

- **2016** Nominal MC:

Tune CUETP8M2T4	TuneCP5
/TT_TuneCUETP8M2T4_13TeV-powheg-pythia8/RunIISummer16MiniAODv3-PUMoriond17_94X_mcRun2_asymptotic_v3-v1/MINIAODSIM	/TTToHadronic_TuneCP5_PSweights_13TeV-powheg-pythia8/RunIISummer16MiniAODv3-PUMoriond17_94X_mcRun2_asymptotic_v3-v1/MINIAODSIM (TTToSemiLeptonic, TTTo2L2Nu)

- Investigate on how to combine the measurements between the three years
  - Combine them in the fiducial level and extract the cross section?
  - Extract the cross sections individually and combine the measurements in the unfolded level?



# BACKUP



## Signal Selection

Variables	Selected Cut
pT (both leading jets)	> 400 GeV
Njets	> 1
N leptons	= 0
eta  (both leading jets)	< 2.4
mJJ	> 1000 GeV
jetMassSoftDrop (only for fit)	(50,300) GeV
Top Tagger	> 0.2, 0, 0.1
B tagging (2 btagged jets)	> Medium WP
Signal Trigger	

## Control Region Selection

Variables	Selected Cut
pT (both leading jets)	> 400 GeV
Njets	> 1
N leptons	= 0
eta  (both leading jets)	< 2.4
mJJ	> 1000 GeV
jetMassSoftDrop (only for fit)	(50,300) GeV
Top Tagger	> 0.2, 0, 0.1
B tagging (0 btagged jets)	< Medium WP
Control Trigger	



Year	Type of File	DAS
2016	TT Mtt 700-1000	<a href="#">/TT_Mtt-700to1000_TuneCUETP8M2T4_13TeV-powheg-pythia8/RunIISummer16MiniAODv3-PUMoriond17_94X_mcRun2_asymptotic_v3-v2/MINIAODSIM</a>
	TT Mtt 1000-Inf	<a href="#">/TT_Mtt-1000toInf_TuneCUETP8M2T4_13TeV-powheg-pythia8/RunIISummer16MiniAODv3-PUMoriond17_94X_mcRun2_asymptotic_v3-v2/MINIAODSIM</a>
	TT Nominal	<a href="#">/TT_TuneCUETP8M2T4_13TeV-powheg-pythia8/RunIISummer16MiniAODv3-PUMoriond17_94X_mcRun2_asymptotic_v3-v1/MINIAODSIM</a>
2017	TT Mtt 700-1000	--
	TT Mtt 1000-Inf	--
	TT Nominal Hadronic	<a href="#">/TTToHadronic_TuneCP5_13TeV-powheg-pythia8/RunIIFall17MiniAODv2-PU2017_12Apr2018_94X_mc2017_realistic_v14-v1/MINIAODSIM</a>
	TT Nominal Semilepton	<a href="#">/TTToSemiLeptonic_TuneCP5_13TeV-powheg-pythia8/RunIIFall17MiniAODv2-PU2017_12Apr2018_94X_mc2017_realistic_v14-v2/MINIAODSIM</a>
	TT Nominal Dilepton	<a href="#">/TTTo2L2Nu_TuneCP5_13TeV-powheg-pythia8/RunIIFall17MiniAODv2-PU2017_12Apr2018_94X_mc2017_realistic_v14-v2/MINIAODSIM</a>
2018	TT Mtt 700-1000	--
	TT Mtt 1000-Inf	--
	TT Nominal Hadronic	<a href="#">/TTToHadronic_TuneCP5_13TeV-powheg-pythia8/RunIIAutumn18MiniAOD-102X_upgrade2018_realistic_v15-v1/MINIAODSIM</a>
	TT Nominal Semilepton	<a href="#">/TTToSemiLeptonic_TuneCP5_13TeV-powheg-pythia8/RunIIAutumn18MiniAOD-102X_upgrade2018_realistic_v15-v1/MINIAODSIM</a>
	TT Nominal Dilepton	<a href="#">/TTTo2L2Nu_TuneCP5_13TeV-powheg-pythia8/RunIIAutumn18MiniAOD-102X_upgrade2018_realistic_v15-v1/MINIAODSIM</a>




# Overview: Discriminator, Efficiency and Acceptance

The discriminator is a BDT trained individually for 2016, 2017 and 2018


Category training: split the sample in categories based on Pt

- Bins:
  - [400, 600] GeV
  - [600, 800] GeV
  - [800, 1200] GeV
  - [1200, inf] GeV
- BDT, used variables:
  - Leading and Sub-leading subjet mass
  - N-Subjetiness variables (tau1, tau2, tau3)
  - fraction of the jetPt over the total pt sum of the event.
  - Energy correlation functions (ecfB1N2,ecfB1N3,ecfB2N2, ecfB2N3)
- BDT Output consistency for the 3 years
- Calculation of Efficiency and acceptance for each year
  - We choose the WP's for each year so that the leading jet  $p_T$  efficiency is similar for all years




2016







2017



2018



# Fit Params Results Comparison






# Response Matrices 2016


b tagging SF's

without b tagging SF's

Response Reco-Parton mJJ 2016 NominalMC



Response Reco-Parton mJJ 2016 NominalMC




# Response Matrices 2017


b tagging SF's

without b tagging SF's

Response Reco-Parton mJJ 2017 NominalMC



Response Reco-Parton mJJ 2017 NominalMC




# Response Matrices 2018


b tagging SF's

without b tagging SF's

Response Reco-Parton mJJ 2018 NominalMC




Response Reco-Parton mJJ 2018 NominalMC




# Fiducial Differential Cross Section Comparison


Fiducial DataOverMC ratio (2016, TOP18013)




Fiducial DataOverMC ratio (2017, TOP18013)




Fiducial DataOverMC ratio (2018, TOP18013)




Normalised Fiducial DataOverMC ratio (2016, TOP18013)



Normalised Fiducial DataOverMC ratio (2017, TOP18013)




Normalised Fiducial DataOverMC ratio (2018, TOP18013)




# Parton Differential Cross Section Comparison

Parton Unfolded Ratio Comparison ('16,'17,'18)




Normalized Parton Unfolded Ratio Comparison ('16,'17,'18)






# Parton Differential Cross Section Comparison

Parton Unfolded Ratio Comparison ('16,'17,'18)




Normalized Parton Unfolded Ratio Comparison ('16,'17,'18)








To scale the ttbar  $\rightarrow$  fit the leading jetMassSoftDrop in each of these regions (ttbar compatible  $\sim$  with SR)



## Tight TopTagger + Probe



# Tag And Probe Calculations 2016

## b tagging SF's

Efficiency--with btagging SF's

eff data:  $0.781 \pm 0.038$

eff ttbar:  $0.772 \pm 0.014$

-----

Efficiency per Pt region

eff data pT[400-600]:  $0.761 \pm 0.042$

eff ttbar pT[400-600]:  $0.778 \pm 0.016$

-----

eff data pT[600-800]:  $0.851 \pm 0.100$

eff ttbar pT[600-800]:  $0.748 \pm 0.031$

-----

eff data pT[800-Inf]:  $0.886 \pm 0.160$

eff ttbar pT[800-Inf]:  $0.775 \pm 0.063$

## without b tagging SF's

Efficiency--without btagging SF's

eff data:  $0.782 \pm 0.039$

eff ttbar:  $0.772 \pm 0.014$

-----

Efficiency per Pt region

eff data pT[400-600]:  $0.762 \pm 0.043$

eff ttbar pT[400-600]:  $0.778 \pm 0.016$

-----

eff data pT[600-800]:  $0.854 \pm 0.103$

eff ttbar pT[600-800]:  $0.748 \pm 0.031$

-----

eff data pT[800-Inf]:  $0.888 \pm 0.161$

eff ttbar pT[800-Inf]:  $0.775 \pm 0.064$





# Tag And Probe Calculations 2017

## b tagging SF's

Efficiency-- with btagging SF's

eff data:  $0.857 \pm 0.040$

**eff ttbar:  $0.875 \pm 0.0072$**

-----

Efficiency per Pt region

eff data pT[400-600]:  $0.872 \pm 0.047$

eff ttbar pT[400-600]:  $0.874 \pm 0.008$

-----

eff data pT[600-800]:  $0.795 \pm 0.088$

eff ttbar pT[600-800]:  $0.876 \pm 0.018$

-----

eff data pT[800-Inf]:  $0.797 \pm 0.186$

eff ttbar pT[800-Inf]:  $0.899 \pm 0.045$

## without b tagging SF's

Efficiency-- without btagging SF's

eff data:  $0.864 \pm 0.043$

**eff ttbar:  $0.875 \pm 0.007$**

-----

Efficiency per Pt region

eff data pT[400-600]:  $0.880 \pm 0.049$

eff ttbar pT[400-600]:  $0.874 \pm 0.008$

-----

eff data pT[600-800]:  $0.8 \pm 0.091$

eff ttbar pT[600-800]:  $0.876 \pm 0.018$

-----

eff data pT[800-Inf]:  $0.796 \pm 0.2$

eff ttbar pT[800-Inf]:  $0.898 \pm 0.045$



# Tag And Probe Calculations 2018

## b tagging SF's

Efficiency-- with tag SF's

eff data:  $0.816 \pm 0.032$

eff ttbar:  $0.839 \pm 0.005$

-----

Efficiency per Pt region

eff data pT[400-600]:  $0.8176 \pm 0.038$

eff ttbar pT[400-600]:  $0.837 \pm 0.006$

-----

eff data pT[600-800]:  $0.809 \pm 0.063$

eff ttbar pT[600-800]:  $0.847 \pm 0.013$

-----

eff data pT[800-Inf]:  $0.772 \pm 0.132$

eff ttbar pT[800-Inf]:  $0.868 \pm 0.032$

## without b tagging SF's

Efficiency-- without tag sf's

eff data:  $0.822 \pm 0.034$

eff ttbar:  $0.839 \pm 0.005$

-----

Efficiency per Pt region

eff data pT[400-600]:  $0.824 \pm 0.039$

eff ttbar pT[400-600]:  $0.837 \pm 0.006$

-----

eff data pT[600-800]:  $0.819 \pm 0.066$

eff ttbar pT[600-800]:  $0.847 \pm 0.013$

-----

eff data pT[800-Inf]:  $0.789 \pm 0.141$

eff ttbar pT[800-Inf]:  $0.868 \pm 0.032$

