

125. (General Topology HW - Dr. Tehrani) If d is a metric on X , and X is given the metric topology, show that $d : X \times X \rightarrow \mathbb{R}$ is continuous.

\mathbb{R}

Let (a, b) be an open interval in \mathbb{R} with $a < b$ WLOG. To show d is continuous, we need to show the set


$$A = d^{-1}((a, b)) = \{(x, x') \in X \times X \mid a < d(x, x') < b\}$$

is open in the product topology on $X \times X$. To do this, we will find an open set $Y = B_\epsilon(x) \times B_\epsilon(x')$ about each $(x, x') \in A$ such that $Y \subseteq A$.

Let $(x, x') \in A$. Let ϵ be small enough so that $B_{2\epsilon}(d(x, x'))$ is properly contained in (a, b) . (for later!)

Claim. $Y = B_\epsilon(x) \times B_\epsilon(x') \subseteq A$.

Proof. Let $(y, y') \in Y$. WTS $(y, y') \in A$.



We have:

$$\begin{aligned} \textcircled{1} \quad d(x, x') &\leq d(x, y) + d(y, x') \\ &\leq \underbrace{d(x, y)}_{< \epsilon} + d(y, y') + \underbrace{d(y', x')}_{< \epsilon} \\ &< d(y, y') + 2\epsilon \end{aligned}$$

and

$$\begin{aligned} \textcircled{2} \quad d(y, y') &\leq d(y, x) + d(x, y') \\ &\leq \underbrace{d(y, x)}_{< \epsilon} + d(x, x') + \underbrace{d(x', y')}_{< \epsilon} \\ &< d(x, x') + 2\epsilon \end{aligned}$$

draw triangles
(pretend $X = \mathbb{R}$)

(Δ inequality)

Now by our choice of ϵ ,

$$a < d(x, x') - 2\epsilon$$

$$< d(y, y') \quad (\text{by } \textcircled{1})$$

$$< d(x, x') + 2\epsilon \quad (\text{by } \textcircled{2})$$

$$< b \quad (\text{again by our choice of } \epsilon)$$

Therefore $(y, y') \in A$ by definition and the claim, ergo the whole question, is proved.