

REAL ANALYSIS FINAL
FALL 2021
RAÚL CURTO

1. (12 points) Define:
 - (a) (2 points) Uniform convergence of a sequence of functions on a set $E \subseteq \mathbb{R}$.
 - (b) (3 points) Variation of a function f on a closed, bounded interval $[a, b]$, with respect to a partition P of $[a, b]$.
 - (c) (2 points) Point of closure of a subset E of a metric space X .
 - (d) (2 points) Equicontinuity for a collection \mathcal{F} of real-valued functions on a metric space X .
 - (e) (3 points) State the Cantor intersection theorem.
2. (8 points) Let f be a continuous function on a closed, bounded, nondegenerate interval $[a, b]$ such that
 - (i) f is of bounded variation on $[a, b]$; and
 - (ii) f maps sets of measure zero to sets of measure zero; that is, for E a measurable subset of $[a, b]$, $m(E) = 0 \implies m(f(E)) = 0$.Prove that f is absolutely continuous on $[a, b]$.

3. (8 points) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a Borel function, and define

$$\mu(E) := m(f^{-1}(E)) \quad (E \subseteq \mathbb{R}, E \text{ Borel}).$$

Prove:

- (a) $\mu(E) \geq 0$ for all E Borel;
 - (b) μ is monotone;
 - (c) μ is countably additive;
 - (d) μ is not translation invariant; e.g., find a counterexample of a Borel function f and a set E such that $\mu(E + 1) \neq \mu(E)$, where $E + 1 := \{x + 1 \mid x \in E\}$.
4. (20 points) Determine if each of the following statements is true or false. If true, provide a proof; if false, provide a counterexample or show in some fashion why the statement is false. In either case, you are free to cite the textbook, and provide a rationale along the lines of "...by a proposition in Section a.b of Royden-Fitzpatrick."
 - (a) (4 points) True or false? Let \mathcal{F} be a collection of measurable functions on \mathbb{R} , and let

$$g := \sup_{f \in \mathcal{F}} f.$$

Then g is measurable on \mathbb{R} .

- (b) (6 points) True or false? Consider the normed linear space X of Riemann integrable functions on $[0, 1]$, with the norm $\|f\|_R := (R) \int_0^1 |f(x)| dx$. Then X is a Banach space. (Hint: the sequence $\{f_n\}_{n=1}^\infty$ of measurable functions given by

$$f_n(x) := \begin{cases} 1 & x = \frac{i}{k}, \text{ if } 1 \leq k \leq n \text{ and } 0 \leq i \leq k \\ 0 & \text{otherwise.} \end{cases}$$

The sequence $\{f_n\}$ is increasing and $f_n \rightarrow \chi_{\mathbb{Q}}$ as $n \rightarrow \infty$.)

- (c) (4 points) True or false? Recall that ℓ^∞ is the Banach space of real bounded sequences, equipped with the supremum norm. The space ℓ^∞ is separable. (Hint: $2^{\mathbb{N}}$ is not countable.)

- (d) (6 points) True or false? Let g be strictly increasing and absolutely continuous on a closed, bounded, nondegenerate interval $[a, b]$, and let \mathcal{O} be an open subset of (a, b) . Then

$$m(g(\mathcal{O})) = \int_{\mathcal{O}} g'.$$

5. (16 points) (a) (6 points) A Hilbert space \mathcal{H} may be characterized as a Banach space in which the so-called parallelogram law holds; that is, for each pair of vectors x and y in \mathcal{H} , we have

$$\|x + y\|_{\mathcal{H}}^2 + \|x - y\|_{\mathcal{H}}^2 = 2 \cdot \|x\|_{\mathcal{H}}^2 + 2 \cdot \|y\|_{\mathcal{H}}^2.$$

Using only characteristic functions of measurable subsets of $[0, 1]$, prove that $L^1([0, 1])$ is not a Hilbert space.

- (b) (10 points) (Embedding of ℓ^p in $L^p[1, \infty)$.) Let $1 \leq p < \infty$. For $a = (a_1, a_2, \dots) \in \ell^p$, define $f_a : [1, \infty) \rightarrow \mathbb{R}$ by

$$f_a(x) := \sum_{k=1}^{\infty} a_k \chi_{[k, k+1)}(x) \quad (x \in [1, \infty)).$$

- i. (2 points) Prove that $f_a \in L^p[1, \infty)$.
 - ii. (2 points) Prove that $\|f_a\|_p = \|a\|_p$.
 - iii. (3 points) Prove that the map $T : \ell^p \rightarrow L^p[1, \infty)$ given by $T(a) := f_a$ is linear and injective, but not surjective.
 - iv. (3 points) True or false? The range of T is dense in $L^p[1, \infty)$.
6. (6 points) Let f be a Lipschitz function on $[0, 1]$ and let g an absolutely continuous function on $[0, 1]$. Prove that $f \circ g$ is absolutely continuous on $[0, 1]$.
7. (16 points) (a) (4 points)
- (b) (6 points)
 - (c) (6 points)
8. (14 points) (a) (4 points)
- (b) (3 points)
 - (c) (4 points)
 - (d) (3 points)