

Key facts

- More than 1 million curable sexually transmitted infections (STIs) are acquired every day worldwide in people 15–49 years old, the majority of which are asymptomatic.
 - In 2020 there were an estimated 374 million new infections in people 15–49 years with 1 of 4 curable STIs: chlamydia, gonorrhoea, syphilis and trichomoniasis.
 - An estimated 8 million adults between 15 and 49 years old were infected with syphilis in 2022.
 - An estimated 520 million people aged 15–49 years (13%) worldwide have herpes simplex virus type 2 (HSV-2) infection, the main cause of genital herpes (1).
 - Human papillomavirus (HPV) infection is associated with over 311 000 cervical cancer deaths each year (2).
 - 1.1 million pregnant women were estimated to be infected with syphilis in 2022, resulting in over 390 000 adverse birth outcomes.
 - STIs have a direct impact on sexual and reproductive health through stigmatization, infertility, cancers and pregnancy complications and can increase the risk of HIV.
 - Drug resistance is a major threat to reducing the burden of STIs worldwide.
-

Overview

More than 30 different bacteria, viruses and parasites are known to be transmitted through sexual contact, including vaginal, anal and oral sex. Some STIs can also be transmitted from mother-to-child during pregnancy, childbirth and breastfeeding. Eight pathogens are linked to the greatest incidence of STIs. Of these, 4 are currently curable: syphilis, gonorrhoea, chlamydia and trichomoniasis. The other 4 are viral infections: hepatitis B, herpes simplex virus (HSV), HIV and human papillomavirus (HPV).

In addition, emerging outbreaks of new infections that can be acquired by sexual contact such as mpox, *Shigella sonnei*, *Neisseria meningitidis*, Ebola and Zika, as well as re-emergence of neglected STIs such as lymphogranuloma venereum. These herald increasing challenges in the provision of adequate services for STIs prevention and control.

Scope of the problem

STIs have a profound impact on sexual and reproductive health worldwide.

More than 1 million curable STIs are acquired every day. In 2020, WHO estimated 374 million new infections with 1 of 4 STIs: chlamydia (129 million),

gonorrhoea (82 million), syphilis (7.1 million) and trichomoniasis (156 million). More than 520 million people were estimated to be living with genital herpes in 2020, and an estimated 300 million women have an HPV infection, the primary cause of cervical cancer and anal cancer among men who have sex with men. In addition, updated [WHO estimates indicate](#) that 254 million people were living with chronic hepatitis B in 2022.

STIs can have serious consequences beyond the immediate impact of the infection itself.

- STIs like herpes, gonorrhoea and syphilis can increase the risk of HIV acquisition.
- Mother-to-child transmission of STIs can result in stillbirth, neonatal death, low-birth weight and prematurity, sepsis, neonatal conjunctivitis and congenital deformities.
- HPV infection causes cervical and other cancers.
- In 2022, hepatitis B resulted in an estimated 1.1 million deaths, mostly from cirrhosis and hepatocellular carcinoma (primary liver cancer).
- STIs such as gonorrhoea and chlamydia are major causes of pelvic inflammatory disease and infertility in women.

Prevention of STIs

When used correctly and consistently, condoms offer one of the most effective methods of protection against STIs, including HIV. Although highly effective, condoms do not offer protection for STIs that cause extra-genital ulcers (i.e., syphilis or genital herpes). When possible, condoms should be used in all vaginal and anal sex.

Safe and highly effective vaccines are available for 2 viral STIs: hepatitis B and HPV. These vaccines have represented major advances in STI prevention. By the end of 2023, the HPV vaccine had been introduced as part of routine immunization programmes in 140 countries, primarily high- and middle-income countries. To eliminate cervical cancer as a public health problem globally, high coverage targets for HPV vaccination, screening and treatment of precancerous lesions, and management of cancer must be reached by 2030 and maintained at this high level for decades.

Research to develop vaccines against genital herpes is advanced, with several vaccine candidates in early clinical development. There is mounting evidence suggesting that the vaccine to prevent meningitis (MenB) provides some cross-protection against gonorrhoea. More research into vaccines for chlamydia, gonorrhoea, syphilis and trichomoniasis are needed.

Other biomedical interventions to prevent some STIs include adult voluntary medical male circumcision, microbicides, and partner treatment. There are ongoing trials to evaluate the benefit of pre- and post-exposure prophylaxis of STIs and their potential safety weighed with antimicrobial resistance (AMR).

Diagnosis of STIs

STIs are often asymptomatic. When symptoms occur, they can be non-specific.

Accurate diagnostic tests for STIs (using molecular technology) are widely used in high-income countries. These are especially useful for the diagnosis of asymptomatic infections. However, they are largely unavailable in low- and middle-income countries (LMICs) for chlamydia and gonorrhoea. Even in countries where testing is available, it is often expensive and not widely accessible. In addition, the time it takes for results to be received is often long. As a result, follow-up can be impeded and care or treatment can be incomplete.

On the other hand, inexpensive, rapid tests are available for syphilis, hepatitis B and HIV. The rapid syphilis test and rapid dual HIV/syphilis tests are used in several resource-limited settings.

Several other rapid tests are under development and have the potential to improve STI diagnosis and treatment, especially in resource-limited settings.

Treatment of STIs

Effective treatment is currently available for several STIs.

- Three bacterial (chlamydia, gonorrhoea and syphilis) and one parasitic STIs (trichomoniasis) are generally curable with existing single-dose regimens of antibiotics.
- For herpes and HIV, the most effective medications available are antivirals that can modulate the course of the disease, though they cannot cure the disease.
- For hepatitis B, antivirals can help fighting the virus and slowing damage to the liver.

AMR of STIs – in particular gonorrhoea – has increased rapidly in recent years and has reduced treatment options. The [Gonococcal AMR Surveillance Programme \(GASP\)](#) has shown high rates of resistance to many antibiotics including quinolone, azithromycin and extended-spectrum cephalosporins, a last-line treatment (3).

AMR for other STIs, like *Mycoplasma genitalium*, also exist but are not systematically monitored.

STI case management

LMICs rely on identifying consistent, easily recognizable signs and symptoms to guide treatment, without the use of laboratory tests. This approach – syndromic management – often relies on clinical algorithms and allows health

workers to diagnose a specific infection based on observed syndromes (e.g., vaginal/urethral discharge, anogenital ulcers, etc). **Syndromic management** is simple, assures rapid, same-day treatment, and avoids expensive or unavailable diagnostic tests for patients with symptoms. However, this approach results in overtreatment and missed treatment as the majority of STIs are asymptomatic. Thus, WHO recommends countries to enhance syndromic management by gradually incorporating laboratory testing to support diagnosis. In settings where quality assured molecular assays are available, it is recommended to treat STIs based on laboratory tests. Moreover, STI screening strategies are essential for those at higher risk of infection, such sex workers, men who have sex with men, adolescents in some settings and pregnant women.

To interrupt transmission and prevent re-infection, treating sexual partners is an important component of STI case management.

Controlling the spread

Behaviour change is complex

Despite considerable efforts to identify simple interventions that can reduce risky sexual behaviour, behaviour change remains a complex challenge.

Information, education and counselling can improve people's ability to recognize the symptoms of STIs and increase the likelihood that they will seek care and encourage a sexual partner to do so. Unfortunately, lack of public awareness, lack of training among health workers, and long-standing, widespread stigma around STIs remain barriers to greater and more effective use of these interventions.

Health services for screening and treatment of STIs remain weak

People seeking screening and treatment for STIs face numerous problems. These include limited resources, stigmatization, poor quality of services and often out-of-pocket expenses.

Some populations with the highest rates of STIs – such as sex workers, men who have sex with men, people who inject drugs, prison inmates, mobile populations and adolescents in high burden countries for HIV – often do not have access to adequate and friendly health services.

In many settings, STI services are often neglected and underfunded. These problems lead to difficulties in providing testing for asymptomatic infections, insufficient number of trained personnel, limited laboratory capacity and inadequate supplies of appropriate medicines.

WHO response

Our work is currently guided by the [Global health sector strategy on HIV, Hepatitis and Sexually Transmitted Infections, 2022–2030](#). Within this framework, WHO:

- develops global targets, norms and standards for STI prevention, testing and treatment;
- supports the estimation and economic burden of STIs and the strengthening of STI surveillance;
- globally monitors AMR to gonorrhoea; and
- leads the setting of the global research agenda on STIs, including the development of diagnostic tests, vaccines and additional drugs for gonorrhoea and syphilis.

As part of its mission, WHO supports countries to:

- develop national strategic plans and guidelines;
- create an encouraging environment allowing individuals to discuss STIs, adopt safer sexual practices, and seek treatment;
- scale-up primary prevention (condom availability and use, etc.);
- increase integration of STI services within primary healthcare services;
- increase accessibility of people-centred quality STI care;
- facilitate adoption of point-of-care tests;
- enhance and scale-up health intervention for impact, such as hepatitis B and HPV vaccination, syphilis screening in priority populations;
- strengthen capacity to monitoring STIs trends; and
- monitor and respond to AMR in gonorrhoea.