

# 16-720B Homework 4 Write-up

Gu, Qiao

November 5, 2019

## Q1.1

Consider the point  $\mathbf{w}$  where the principle axes of the two cameras intersect, and we can see that  $\tilde{\mathbf{x}}_1 = [0, 0, 1]^T$  and  $\tilde{\mathbf{x}}_2 = [0, 0, 1]^T$  corresponding one point in 3D. Therefore

$$\tilde{\mathbf{x}}_2^T \mathbf{E} \tilde{\mathbf{x}}_1 = \begin{bmatrix} 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \mathbf{E}_{11} & \mathbf{E}_{12} & \mathbf{E}_{13} \\ \mathbf{E}_{21} & \mathbf{E}_{22} & \mathbf{E}_{23} \\ \mathbf{E}_{31} & \mathbf{E}_{32} & \mathbf{E}_{33} \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \mathbf{E}_{33} = 0 \quad (1)$$

Since two cameras are normalized, the intrinsic matrices for them are identity:  $\mathbf{K}_1 = \mathbf{K}_2 = \mathbf{I}$ . Then  $\mathbf{E} = \mathbf{K}_1^T \mathbf{F} \mathbf{K}_2 = \mathbf{E}$ . Therefore,  $\mathbf{E}_{33} = \mathbf{F}_{33} = 0$ .

### Q1.2

Suppose the cameras are normalized in the sense that their intrinsic matrices are both identity:  $\mathbf{K}_1 = \mathbf{K}_2 = \mathbf{I}$ .

Now that the translation and rotation from camera 1 to camera 2 are

$$\mathbf{R} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad \mathbf{t} = \begin{bmatrix} t_x \\ 0 \\ 0 \end{bmatrix} \quad (2)$$

And thus the essential matrix are

$$\mathbf{E} = \mathbf{t}_{\times} \mathbf{R} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -t_x \\ 0 & t_x & 0 \end{bmatrix} \quad (3)$$

Therefore for an epipolar line in camera 1  $\mathbf{l}_1^T \tilde{\mathbf{x}}_1 = 0$  and  $\tilde{\mathbf{x}}_2^T \mathbf{E} \tilde{\mathbf{x}}_1 = 0$ , where  $\tilde{\mathbf{x}}_2$  is a fixed point on the image plane of camera 2 resulting from the ray corresponding to the epipolar line, then we can see that

$$\mathbf{l}_1^T = \tilde{\mathbf{x}}_2^T \mathbf{E} = \begin{bmatrix} x_2 & y_2 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -t_x \\ 0 & t_x & 0 \end{bmatrix} = \begin{bmatrix} 0 & t_x & -t_x y_2 \end{bmatrix} \quad (4)$$

Similarly we can see that any epipolar line in camera 1 has  $\mathbf{l}_2^T = [0 \ -t_x \ t_x y_1]$ . Since the first elements in both  $\mathbf{l}_1$  and  $\mathbf{l}_2$  are zero, the epipolar lines are parallel to  $x$  axis.

### Q1.3

Assume  $(\mathbf{R}_i, \mathbf{t}_i)$  and  $(\mathbf{R}_j, \mathbf{t}_j)$  are the rotation and translation at time  $i$  and time  $j$ . Then for a point  $\mathbf{w}$  in the 3D world

$$\begin{aligned}\lambda_i \tilde{\mathbf{x}}_i &= \mathbf{R}_i \mathbf{w} + \mathbf{t}_i, & \lambda_j \tilde{\mathbf{x}}_j &= \mathbf{R}_j \mathbf{w} + \mathbf{t}_j \\ \Rightarrow \mathbf{w} &= \mathbf{R}_i^T (\lambda_i \tilde{\mathbf{x}}_i - \mathbf{t}_i) \\ \Rightarrow \lambda_j \tilde{\mathbf{x}}_j &= \mathbf{R}_j \mathbf{R}_i^T (\lambda_i \tilde{\mathbf{x}}_i - \mathbf{t}_i) + \mathbf{t}_j \\ \Rightarrow \lambda_j \tilde{\mathbf{x}}_j &= \mathbf{R}_j \mathbf{R}_i^T \lambda_i \tilde{\mathbf{x}}_i - \mathbf{R}_j \mathbf{R}_i^T \mathbf{t}_i + \mathbf{t}_j \\ \Rightarrow \lambda_j \tilde{\mathbf{x}}_j &= \lambda_i \mathbf{R}_{rel} \tilde{\mathbf{x}}_i + \mathbf{t}_{rel}\end{aligned}\tag{5}$$

Therefore

$$\mathbf{R}_{rel} = \mathbf{R}_j \mathbf{R}_i^T, \quad \mathbf{t}_{rel} = \mathbf{t}_j - \mathbf{R}_j \mathbf{R}_i^T \mathbf{t}_i\tag{6}$$

Q1.4