

# Imbalanced kinetic Alfvén wave turbulence

George Miloshevich, Thierry Passot, Pierre-Louis Sulem and Dimitri Laveder



Observatoire  
de la CÔTE d'AZUR

Festival de Théorie  
Aix en Provence, 2019

# Alfvén Wave turbulence in the Solar Wind (SW)

- Alfvén Waves (AW) play an important role in SW [1]


- Reflection leads to backward propagating waves
- SW turbulence results from these interactions.
- SW turbulence has mostly been studied using MHD




Schematics of a forward cascade to small scales

- SW is collisionless so kinetical treatment is ideal

- Microscale dispersive gyro-scales play a role
- Imbalanced turbulence [2] is of interest for (PSP)
- Direct Vlasov-Maxwell simulations are too costly
- Recourse to gyrofluids is more feasible [3]
- Describe the transition to imbalanced dispersive range



Parker Solar Probe (PSP)



Wind Data.

- [1] F. Sahraoui et al., PRL **102**, 231102 (2009)
- [2] R. T. Wicks et al., PRL **106**, 045001 (2011)
- [3] T. Passot and P. L. Sulem, JPP **85**, 905850301 (2019)

# Alfvén Wave turbulence in the Solar Wind (SW)

- Alfvén Waves (AW) play an important role in SW [1]


- Reflection leads to backward propagating waves
- SW turbulence results from these interactions.
- SW turbulence has mostly been studied using MHD




Schematics of a forward cascade to small scales

- SW is collisionless so kinetical treatment is ideal

- Microscale dispersive gyro-scales play a role
- Imbalanced turbulence [2] is of interest for (PSP)
- Direct Vlasov-Maxwell simulations are too costly
- Recourse to gyrofluids is more feasible [3]
- Describe the transition to imbalanced dispersive range



## Parker Solar Probe (PSP)




Wind Data.

- [1] F. Sahraoui et al., PRL **102**, 231102 (2009)
- [2] R. T. Wicks et al., PRL **106**, 045001 (2011)
- [3] T. Passot and P. L. Sulem, JPP **85**, 905850301 (2019)

# Alfvén Wave turbulence in the Solar Wind (SW)


- *Alfvén Waves (AW)* play an important role in SW <sup>[1]</sup>

- Reflection leads to backward propagating waves
- SW turbulence results from these interactions.
- SW turbulence has mostly been studied using MHD




Schematics of a forward cascade to small scales

- SW is collisionless so kinetical treatment is ideal
  - Microscale dispersive gyro-scales play a role
  - Imbalanced turbulence <sup>[2]</sup> is of interest for (PSP)
  - Direct Vlasov-Maxwell simulations are too costly
  - Recourse to gyrofluids is more feasible <sup>[3]</sup>
- Describe the transition to imbalanced dispersive range



Parker Solar Probe (PSP)




Wind Data.

- [1] F. Sahraoui et al., PRL **102**, 231102 (2009)
- [2] R. T. Wicks et al., PRL **106**, 045001 (2011)
- [3] T. Passot and P. L. Sulem, JPP **85**, 905850301 (2019)


# Alfvén Wave turbulence in the Solar Wind (SW)

- *Alfvén Waves (AW)* play an important role in SW <sup>[1]</sup>
  - Reflection leads to backward propagating waves
  - SW turbulence results from these interactions.
  - SW turbulence has mostly been studied using MHD




Schematics of a forward cascade to small scales

- SW is collisionless so kinetical treatment is ideal
  - Microscale dispersive *gyro-scales* play a role
  - *Imbalanced turbulence* <sup>[2]</sup> is of interest for (PSP)
  - Direct Vlasov-Maxwell simulations are too costly
  - Recourse to gyrofluids is more feasible <sup>[3]</sup>
- Describe the transition to imbalanced dispersive range



Parker Solar Probe (PSP)




Wind Data.

- [1] F. Sahraoui et al., PRL **102**, 231102 (2009)
- [2] R. T. Wicks et al., PRL **106**, 045001 (2011)
- [3] T. Passot and P. L. Sulem, JPP **85**, 905850301 (2019)


# Alfvén Wave turbulence in the Solar Wind (SW)

- Alfvén Waves (AW) play an important role in SW [1]
  - Reflection leads to backward propagating waves
  - SW turbulence results from these interactions.
  - SW turbulence has mostly been studied using MHD




Schematics of a forward cascade to small scales

- SW is collisionless so kinetical treatment is ideal
  - Microscale dispersive **gyro-scales** play a role
  - **Imbalanced turbulence** [2] is of interest for (PSP)
  - Direct Vlasov-Maxwell simulations are too costly
  - Recourse to gyrofluids is more feasible [3]
- Describe the transition to imbalanced dispersive range



Parker Solar Probe (PSP)




Wind Data.

- [1] F. Sahraoui et al., PRL **102**, 231102 (2009)
- [2] R. T. Wicks et al., PRL **106**, 045001 (2011)
- [3] T. Passot and P. L. Sulem, JPP **85**, 905850301 (2019)

# Alfvén Wave turbulence in the Solar Wind (SW)

- Alfvén Waves (AW) play an important role in SW [1]


- Reflection leads to backward propagating waves
- SW turbulence results from these interactions.
- SW turbulence has mostly been studied using MHD




Schematics of a forward cascade to small scales

- SW is collisionless so kinetical treatment is ideal

- Microscale dispersive **gyro-scales** play a role
- **Imbalanced turbulence** [2] is of interest for (PSP)
- Direct Vlasov-Maxwell simulations are too costly
- Recourse to gyrofluids is more feasible [3]
- Describe the transition to imbalanced dispersive range



Parker Solar Probe (PSP)




Wind Data.

- [1] F. Sahraoui et al., PRL **102**, 231102 (2009)
- [2] R. T. Wicks et al., PRL **106**, 045001 (2011)
- [3] T. Passot and P. L. Sulem, JPP **85**, 905850301 (2019)


# Alfvén Wave turbulence in the Solar Wind (SW)

- Alfvén Waves (AW) play an important role in SW [1]
  - Reflection leads to backward propagating waves
  - SW turbulence results from these interactions.
  - SW turbulence has mostly been studied using MHD




Schematics of a forward cascade to small scales

- SW is collisionless so kinetical treatment is ideal
  - Microscale dispersive **gyro-scales** play a role
  - **Imbalanced turbulence** [2] is of interest for (PSP)
    - Direct Vlasov-Maxwell simulations are too costly
    - Recourse to gyrofluids is more feasible [3]
- Describe the transition to imbalanced dispersive range



Parker Solar Probe (PSP)



Wind Data.


- [1] F. Sahraoui et al., PRL **102**, 231102 (2009)
- [2] R. T. Wicks et al., PRL **106**, 045001 (2011)
- [3] T. Passot and P. L. Sulem, JPP **85**, 905850301 (2019)



# Alfvén Wave turbulence in the Solar Wind (SW)

- Alfvén Waves (AW) play an important role in SW [1]

- Reflection leads to backward propagating waves
- SW turbulence results from these interactions.
- SW turbulence has mostly been studied using MHD




Schematics of a forward cascade to small scales


- SW is collisionless so kinetical treatment is ideal

- Microscale dispersive **gyro-scales** play a role
- **Imbalanced turbulence** [2] is of interest for (PSP)
- Direct Vlasov-Maxwell simulations are too costly
  - Recourse to gyrofluids is more feasible [3]

- Describe the transition to imbalanced dispersive range



Parker Solar Probe (PSP)




Wind Data.

- [1] F. Sahraoui et al., PRL **102**, 231102 (2009)
- [2] R. T. Wicks et al., PRL **106**, 045001 (2011)
- [3] T. Passot and P. L. Sulem, JPP **85**, 905850301 (2019)


# Alfvén Wave turbulence in the Solar Wind (SW)

- Alfvén Waves (AW) play an important role in SW [1]
  - Reflection leads to backward propagating waves
  - SW turbulence results from these interactions.
  - SW turbulence has mostly been studied using MHD




Schematics of a forward cascade to small scales

- SW is collisionless so kinetical treatment is ideal
  - Microscale dispersive **gyro-scales** play a role
  - **Imbalanced turbulence** [2] is of interest for (PSP)
  - Direct Vlasov-Maxwell simulations are too costly
  - Recourse to gyrofluids is more feasible [3]
- Describe the transition to imbalanced dispersive range



Parker Solar Probe (PSP)



Wind Data.

- [1] F. Sahraoui et al., PRL **102**, 231102 (2009)
- [2] R. T. Wicks et al., PRL **106**, 045001 (2011)
- [3] T. Passot and P. L. Sulem, JPP **85**, 905850301 (2019)


# Alfvén Wave turbulence in the Solar Wind (SW)

- Alfvén Waves (AW) play an important role in SW [1]
  - Reflection leads to backward propagating waves
  - SW turbulence results from these interactions.
  - SW turbulence has mostly been studied using MHD




Schematics of a forward cascade to small scales

- SW is collisionless so kinetical treatment is ideal
  - Microscale dispersive **gyro-scales** play a role
  - **Imbalanced turbulence** [2] is of interest for (PSP)
  - Direct Vlasov-Maxwell simulations are too costly
  - Recourse to gyrofluids is more feasible [3]
- Describe the transition to imbalanced dispersive range



Parker Solar Probe (PSP)



Wind Data.

- [1] F. Sahraoui et al., PRL **102**, 231102 (2009)
- [2] R. T. Wicks et al., PRL **106**, 045001 (2011)
- [3] T. Passot and P. L. Sulem, JPP **85**, 905850301 (2019)

- 1 Introduction
  - MHD Turbulence
  - Hamiltonian gyrofluid model
- 2 Influence of the dispersive range
  - Nonlinear diffusion equation
  - Landau damping
  - Inverse Cascade
- 3 Conclusion
  - Comparisons with 3D gyrofluid simulations
  - Future Work

- 1 Introduction
  - MHD Turbulence
  - Hamiltonian gyrofluid model
- 2 Influence of the dispersive range
  - Nonlinear diffusion equation
  - Landau damping
  - Inverse Cascade
- 3 Conclusion
  - Comparisons with 3D gyrofluid simulations
  - Future Work

- 1 Introduction
  - MHD Turbulence
  - Hamiltonian gyrofluid model
- 2 Influence of the dispersive range
  - Nonlinear diffusion equation
  - Landau damping
  - Inverse Cascade
- 3 Conclusion
  - Comparisons with 3D gyrofluid simulations
  - Future Work

- 1 Introduction
  - MHD Turbulence
  - Hamiltonian gyrofluid model
- 2 Influence of the dispersive range
  - Nonlinear diffusion equation
  - Landau damping
  - Inverse Cascade
- 3 Conclusion
  - Comparisons with 3D gyrofluid simulations
  - Future Work

# Weak balanced magnetohydrodynamic (MHD) turbulence

- Before discussing dispersive effects we review what is known about MHD

- Using Elsässer variables  $\mathbf{w}^\pm := \mathbf{v} \pm \mathbf{b}$ , MHD can be cast:

$$\partial_t \mathbf{w}^\pm \pm V_A \partial_z \mathbf{w}^\pm = -\mathbf{w}^\mp \cdot \nabla \mathbf{w}^\pm - \nabla P, \quad \nabla \cdot \mathbf{w}^\pm = 0, \quad P = -\nabla^{-2}(\nabla \mathbf{w}^+ : \nabla \mathbf{w}^-) \quad (1)$$

only counter propagating waves interact

- Assuming *weak balanced* cascade<sup>[4]</sup>, many  $w_\pm$  collisions before cascading  $\tau_{nl} \gg \tau_A$

$$w_{k_\perp}^+ = w_{k_\perp}^-, \quad w_{k_\perp}^+ w_{k_\perp}^- \propto k_\perp^{-1} \Rightarrow w_{k_\perp}^\pm \propto k_\perp^{-1/2} \quad (2)$$

- Criticism of weak MHD turbulence<sup>[5]</sup>: Because AW have  $\omega_p^\pm = \pm V_A k_\parallel$

$$\left. \begin{aligned} p + q = k \\ \omega_p^\pm + \omega_q^\mp = \omega_k^\pm \Rightarrow p_\parallel - q_\parallel = k_\parallel \end{aligned} \right\} \Rightarrow p_\parallel = k_\parallel, \quad q_\parallel = 0 \quad (3)$$

- So in MHD it involves scattering of AW off a 2D perturbation ( $q_\parallel = 0$ )

- As weak cascade proceeds eventually  $\tau_{nl} \sim k_\perp w_{k_\perp} \sim \tau_A \rightarrow$  and becomes *strong*

[4] C. S. Ng and A. Bhattacharjee, PoP 4, 605–610 (1997)

[5] A. A. Schekochihin, MHD Turbulence: A Biased Review, submitted to JPLP



# Weak balanced magnetohydrodynamic (MHD) turbulence

- Before discussing dispersive effects we review what is known about MHD
- Using Elsässer variables  $\mathbf{w}^\pm := \mathbf{v} \pm \mathbf{b}$ , MHD can be cast:

$$\partial_t \mathbf{w}^\pm \pm V_A \partial_z \mathbf{w}^\pm = -\mathbf{w}^\mp \cdot \nabla \mathbf{w}^\pm - \nabla P, \quad \nabla \cdot \mathbf{w}^\pm = 0, \quad P = -\nabla^{-2}(\nabla \mathbf{w}^+ : \nabla \mathbf{w}^-) \quad (1)$$

only counter propagating waves interact

- Assuming *weak balanced* cascade<sup>[4]</sup>, many  $w_\pm$  collisions before cascading  $\tau_{nl} \gg \tau_A$

$$w_{k_\perp}^+ = w_{k_\perp}^-, \quad w_{k_\perp}^+ w_{k_\perp}^- \propto k_\perp^{-1} \Rightarrow w_{k_\perp}^\pm \propto k_\perp^{-1/2} \quad (2)$$

- Criticism of weak MHD turbulence<sup>[5]</sup>: Because AW have  $\omega_p^\pm = \pm V_A k_\parallel$

$$\left. \begin{aligned} p + q &= k \\ \omega_p^\pm + \omega_q^\mp &= \omega_k^\pm \Rightarrow p_\parallel - q_\parallel = k_\parallel \end{aligned} \right\} \Rightarrow p_\parallel = k_\parallel, \quad q_\parallel = 0 \quad (3)$$

- So in MHD it involves scattering of AW off a 2D perturbation ( $q_\parallel = 0$ )
- As weak cascade proceeds eventually  $\tau_{nl} \sim k_\perp w_{k_\perp} \sim \tau_A \rightarrow$  and becomes *strong*

[4] C. S. Ng and A. Bhattacharjee, PoP 4, 605–610 (1997)

[5] A. A. Schekochihin, MHD Turbulence: A Biased Review, submitted to JPLP

# Weak balanced magnetohydrodynamic (MHD) turbulence

- Before discussing dispersive effects we review what is known about MHD
- Using Elsässer variables  $\mathbf{w}^\pm := \mathbf{v} \pm \mathbf{b}$ , MHD can be cast:

$$\partial_t \mathbf{w}^\pm \pm V_A \partial_z \mathbf{w}^\pm = -\mathbf{w}^\mp \cdot \nabla \mathbf{w}^\pm - \nabla P, \quad \nabla \cdot \mathbf{w}^\pm = 0, \quad P = -\nabla^{-2}(\nabla \mathbf{w}^+ : \nabla \mathbf{w}^-) \quad (1)$$

only counter propagating waves interact

- Assuming *weak balanced* cascade<sup>[4]</sup>, many  $w_\pm$  collisions before cascading  $\tau_{nl} \gg \tau_A$

$$w_{k_\perp}^+ = w_{k_\perp}^-, \quad w_{k_\perp}^+ w_{k_\perp}^- \propto k_\perp^{-1} \Rightarrow w_{k_\perp}^\pm \propto k_\perp^{-1/2} \quad (2)$$

- Criticism of weak MHD turbulence<sup>[5]</sup>: Because AW have  $\omega_p^\pm = \pm V_A k_\parallel$

$$\left. \begin{aligned} p + q &= k \\ \omega_p^\pm + \omega_q^\mp &= \omega_k^\pm \Rightarrow p_\parallel - q_\parallel = k_\parallel \end{aligned} \right\} \Rightarrow p_\parallel = k_\parallel, \quad q_\parallel = 0 \quad (3)$$

- So in MHD it involves scattering of AW off a 2D perturbation ( $q_\parallel = 0$ )
- As weak cascade proceeds eventually  $\tau_{nl} \sim k_\perp w_{k_\perp} \sim \tau_A \rightarrow$  and becomes *strong*

[4] C. S. Ng and A. Bhattacharjee, PoP 4, 605–610 (1997)

[5] A. A. Schekochihin, MHD Turbulence: A Biased Review, submitted to JPLP

# Weak balanced magnetohydrodynamic (MHD) turbulence

- Before discussing dispersive effects we review what is known about MHD
- Using Elsässer variables  $\mathbf{w}^\pm := \mathbf{v} \pm \mathbf{b}$ , MHD can be cast:

$$\partial_t \mathbf{w}^\pm \pm V_A \partial_z \mathbf{w}^\pm = -\mathbf{w}^\mp \cdot \nabla \mathbf{w}^\pm - \nabla P, \quad \nabla \cdot \mathbf{w}^\pm = 0, \quad P = -\nabla^{-2}(\nabla \mathbf{w}^+ : \nabla \mathbf{w}^-) \quad (1)$$

 only counter propagating waves interact

- Assuming *weak balanced* cascade<sup>[4]</sup>, many  $w_\pm$  collisions before cascading  $\tau_{nl} \gg \tau_A$

$$w_{k_\perp}^+ = w_{k_\perp}^-, \quad w_{k_\perp}^+ w_{k_\perp}^- \propto k_\perp^{-1} \Rightarrow w_{k_\perp}^\pm \propto k_\perp^{-1/2} \quad (2)$$

- Criticism of weak MHD turbulence<sup>[5]</sup>: Because AW have  $\omega_p^\pm = \pm V_A k_\parallel$

$$\left. \begin{aligned} p + q &= k \\ \omega_p^\pm + \omega_q^\mp &= \omega_k^\pm \Rightarrow p_\parallel - q_\parallel = k_\parallel \end{aligned} \right\} \Rightarrow p_\parallel = k_\parallel, \quad q_\parallel = 0 \quad (3)$$

- So in MHD it involves scattering of AW off a 2D perturbation ( $q_\parallel = 0$ )
- As weak cascade proceeds eventually  $\tau_{nl} \sim k_\perp w_{k_\perp} \sim \tau_A \rightarrow$  and becomes *strong*

[4] C. S. Ng and A. Bhattacharjee, PoP **4**, 605–610 (1997)

[5] A. A. Schekochihin, MHD Turbulence: A Biased Review, submitted to JPLP

# Weak balanced magnetohydrodynamic (MHD) turbulence

- Before discussing dispersive effects we review what is known about MHD
- Using Elsässer variables  $\mathbf{w}^\pm := \mathbf{v} \pm \mathbf{b}$ , MHD can be cast:

$$\partial_t \mathbf{w}^\pm \pm V_A \partial_z \mathbf{w}^\pm = -\mathbf{w}^\mp \cdot \nabla \mathbf{w}^\pm - \nabla P, \quad \nabla \cdot \mathbf{w}^\pm = 0, \quad P = -\nabla^{-2}(\nabla \mathbf{w}^+ : \nabla \mathbf{w}^-) \quad (1)$$

 only counter propagating waves interact

- Assuming *weak balanced* cascade<sup>[4]</sup>, many  $w_\pm$  collisions before cascading  $\tau_{nl} \gg \tau_A$

$$w_{k_\perp}^+ = w_{k_\perp}^-, \quad w_{k_\perp}^+ w_{k_\perp}^- \propto k_\perp^{-1} \Rightarrow w_{k_\perp}^\pm \propto k_\perp^{-1/2} \quad (2)$$

- Criticism of weak MHD turbulence<sup>[5]</sup>: Because AW have  $\omega_p^\pm = \pm V_A k_\parallel$

$$\left. \begin{aligned} p + q &= k \\ \omega_p^\pm + \omega_q^\mp &= \omega_k^\pm \Rightarrow p_\parallel - q_\parallel = k_\parallel \end{aligned} \right\} \Rightarrow p_\parallel = k_\parallel, \quad q_\parallel = 0 \quad (3)$$

- So in MHD it involves scattering of AW off a 2D perturbation ( $q_\parallel = 0$ )
- As weak cascade proceeds eventually  $\tau_{nl} \sim k_\perp w_{k_\perp} \sim \tau_A \rightarrow$  and becomes *strong*

[4] C. S. Ng and A. Bhattacharjee, PoP **4**, 605–610 (1997)

[5] A. A. Schekochihin, MHD Turbulence: A Biased Review, submitted to JPP 

# Weak balanced magnetohydrodynamic (MHD) turbulence

- Before discussing dispersive effects we review what is known about MHD
- Using Elsässer variables  $\mathbf{w}^\pm := \mathbf{v} \pm \mathbf{b}$ , MHD can be cast:

$$\partial_t \mathbf{w}^\pm \pm V_A \partial_z \mathbf{w}^\pm = -\mathbf{w}^\mp \cdot \nabla \mathbf{w}^\pm - \nabla P, \quad \nabla \cdot \mathbf{w}^\pm = 0, \quad P = -\nabla^{-2}(\nabla \mathbf{w}^+ : \nabla \mathbf{w}^-) \quad (1)$$

only counter propagating waves interact

- Assuming *weak balanced* cascade<sup>[4]</sup>, many  $w_\pm$  collisions before cascading  $\tau_{nl} \gg \tau_A$

$$w_{k_\perp}^+ = w_{k_\perp}^-, \quad w_{k_\perp}^+ w_{k_\perp}^- \propto k_\perp^{-1} \Rightarrow w_{k_\perp}^\pm \propto k_\perp^{-1/2} \quad (2)$$

- Criticism of weak MHD turbulence<sup>[5]</sup>: Because AW have  $\omega_p^\pm = \pm V_A k_\parallel$

$$\left. \begin{aligned} \mathbf{p} + \mathbf{q} &= \mathbf{k} \\ \omega_p^\pm + \omega_q^\mp &= \omega_k^\pm \Rightarrow p_\parallel - q_\parallel = k_\parallel \end{aligned} \right\} \Rightarrow p_\parallel = k_\parallel, \quad q_\parallel = 0 \quad (3)$$

- So in MHD it involves scattering of AW off a 2D perturbation ( $q_\parallel = 0$ )

- As weak cascade proceeds eventually  $\tau_{nl} \sim k_\perp w_{k_\perp} \sim \tau_A \rightarrow$  and becomes *strong*

[4] C. S. Ng and A. Bhattacharjee, PoP **4**, 605–610 (1997)

[5] A. A. Schekochihin, MHD Turbulence: A Biased Review, submitted to JPP

# Weak balanced magnetohydrodynamic (MHD) turbulence

- Before discussing dispersive effects we review what is known about MHD
- Using Elsässer variables  $\mathbf{w}^\pm := \mathbf{v} \pm \mathbf{b}$ , MHD can be cast:

$$\partial_t \mathbf{w}^\pm \pm V_A \partial_z \mathbf{w}^\pm = -\mathbf{w}^\mp \cdot \nabla \mathbf{w}^\pm - \nabla P, \quad \nabla \cdot \mathbf{w}^\pm = 0, \quad P = -\nabla^{-2}(\nabla \mathbf{w}^+ : \nabla \mathbf{w}^-) \quad (1)$$

 only counter propagating waves interact

- Assuming *weak balanced* cascade<sup>[4]</sup>, many  $w_\pm$  collisions before cascading  $\tau_{nl} \gg \tau_A$

$$w_{k_\perp}^+ = w_{k_\perp}^-, \quad w_{k_\perp}^+ w_{k_\perp}^- \propto k_\perp^{-1} \Rightarrow w_{k_\perp}^\pm \propto k_\perp^{-1/2} \quad (2)$$

- Criticism of weak MHD turbulence<sup>[5]</sup>: Because AW have  $\omega_p^\pm = \pm V_A k_\parallel$

$$\left. \begin{aligned} \mathbf{p} + \mathbf{q} &= \mathbf{k} \\ \omega_p^\pm + \omega_q^\mp &= \omega_k^\pm \Rightarrow p_\parallel - q_\parallel = k_\parallel \end{aligned} \right\} \Rightarrow p_\parallel = k_\parallel, \quad q_\parallel = 0 \quad (3)$$


- So in MHD it involves scattering of AW off a 2D perturbation ( $q_\parallel = 0$ )
- As weak cascade proceeds eventually  $\tau_{nl} \sim k_\perp w_{k_\perp} \sim \tau_A \rightarrow$  and becomes *strong*

[4] C. S. Ng and A. Bhattacharjee, PoP **4**, 605–610 (1997)

[5] A. A. Schekochihin, MHD Turbulence: A Biased Review, submitted to JPP 

# Pinning in case of imbalance and strong MHD turbulence

- *Imbalanced cascades* are much more common
- When  $w_{k_\perp}^+ \neq w_{k_\perp}^-$  additional criterion is required
- This is provided by the phenomenon of **pinning** [6][7]
- Weak turbulence proceeds to become strong
- Goldreich and Sridhar[8] introduced anisotropic theory postulating *critical balance*



Weak  $E_\pm(k_\perp) := w_{k_\perp}^2 / k_\perp$


$$k_\parallel V_A \sim k_\perp v_\perp \quad \epsilon \sim \frac{V_A^3}{L}, \quad \epsilon \sim \frac{v_\perp^2}{t_{cas}}, \quad t_{cas} \sim \frac{1}{k_\parallel V_A} \Rightarrow k_\parallel \sim k_\perp^{2/3} L^{-1/3} \quad (4)$$

- Role of dynamical alignment leading to  $-3/2$  energy spectrum[9]
- High resolution MHD simulations suggest  $-5/3$  energy spectrum[10]

- [6] R. Grappin et al., *Astron. and Astroph.* **126**, 51–58 (1983)
- [7] Y. Lithwick et al., *ApJ* **582**, 1220–1240 (2003)
- [8] P. Goldreich and S. Sridhar, *ApJ* **438**, 763–775 (1995)
- [9] S. Boldyrev et al., *ApJ* **699**, L39–L42 (2009)
- [10] A. Beresnyak, *ApJ* **784**, L20 (2014)

# Pinning in case of imbalance and strong MHD turbulence

- *Imbalanced cascades* are much more common
- When  $w_{k_\perp}^+ \neq w_{k_\perp}^-$  additional criterion is required
- This is provided by the phenomenon of **pinning** [6][7]
- Weak turbulence proceeds to become strong
- Goldreich and Sridhar[8] introduced anisotropic theory postulating *critical balance*



Weak  $E_\pm(k_\perp) := w_{k_\perp}^2 / k_\perp$

$$k_\parallel V_A \sim k_\perp v_\perp \quad \epsilon \sim \frac{V_A^3}{L}, \quad \epsilon \sim \frac{v_\perp^2}{t_{cas}}, \quad t_{cas} \sim \frac{1}{k_\parallel V_A} \Rightarrow k_\parallel \sim k_\perp^{2/3} L^{-1/3} \quad (4)$$


- Role of dynamical alignment leading to  $-3/2$  energy spectrum[9]
- High resolution MHD simulations suggest  $-5/3$  energy spectrum[10]

- [6] R. Grappin et al., *Astron. and Astroph.* **126**, 51–58 (1983)
- [7] Y. Lithwick et al., *ApJ* **582**, 1220–1240 (2003)
- [8] P. Goldreich and S. Sridhar, *ApJ* **438**, 763–775 (1995)
- [9] S. Boldyrev et al., *ApJ* **699**, L39–L42 (2009)
- [10] A. Beresnyak, *ApJ* **784**, L20 (2014)



# Pinning in case of imbalance and strong MHD turbulence

- *Imbalanced cascades* are much more common
- When  $w_{k_\perp}^+ \neq w_{k_\perp}^-$  additional criterion is required
- This is provided by the phenomenon of **pinning** [6][7]
- Weak turbulence proceeds to become strong
- Goldreich and Sridhar[8] introduced anisotropic theory postulating *critical balance*



Weak  $E_\pm(k_\perp) := w_{k_\perp}^2 / k_\perp$


$$k_\parallel V_A \sim k_\perp v_\perp \quad \epsilon \sim \frac{V_A^3}{L}, \quad \epsilon \sim \frac{v_\perp^2}{t_{cas}}, \quad t_{cas} \sim \frac{1}{k_\parallel V_A} \Rightarrow k_\parallel \sim k_\perp^{2/3} L^{-1/3} \quad (4)$$

- Role of dynamical alignment leading to  $-3/2$  energy spectrum[9]
- High resolution MHD simulations suggest  $-5/3$  energy spectrum[10]

- [6] R. Grappin et al., *Astron. and Astroph.* **126**, 51–58 (1983)
- [7] Y. Lithwick et al., *ApJ* **582**, 1220–1240 (2003)
- [8] P. Goldreich and S. Sridhar, *ApJ* **438**, 763–775 (1995)
- [9] S. Boldyrev et al., *ApJ* **699**, L39–L42 (2009)
- [10] A. Beresnyak, *ApJ* **784**, L20 (2014)

# Pinning in case of imbalance and strong MHD turbulence

- *Imbalanced cascades* are much more common
- When  $w_{k_\perp}^+ \neq w_{k_\perp}^-$  additional criterion is required
- This is provided by the phenomenon of **pinning** [6][7]
- Weak turbulence proceeds to become strong
- Goldreich and Sridhar<sup>[8]</sup> introduced anisotropic theory postulating *critical balance*



Weak  $E_\pm(k_\perp) := w_{k_\perp}^2 / k_\perp$


$$k_\parallel V_A \sim k_\perp v_\perp \quad \epsilon \sim \frac{V_A^3}{L}, \quad \epsilon \sim \frac{v_\perp^2}{t_{cas}}, \quad t_{cas} \sim \frac{1}{k_\parallel V_A} \Rightarrow k_\parallel \sim k_\perp^{2/3} L^{-1/3} \quad (4)$$

- Role of dynamical alignment leading to  $-3/2$  energy spectrum<sup>[9]</sup>
- High resolution MHD simulations suggest  $-5/3$  energy spectrum<sup>[10]</sup>

- [6] R. Grappin et al., *Astron. and Astroph.* **126**, 51–58 (1983)
- [7] Y. Lithwick et al., *ApJ* **582**, 1220–1240 (2003)
- [8] P. Goldreich and S. Sridhar, *ApJ* **438**, 763–775 (1995)
- [9] S. Boldyrev et al., *ApJ* **699**, L39–L42 (2009)
- [10] A. Beresnyak, *ApJ* **784**, L20 (2014)

# Pinning in case of imbalance and strong MHD turbulence

- *Imbalanced cascades* are much more common
- When  $w_{k_\perp}^+ \neq w_{k_\perp}^-$  additional criterion is required
- This is provided by the phenomenon of **pinning** [6][7]
- Weak turbulence proceeds to become strong
- Goldreich and Sridhar<sup>[8]</sup> introduced anisotropic theory postulating *critical balance*



Weak  $E_\pm(k_\perp) := w_{k_\perp}^2 / k_\perp$


$$k_\parallel V_A \sim k_\perp v_\perp \quad \epsilon \sim \frac{V_A^3}{L}, \quad \epsilon \sim \frac{v_\perp^2}{t_{cas}}, \quad t_{cas} \sim \frac{1}{k_\parallel V_A} \Rightarrow k_\parallel \sim k_\perp^{2/3} L^{-1/3} \quad (4)$$

- Role of dynamical alignment leading to  $-3/2$  energy spectrum<sup>[9]</sup>
- High resolution MHD simulations suggest  $-5/3$  energy spectrum<sup>[10]</sup>

- [6] R. Grappin et al., *Astron. and Astroph.* **126**, 51–58 (1983)
- [7] Y. Lithwick et al., *ApJ* **582**, 1220–1240 (2003)
- [8] P. Goldreich and S. Sridhar, *ApJ* **438**, 763–775 (1995)
- [9] S. Boldyrev et al., *ApJ* **699**, L39–L42 (2009)
- [10] A. Beresnyak, *ApJ* **784**, L20 (2014)

# Pinning in case of imbalance and strong MHD turbulence

- *Imbalanced cascades* are much more common
- When  $w_{k_\perp}^+ \neq w_{k_\perp}^-$  additional criterion is required
- This is provided by the phenomenon of **pinning** [6][7]
- Weak turbulence proceeds to become strong
- Goldreich and Sridhar<sup>[8]</sup> introduced anisotropic theory postulating *critical balance*



Weak  $E_\pm(k_\perp) := w_{k_\perp}^2 / k_\perp$


$$k_{\parallel} V_A \sim k_{\perp} v_{\perp} \quad \epsilon \sim \frac{V_A^3}{L}, \quad \epsilon \sim \frac{v_{\perp}^2}{t_{cas}}, \quad t_{cas} \sim \frac{1}{k_{\parallel} V_A} \Rightarrow k_{\parallel} \sim k_{\perp}^{2/3} L^{-1/3} \quad (4)$$

- Role of dynamical alignment leading to  $-3/2$  energy spectrum<sup>[9]</sup>
- High resolution MHD simulations suggest  $-5/3$  energy spectrum<sup>[10]</sup>

- [6] R. Grappin et al., *Astron. and Astroph.* **126**, 51–58 (1983)  
 [7] Y. Lithwick et al., *ApJ* **582**, 1220–1240 (2003)  
 [8] P. Goldreich and S. Sridhar, *ApJ* **438**, 763–775 (1995)  
 [9] S. Boldyrev et al., *ApJ* **699**, L39–L42 (2009)  
 [10] A. Beresnyak, *ApJ* **784**, L20 (2014)

# Pinning in case of imbalance and strong MHD turbulence

- *Imbalanced cascades* are much more common
- When  $w_{k_\perp}^+ \neq w_{k_\perp}^-$  additional criterion is required
- This is provided by the phenomenon of **pinning** [6][7]
- Weak turbulence proceeds to become strong
- Goldreich and Sridhar<sup>[8]</sup> introduced anisotropic theory postulating *critical balance*



Weak  $E_\pm(k_\perp) := w_{k_\perp}^2 / k_\perp$


$$k_{\parallel} V_A \sim k_{\perp} v_{\perp} \quad \epsilon \sim \frac{V_A^3}{L}, \quad \epsilon \sim \frac{v_{\perp}^2}{t_{cas}}, \quad t_{cas} \sim \frac{1}{k_{\parallel} V_A} \Rightarrow k_{\parallel} \sim k_{\perp}^{2/3} L^{-1/3} \quad (4)$$

- Role of dynamical alignment leading to  $-3/2$  energy spectrum<sup>[9]</sup>
- High resolution MHD simulations suggest  $-5/3$  energy spectrum<sup>[10]</sup>

- [6] R. Grappin et al., *Astron. and Astroph.* **126**, 51–58 (1983)
- [7] Y. Lithwick et al., *ApJ* **582**, 1220–1240 (2003)
- [8] P. Goldreich and S. Sridhar, *ApJ* **438**, 763–775 (1995)
- [9] S. Boldyrev et al., *ApJ* **699**, L39–L42 (2009)
- [10] A. Beresnyak, *ApJ* **784**, L20 (2014)

# Pinning in case of imbalance and strong MHD turbulence

- *Imbalanced cascades* are much more common
- When  $w_{k_\perp}^+ \neq w_{k_\perp}^-$  additional criterion is required
- This is provided by the phenomenon of **pinning** [6][7]
- Weak turbulence proceeds to become strong
- Goldreich and Sridhar<sup>[8]</sup> introduced anisotropic theory postulating *critical balance*



Weak  $E_\pm(k_\perp) := w_{k_\perp}^2 / k_\perp$

$$k_{\parallel} V_A \sim k_{\perp} v_{\perp} \quad \epsilon \sim \frac{V_A^3}{L}, \quad \epsilon \sim \frac{v_{\perp}^2}{t_{cas}}, \quad t_{cas} \sim \frac{1}{k_{\parallel} V_A} \Rightarrow k_{\parallel} \sim k_{\perp}^{2/3} L^{-1/3} \quad (4)$$

- Role of dynamical alignment leading to  $-3/2$  energy spectrum<sup>[9]</sup>
- High resolution MHD simulations suggest  $-5/3$  energy spectrum<sup>[10]</sup>

- [6] R. Grappin et al., *Astron. and Astroph.* **126**, 51–58 (1983)
- [7] Y. Lithwick et al., *ApJ* **582**, 1220–1240 (2003)
- [8] P. Goldreich and S. Sridhar, *ApJ* **438**, 763–775 (1995)
- [9] S. Boldyrev et al., *ApJ* **699**, L39–L42 (2009)
- [10] A. Beresnyak, *ApJ* **784**, L20 (2014)

# Addressing the dispersive range in Solar wind

- We leave the controversy of MHD and instead address dispersive scales
- Our description must involve small  $\beta_e$  relevant for the solar wind near the sun
- kinetic scales of interest are: ion  $\rho_i$  and sonic  $\rho_s$  Larmor radii

Relative kinetic plasma modes<sup>[11]</sup>


Electron beta

---


[11] V. Roytershteyn et al., ApJ **870**, 103 (2019)

# Addressing the dispersive range in Solar wind

- We leave the controversy of MHD and instead address dispersive scales
- Our description must involve small  $\beta_e$  relevant for the solar wind near the sun
- kinetic scales of interest are: ion  $\rho_i$  and sonic  $\rho_s$  Larmor radii



Relative kinetic plasma modes<sup>[11]</sup>




Electron beta

[11] V. Roytershteyn et al., ApJ 870, 103 (2019)




# Addressing the dispersive range in Solar wind

- We leave the controversy of MHD and instead address dispersive scales
- Our description must involve small  $\beta_e$  relevant for the solar wind near the sun
- kinetic scales of interest are: ion  $\rho_i$  and sonic  $\rho_s$  Larmor radii



Relative kinetic plasma modes<sup>[11]</sup>




Electron beta

[11] V. Roytershteyn et al., ApJ **870**, 103 (2019)

# Hamiltonian reduced gyrofluid model (GYRO)

- In order to understand **gyroscale physics** gyrokinetics <sup>[12]</sup>(5D) is often employed
- Numerical solutions of developed spectra are too costly even for this model
- Therefore, it is desirable to find some closure for a 3D gyrofluid model
- A Hamiltonian closure of this kind has been found recently <sup>[13]</sup>.
- Omitting effects of *electron inertia*  $m_e/m_i = 0$  the simplified equations read:



$$\partial_t N_e + V_A^2 \nabla_{\parallel} \Delta_{\perp} A_{\parallel} = \nabla(B_z - \varphi) \times \nabla N_e \cdot \hat{z} \quad (5)$$

gyrocenter density

$$\partial_t A_{\parallel} + \nabla_{\parallel}(\varphi - N_e - B_z) = 0, \quad (6)$$

$$\nabla_{\parallel} f := \partial f / \partial z + \nabla(A_{\parallel}) \times \nabla f \cdot \hat{z} \quad (7)$$

electrostatic pot.

magnetic potential

$$B_z = \hat{M}_1(\tau \Delta_{\perp})[\varphi] \quad N_e = -\hat{M}_2(\tau \Delta_{\perp})[\varphi] \quad (8)$$

Here  $\hat{M}_{1,2}$  are operators related to modified Bessel functions

Larmor gyration


$$\beta_e = 8\pi n_0 T_{0e} / B_0^2, \quad \Omega_i = eB_0 / m_i c, \quad \tau = T_{0i} / T_{0e} \quad (9)$$

[12] A. Brizard, Phys. Fluids B **4**, 1213–1228 (1992)

[13] T. Passot et al., PoP **25**, 042107 (2018)

# Hamiltonian reduced gyrofluid model (GYRO)

- In order to understand **gyroscale physics** gyrokinetics <sup>[12]</sup>(5D) is often employed
- Numerical solutions of developed spectra are too costly even for this model
- Therefore, it is desirable to find some closure for a 3D gyrofluid model
- A Hamiltonian closure of this kind has been found recently <sup>[13]</sup>.
- Omitting effects of *electron inertia*  $m_e/m_i = 0$  the simplified equations read:



gyrocenter density

$$\partial_t N_e + V_A^2 \nabla_{\parallel} \Delta_{\perp} A_{\parallel} = \nabla(B_z - \varphi) \times \nabla N_e \cdot \hat{z} \quad (5)$$

$$\partial_t A_{\parallel} + \nabla_{\parallel}(\varphi - N_e - B_z) = 0, \quad (6)$$

$$\nabla_{\parallel} f := \partial f / \partial z + \nabla(A_{\parallel}) \times \nabla f \cdot \hat{z} \quad (7)$$

electrostatic pot.

magnetic potential

$$B_z = \hat{M}_1(\tau \Delta_{\perp})[\varphi] \quad N_e = -\hat{M}_2(\tau \Delta_{\perp})[\varphi] \quad (8)$$

Here  $\hat{M}_{1,2}$  are operators related to modified Bessel functions

$$\beta_e = 8\pi n_0 T_{0e} / B_0^2, \quad \Omega_i = eB_0 / m_i c, \quad \tau = T_{0i} / T_{0e} \quad (9)$$


Larmor gyration

[12] A. Brizard, Phys. Fluids B **4**, 1213–1228 (1992)

[13] T. Passot et al., PoP **25**, 042107 (2018)

# Hamiltonian reduced gyrofluid model (GYRO)

- In order to understand **gyroscale physics** gyrokinetics <sup>[12]</sup>(5D) is often employed
- Numerical solutions of developed spectra are too costly even for this model
- Therefore, it is desirable to find some closure for a 3D gyrofluid model
- A Hamiltonian closure of this kind has been found recently <sup>[13]</sup>.
- Omitting effects of *electron inertia*  $m_e/m_i = 0$  the simplified equations read:



$$\partial_t N_e + V_A^2 \nabla_{\parallel} \Delta_{\perp} A_{\parallel} = \nabla(B_z - \varphi) \times \nabla N_e \cdot \hat{z} \quad (5)$$

gyrocenter density

$$\partial_t A_{\parallel} + \nabla_{\parallel}(\varphi - N_e - B_z) = 0, \quad (6)$$

$$\nabla_{\parallel} f := \partial f / \partial z + \nabla(A_{\parallel}) \times \nabla f \cdot \hat{z} \quad (7)$$

electrostatic pot.

magnetic potential

$$B_z = \hat{M}_1(\tau \Delta_{\perp})[\varphi] \quad N_e = -\hat{M}_2(\tau \Delta_{\perp})[\varphi] \quad (8)$$

Here  $\hat{M}_{1,2}$  are operators related to modified Bessel functions

Larmor gyration


$$\beta_e = 8\pi n_0 T_{0e} / B_0^2, \quad \Omega_i = eB_0 / m_i c, \quad \tau = T_{0i} / T_{0e} \quad (9)$$

[12] A. Brizard, Phys. Fluids B **4**, 1213–1228 (1992)

[13] T. Passot et al., PoP **25**, 042107 (2018)

# Hamiltonian reduced gyrofluid model (GYRO)

- In order to understand **gyroscale physics** gyrokinetics <sup>[12]</sup>(5D) is often employed
- Numerical solutions of developed spectra are too costly even for this model
- Therefore, it is desirable to find some closure for a 3D gyrofluid model
- A Hamiltonian closure of this kind has been found recently <sup>[13]</sup>.
- Omitting effects of *electron inertia*  $m_e/m_i = 0$  the simplified equations read:



$$\partial_t N_e + V_A^2 \nabla_{\parallel} \Delta_{\perp} A_{\parallel} = \nabla(B_z - \varphi) \times \nabla N_e \cdot \hat{z} \quad (5)$$

gyrocenter density

$$\partial_t A_{\parallel} + \nabla_{\parallel}(\varphi - N_e - B_z) = 0, \quad (6)$$

$$\nabla_{\parallel} f := \partial f / \partial z + \nabla(A_{\parallel}) \times \nabla f \cdot \hat{z} \quad (7)$$

electrostatic pot.

magnetic potential

$$B_z = \hat{M}_1(\tau \Delta_{\perp})[\varphi] \quad N_e = -\hat{M}_2(\tau \Delta_{\perp})[\varphi] \quad (8)$$

Here  $\hat{M}_{1,2}$  are operators related to modified Bessel functions

Larmor gyration


$$\beta_e = 8\pi n_0 T_{0e} / B_0^2, \quad \Omega_i = eB_0 / m_i c, \quad \tau = T_{0i} / T_{0e} \quad (9)$$

[12] A. Brizard, Phys. Fluids B **4**, 1213–1228 (1992)

[13] T. Passot et al., PoP **25**, 042107 (2018)

# Hamiltonian reduced gyrofluid model (GYRO)

- In order to understand **gyroscale physics** gyrokinetics <sup>[12]</sup> (5D) is often employed
- Numerical solutions of developed spectra are too costly even for this model
- Therefore, it is desirable to find some closure for a 3D gyrofluid model
- A Hamiltonian closure of this kind has been found recently <sup>[13]</sup>.
- Omitting effects of *electron inertia*  $m_e/m_i = 0$  the simplified equations read:



gyrocenter density

$$\partial_t N_e + V_A^2 \nabla_{\parallel} \Delta_{\perp} A_{\parallel} = \nabla(B_z - \varphi) \times \nabla N_e \cdot \hat{z} \quad (5)$$

$$\partial_t A_{\parallel} + \nabla_{\parallel}(\varphi - N_e - B_z) = 0, \quad (6)$$

$$\nabla_{\parallel} f := \partial f / \partial z + \nabla(A_{\parallel}) \times \nabla f \cdot \hat{z} \quad (7)$$

electrostatic pot.

magnetic potential

$$B_z = \hat{M}_1(\tau \Delta_{\perp})[\varphi] \quad N_e = -\hat{M}_2(\tau \Delta_{\perp})[\varphi] \quad (8)$$

Here  $\hat{M}_{1,2}$  are operators related to modified Bessel functions

$$\beta_e = 8\pi n_0 T_{0e} / B_0^2, \quad \Omega_i = eB_0 / m_i c, \quad \tau = T_{0i} / T_{0e} \quad (9)$$


Larmor gyration

[12] A. Brizard, Phys. Fluids B **4**, 1213–1228 (1992)

[13] T. Passot et al., PoP **25**, 042107 (2018)

# Hamiltonian reduced gyrofluid model (GYRO)

- In order to understand **gyroscale physics** gyrokinetics <sup>[12]</sup> (5D) is often employed
- Numerical solutions of developed spectra are too costly even for this model
- Therefore, it is desirable to find some closure for a 3D gyrofluid model
- A Hamiltonian closure of this kind has been found recently <sup>[13]</sup>.
- Omitting effects of *electron inertia*  $m_e/m_i = 0$  the simplified equations read:



gyrocenter density

$$\partial_t N_e + V_A^2 \nabla_{\parallel} \Delta_{\perp} A_{\parallel} = \nabla(B_z - \varphi) \times \nabla N_e \cdot \hat{z} \quad (5)$$

$$\partial_t A_{\parallel} + \nabla_{\parallel}(\varphi - N_e - B_z) = 0, \quad (6)$$

$$\nabla_{\parallel} f := \partial f / \partial z + \nabla(A_{\parallel}) \times \nabla f \cdot \hat{z} \quad (7)$$

electrostatic pot.

magnetic potential

$$B_z = \hat{M}_1(\tau \Delta_{\perp})[\varphi] \quad N_e = -\hat{M}_2(\tau \Delta_{\perp})[\varphi] \quad (8)$$

Here  $\hat{M}_{1,2}$  are operators related to modified Bessel functions

$$\beta_e = 8\pi n_0 T_{0e} / B_0^2, \quad \Omega_i = eB_0 / m_i c, \quad \tau = T_{0i} / T_{0e} \quad (9)$$


Larmor gyration

[12] A. Brizard, Phys. Fluids B **4**, 1213–1228 (1992)

[13] T. Passot et al., PoP **25**, 042107 (2018)

# Hamiltonian reduced gyrofluid model (GYRO)

- In order to understand **gyroscale physics** gyrokinetics <sup>[12]</sup>(5D) is often employed
- Numerical solutions of developed spectra are too costly even for this model
- Therefore, it is desirable to find some closure for a 3D gyrofluid model
- A Hamiltonian closure of this kind has been found recently <sup>[13]</sup>.
- Omitting effects of *electron inertia*  $m_e/m_i = 0$  the simplified equations read:



gyrocenter density

$$\partial_t N_e + V_A^2 \nabla_{\parallel} \Delta_{\perp} A_{\parallel} = \nabla(B_z - \varphi) \times \nabla N_e \cdot \hat{z} \quad (5)$$

$$\partial_t A_{\parallel} + \nabla_{\parallel}(\varphi - N_e - B_z) = 0, \quad (6)$$

$$\nabla_{\parallel} f := \partial f / \partial z + \nabla(A_{\parallel}) \times \nabla f \cdot \hat{z} \quad (7)$$

electrostatic pot.

magnetic potential

$$B_z = \hat{M}_1(\tau \Delta_{\perp})[\varphi] \quad N_e = -\hat{M}_2(\tau \Delta_{\perp})[\varphi] \quad (8)$$

Here  $\hat{M}_{1,2}$  are operators related to modified Bessel functions

$$\beta_e = 8\pi n_0 T_{0e} / B_0^2, \quad \Omega_i = eB_0 / m_i c, \quad \tau = T_{0i} / T_{0e} \quad (9)$$

Larmor gyration


[12] A. Brizard, Phys. Fluids B **4**, 1213–1228 (1992)

[13] T. Passot et al., PoP **25**, 042107 (2018)



# Hamiltonian reduced gyrofluid model (GYRO)

- In order to understand **gyroscale physics** gyrokinetics <sup>[12]</sup>(5D) is often employed
- Numerical solutions of developed spectra are too costly even for this model
- Therefore, it is desirable to find some closure for a 3D gyrofluid model
- A Hamiltonian closure of this kind has been found recently <sup>[13]</sup>.
- Omitting effects of *electron inertia*  $m_e/m_i = 0$  the simplified equations read:



$$\partial_t N_e + V_A^2 \nabla_{\parallel} \Delta_{\perp} A_{\parallel} = \nabla(B_z - \varphi) \times \nabla N_e \cdot \hat{z} \quad (5)$$

gyrocenter density

$$\partial_t A_{\parallel} + \nabla_{\parallel}(\varphi - N_e - B_z) = 0, \quad (6)$$

$$\nabla_{\parallel} f := \partial f / \partial z + \nabla(A_{\parallel}) \times \nabla f \cdot \hat{z} \quad (7)$$

electrostatic pot.

magnetic potential

$$B_z = \hat{M}_1(\tau \Delta_{\perp})[\varphi] \quad N_e = -\hat{M}_2(\tau \Delta_{\perp})[\varphi] \quad (8)$$

Here  $\hat{M}_{1,2}$  are operators related to modified Bessel functions

Larmor gyration


$$\beta_e = 8\pi n_0 T_{0e} / B_0^2, \quad \Omega_i = eB_0 / m_i c, \quad \tau = T_{0i} / T_{0e} \quad (9)$$

[12] A. Brizard, Phys. Fluids B **4**, 1213–1228 (1992)

[13] T. Passot et al., PoP **25**, 042107 (2018)

# Hamiltonian reduced gyrofluid model (GYRO)

- In order to understand **gyroscale physics** gyrokinetics <sup>[12]</sup>(5D) is often employed
- Numerical solutions of developed spectra are too costly even for this model
- Therefore, it is desirable to find some closure for a 3D gyrofluid model
- A Hamiltonian closure of this kind has been found recently <sup>[13]</sup>.
- Omitting effects of *electron inertia*  $m_e/m_i = 0$  the simplified equations read:



$$\partial_t N_e + V_A^2 \nabla_{\parallel} \Delta_{\perp} A_{\parallel} = \nabla(B_z - \varphi) \times \nabla N_e \cdot \hat{z} \quad (5)$$

gyrocenter density

$$\partial_t A_{\parallel} + \nabla_{\parallel}(\varphi - N_e - B_z) = 0, \quad (6)$$

$$\nabla_{\parallel} f := \partial f / \partial z + \nabla(A_{\parallel}) \times \nabla f \cdot \hat{z} \quad (7)$$

electrostatic pot.

magnetic potential

$$B_z = \hat{M}_1(\tau \Delta_{\perp})[\varphi] \quad N_e = -\hat{M}_2(\tau \Delta_{\perp})[\varphi] \quad (8)$$

Here  $\hat{M}_{1,2}$  are operators related to modified Bessel functions

Larmor gyration

$$\beta_e = 8\pi n_0 T_{0e} / B_0^2, \quad \Omega_i = eB_0 / m_i c, \quad \tau = T_{0i} / T_{0e} \quad (9)$$

[12] A. Brizard, Phys. Fluids B **4**, 1213–1228 (1992)

[13] T. Passot et al., PoP **25**, 042107 (2018)

# Kinetic Alfvén Waves & invariants of a gyrofluid model

- In nondimensional form the length is normalized to **sonic Larmor radius**  $\rho_s$ .

$$\rho_s = c_s/\Omega_i, \quad c_s := \sqrt{T_{oe}/m_i}, \quad \rho_i = \sqrt{2\tau}\rho_s, \quad \tau = T_{0i}/T_{0e} \quad (10)$$

- Introducing  $\mu^\pm := \hat{\Lambda}\phi \pm V_A A_\parallel$  - Elsässer potentials, where  $\hat{\Lambda} := \hat{\Lambda}(\Delta_\perp, \hat{M}_1, \hat{M}_2)$
- In MHD  $\tau \ll 1, \Delta_\perp \ll 1 \Rightarrow \hat{\Lambda} \rightarrow 1 \Rightarrow \mathbf{w}^\pm \rightarrow \nabla \mu^\pm \times \hat{\mathbf{z}}$  are like MHD Elsässer fields
- Model describes **Kinetic Alfvén Waves (KAW)**:  $\omega \rightarrow \alpha(\beta_e, \tau) k_\parallel k_\perp$  as  $k \rightarrow \infty$
- **Energy** and **generalized cross-helicity** conserved:

$$\mathcal{E} = \frac{1}{2} \int \left( V_A^2 |\nabla_\perp A_\parallel|^2 - N_e (\varphi - N_e - B_z) \right) d^3x, \quad (11)$$

$$\mathcal{E}_C = - \int N_e A_\parallel d^3x \rightarrow \begin{cases} - \int \omega_{iz} A_\parallel d^3x, & \text{In MHD} \\ \left( \frac{2}{\beta_e + \beta_i} + 1 \right) \int B_z A_\parallel d^3x, & \text{sub-ion} \end{cases} \quad (12)$$

**KAW** have  $V_{ph}$

We can also introduce energies associated with “backward” and “forward” propagating waves:  $\mathcal{E}_\pm := \frac{1}{2} \int (\mu_\pm^\pm)^2 d^3x$

# Kinetic Alfvén Waves & invariants of a gyrofluid model

- In nondimensional form the length is normalized to **sonic Larmor radius**  $\rho_s$ .

$$\rho_s = c_s / \Omega_i, \quad c_s := \sqrt{T_{0e} / m_i}, \quad \rho_i = \sqrt{2\tau} \rho_s, \quad \tau = T_{0i} / T_{0e} \quad (10)$$

- Introducing  $\mu^\pm := \hat{\Lambda} \phi \pm V_A A_\parallel$  - Elsässer potentials, where  $\hat{\Lambda} := \hat{\Lambda}(\Delta_\perp, \hat{M}_1, \hat{M}_2)$
- In MHD  $\tau \ll 1, \Delta_\perp \ll 1 \Rightarrow \hat{\Lambda} \rightarrow 1 \Rightarrow w^\pm \rightarrow \nabla \mu^\pm \times \hat{z}$  are like MHD Elsässer fields
- Model describes **Kinetic Alfvén Waves (KAW)**:  $\omega \rightarrow \alpha(\beta_e, \tau) k_\parallel k_\perp$  as  $k \rightarrow \infty$
- Energy** and **generalized cross-helicity** conserved:

$$\mathcal{E} = \frac{1}{2} \int \left( V_A^2 |\nabla_\perp A_\parallel|^2 - N_e (\varphi - N_e - B_z) \right) d^3x, \quad (11)$$

$$\mathcal{E}_C = - \int N_e A_\parallel d^3x \rightarrow \begin{cases} - \int \omega_{iz} A_\parallel d^3x, & \text{In MHD} \\ \left( \frac{2}{\beta_e + \beta_i} + 1 \right) \int B_z A_\parallel d^3x, & \text{sub-ion} \end{cases} \quad (12)$$

**KAW** have  $V_{ph}$

We can also introduce energies associated with “backward” and “forward” propagating waves:  $\mathcal{E}_\pm := \frac{1}{2} \int (\mu_\pm^\pm)^2 d^3x$

# Kinetic Alfvén Waves & invariants of a gyrofluid model

- In nondimensional form the length is normalized to **sonic Larmor radius**  $\rho_s$ .

$$\rho_s = c_s / \Omega_i, \quad c_s := \sqrt{T_{oe} / m_i}, \quad \rho_i = \sqrt{2\tau} \rho_s, \quad \tau = T_{0i} / T_{0e} \quad (10)$$

- Introducing  $\mu^\pm := \hat{\Lambda} \phi \pm V_A A_\parallel$  - Elsässer potentials, where  $\hat{\Lambda} := \hat{\Lambda}(\Delta_\perp, \hat{M}_1, \hat{M}_2)$
- In MHD  $\tau \ll 1, \Delta_\perp \ll 1 \Rightarrow \hat{\Lambda} \rightarrow 1 \Rightarrow \mathbf{w}^\pm \rightarrow \nabla \mu^\pm \times \hat{\mathbf{z}}$  are like MHD Elsässer fields

- Model describes **Kinetic Alfvén Waves (KAW)**:

$$\omega \rightarrow \alpha(\beta_e, \tau) k_\parallel k_\perp \text{ as } k \rightarrow \infty$$

- Energy** and **generalized cross-helicity** conserved:

$$\mathcal{E} = \frac{1}{2} \int \left( V_A^2 |\nabla_\perp A_\parallel|^2 - N_e (\varphi - N_e - B_z) \right) d^3 x, \quad (11)$$

$$\mathcal{E}_C = - \int N_e A_\parallel d^3 x \rightarrow \begin{cases} - \int \omega_{iz} A_\parallel d^3 x, & \text{In MHD} \\ \left( \frac{2}{\beta_e + \beta_i} + 1 \right) \int B_z A_\parallel d^3 x, & \text{sub-ion} \end{cases} \quad (12)$$

**KAW** have  $V_{ph}$

We can also introduce energies associated with “backward” and “forward” propagating waves:  $\mathcal{E}_\pm := \frac{1}{2} \int (\mu_\pm^\pm)^2 d^3 x$

# Kinetic Alfvén Waves & invariants of a gyrofluid model

- In nondimensional form the length is normalized to **sonic Larmor radius**  $\rho_s$ .

$$\rho_s = c_s / \Omega_i, \quad c_s := \sqrt{T_{0e} / m_i}, \quad \rho_i = \sqrt{2\tau} \rho_s, \quad \tau = T_{0i} / T_{0e} \quad (10)$$


- Introducing  $\mu^\pm := \hat{\Lambda} \phi \pm V_A A_\parallel$  - Elsässer potentials, where  $\hat{\Lambda} := \hat{\Lambda}(\Delta_\perp, \hat{M}_1, \hat{M}_2)$
- In MHD  $\tau \ll 1, \Delta_\perp \ll 1 \Rightarrow \hat{\Lambda} \rightarrow 1 \Rightarrow \mathbf{w}^\pm \rightarrow \nabla \mu^\pm \times \hat{\mathbf{z}}$  are like MHD Elsässer fields

- Model describes **Kinetic Alfvén Waves (KAW)**:

$$\omega \rightarrow \alpha(\beta_e, \tau) k_\parallel k_\perp \text{ as } k \rightarrow \infty$$

- Energy** and **generalized cross-helicity** conserved:

$$\mathcal{E} = \frac{1}{2} \int \left( V_A^2 |\nabla_\perp A_\parallel|^2 - N_e (\varphi - N_e - B_z) \right) d^3 x, \quad (11)$$



**KAW** have  $V_{ph}$

$$\mathcal{E}_C = - \int N_e A_\parallel d^3 x \rightarrow \begin{cases} - \int \omega_{iz} A_\parallel d^3 x, & \text{In MHD} \\ \left( \frac{2}{\beta_e + \beta_i} + 1 \right) \int B_z A_\parallel d^3 x, & \text{sub-ion} \end{cases} \quad (12)$$

We can also introduce energies associated with “backward” and “forward” propagating waves:  $\mathcal{E}_\pm := \frac{1}{2} \int (\mu_\pm^\pm)^2 d^3 x$

# Kinetic Alfvén Waves & invariants of a gyrofluid model

- In nondimensional form the length is normalized to **sonic Larmor radius**  $\rho_s$ .

$$\rho_s = c_s / \Omega_i, \quad c_s := \sqrt{T_{0e} / m_i}, \quad \rho_i = \sqrt{2\tau} \rho_s, \quad \tau = T_{0i} / T_{0e} \quad (10)$$


- Introducing  $\mu^\pm := \hat{\Lambda} \phi \pm V_A A_\parallel$  - Elsässer potentials, where  $\hat{\Lambda} := \hat{\Lambda}(\Delta_\perp, \hat{M}_1, \hat{M}_2)$
- In MHD  $\tau \ll 1, \Delta_\perp \ll 1 \Rightarrow \hat{\Lambda} \rightarrow 1 \Rightarrow \mathbf{w}^\pm \rightarrow \nabla \mu^\pm \times \hat{\mathbf{z}}$  are like MHD Elsässer fields

- Model describes **Kinetic Alfvén Waves (KAW)**:

$$\omega \rightarrow \alpha(\beta_e, \tau) k_\parallel k_\perp \text{ as } k \rightarrow \infty$$

- Energy** and **generalized cross-helicity** conserved:

$$\mathcal{E} = \frac{1}{2} \int \left( V_A^2 |\nabla_\perp A_\parallel|^2 - N_e (\varphi - N_e - B_z) \right) d^3x, \quad (11)$$



**KAW** have  $V_{ph}$

$$\mathcal{E}_C = - \int N_e A_\parallel d^3x \rightarrow \begin{cases} - \int \omega_{iz} A_\parallel d^3x, & \text{In MHD} \\ \left( \frac{2}{\beta_e + \beta_i} + 1 \right) \int B_z A_\parallel d^3x, & \text{sub-ion} \end{cases} \quad (12)$$

We can also introduce energies associated with “backward” and “forward” propagating waves:  $\mathcal{E}_\pm := \frac{1}{2} \int (\mu_\pm^\pm)^2 d^3x$

# Kinetic Alfvén Waves & invariants of a gyrofluid model

- In nondimensional form the length is normalized to **sonic Larmor radius**  $\rho_s$ .

$$\rho_s = c_s / \Omega_i, \quad c_s := \sqrt{T_{0e} / m_i}, \quad \rho_i = \sqrt{2\tau} \rho_s, \quad \tau = T_{0i} / T_{0e} \quad (10)$$


- Introducing  $\mu^\pm := \hat{\Lambda} \phi \pm V_A A_\parallel$  - Elsässer potentials, where  $\hat{\Lambda} := \hat{\Lambda}(\Delta_\perp, \hat{M}_1, \hat{M}_2)$
- In MHD  $\tau \ll 1, \Delta_\perp \ll 1 \Rightarrow \hat{\Lambda} \rightarrow 1 \Rightarrow \mathbf{w}^\pm \rightarrow \nabla \mu^\pm \times \hat{\mathbf{z}}$  are like MHD Elsässer fields

- Model describes **Kinetic Alfvén Waves (KAW)**:

$$\omega \rightarrow \alpha(\beta_e, \tau) k_\parallel k_\perp \text{ as } k \rightarrow \infty$$

- Energy** and **generalized cross-helicity** conserved:

$$\mathcal{E} = \frac{1}{2} \int \left( V_A^2 |\nabla_\perp A_\parallel|^2 - N_e (\varphi - N_e - B_z) \right) d^3x, \quad (11)$$



**KAW** have  $V_{ph}$

$$\mathcal{E}_C = - \int N_e A_\parallel d^3x \rightarrow \begin{cases} - \int \omega_{iz} A_\parallel d^3x, & \text{In MHD} \\ \left( \frac{2}{\beta_e + \beta_i} + 1 \right) \int B_z A_\parallel d^3x, & \text{sub-ion} \end{cases} \quad (12)$$

We can also introduce energies associated with “backward” and “forward” propagating waves:  $\mathcal{E}_\pm := \frac{1}{2} \int (\mu_\pm^\pm)^2 d^3x$



# Kinetic Alfvén Waves & invariants of a gyrofluid model

- In nondimensional form the length is normalized to **sonic Larmor radius**  $\rho_s$ .

$$\rho_s = c_s/\Omega_i, \quad c_s := \sqrt{T_{0e}/m_i}, \quad \rho_i = \sqrt{2\tau}\rho_s, \quad \tau = T_{0i}/T_{0e} \quad (10)$$


- Introducing  $\mu^\pm := \hat{\Lambda}\phi \pm V_A A_\parallel$  - Elsässer potentials, where  $\hat{\Lambda} := \hat{\Lambda}(\Delta_\perp, \hat{M}_1, \hat{M}_2)$
- In MHD  $\tau \ll 1, \Delta_\perp \ll 1 \Rightarrow \hat{\Lambda} \rightarrow 1 \Rightarrow \mathbf{w}^\pm \rightarrow \nabla\mu^\pm \times \hat{\mathbf{z}}$  are like MHD Elsässer fields

- Model describes **Kinetic Alfvén Waves (KAW)**:

$$\omega \rightarrow \alpha(\beta_e, \tau) k_\parallel k_\perp \text{ as } k \rightarrow \infty$$

- Energy** and **generalized cross-helicity** conserved:

$$\mathcal{E} = \frac{1}{2} \int \left( V_A^2 |\nabla_\perp A_\parallel|^2 - N_e (\varphi - N_e - B_z) \right) d^3x, \quad (11)$$



**KAW** have  $V_{ph}$

$$\mathcal{E}_C = - \int N_e A_\parallel d^3x \rightarrow \begin{cases} - \int \omega_{iz} A_\parallel d^3x, & \text{In MHD} \\ \left( \frac{2}{\beta_e + \beta_i} + 1 \right) \int B_z A_\parallel d^3x, & \text{sub-ion} \end{cases} \quad (12)$$

We can also introduce energies associated with “backward” and “forward” propagating waves:  $\mathcal{E}_\pm := \frac{1}{2} \int (\mu^\pm)^2 d^3x$

- 1 Introduction
  - MHD Turbulence
  - Hamiltonian gyrofluid model
- 2 Influence of the dispersive range
  - Nonlinear diffusion equation
  - Landau damping
  - Inverse Cascade
- 3 Conclusion
  - Comparisons with 3D gyrofluid simulations
  - Future Work

# Steps to derive wave kinetic equation: Ask me later?

- Using Fourier decomposition,  $a_{\mathbf{k}}^{\sigma_k} := e^{i\omega_k^{\sigma_k} t} k \mu_{\mathbf{k}}^{\sigma_k}$ ,  $\tau_{NL} \gg \omega^{-1}$ ,  $\sigma = \pm$  and  $\Omega_{\mathbf{k};pq}^{\sigma_k \sigma_p \sigma_q} = \omega_{\mathbf{k}}^{\sigma_k} - \omega_{\mathbf{p}}^{\sigma_p} - \omega_{\mathbf{q}}^{\sigma_q} = \sigma_k v_{ph}(k_{\perp}) k_{\parallel} - \sigma_p v_{ph}(p_{\perp}) p_{\parallel} - \sigma_q v_{ph}(q_{\perp}) q_{\parallel}$ . (13)

resonance condition one obtains **wave kinetic equation**:

$$\partial_t Q_{\mathbf{k}}^{\sigma} = 4\pi \int \sum_{\sigma_p, \sigma_q} \delta(\mathbf{k} + \mathbf{p} + \mathbf{q}) \delta(\Omega_{\mathbf{k}pq}^{\sigma \sigma_p \sigma_q}) V_{\mathbf{k}pq}^{\sigma \sigma_p \sigma_q} \left\{ \left( V_{\mathbf{p}q\mathbf{k}}^{\sigma_p \sigma_q \sigma} Q_{\mathbf{q}}^{\sigma_q} + V_{\mathbf{q}p\mathbf{k}}^{\sigma_q \sigma_p \sigma} Q_{\mathbf{p}}^{\sigma_p} \right) Q_{\mathbf{k}}^{\sigma} + V_{\mathbf{k}pq}^{\sigma \sigma_p \sigma_q} Q_{\mathbf{p}}^{\sigma_p} Q_{\mathbf{q}}^{\sigma_q} \right\} d\mathbf{p} d\mathbf{q}, \quad (14)$$

where

$$\langle a_{\mathbf{k}}^{\sigma_k} a_{\mathbf{k}'}^{\sigma_{k'}} \rangle =: Q_{\mathbf{k}}^{\sigma_k \sigma_{k'}} \delta(\mathbf{k} + \mathbf{k}') \Leftrightarrow Q_{\mathbf{k}}^{\sigma} = \frac{1}{\pi k_{\perp}} (E(k_{\perp}, k_{\parallel}) + \sigma v_{ph}(k_{\perp}) E_C(k_{\perp}, k_{\parallel})). \quad (15)$$

and introducing notation  $\xi := V_A/V_{ph}$ , the **vertex** is defined as

$$V_{\mathbf{k}pq}^{\sigma_k \sigma_p \sigma_q} := \frac{\hat{\mathbf{z}} \cdot (\mathbf{p} \times \mathbf{q})}{8 \xi(k_{\perp})} \left( \frac{\sigma_p}{\xi(p_{\perp})} - \frac{\sigma_q}{\xi(q_{\perp})} \right) \frac{\sigma_p \sigma_q}{k_{\perp} p_{\perp} q_{\perp}} \left( \sigma_k k_{\perp}^2 \xi(k_{\perp}) + \overset{\cup}{p.q.k} \right) \quad (16)$$

# Nonlinear diffusion equations (NDE) for weak turbulence

- Assuming isotropy in the transverse plane we can collapse

$$\int dp dq \delta(\mathbf{k} + \mathbf{p} + \mathbf{q}) \rightarrow 2\pi \int dp_{\parallel} dq_{\parallel} \delta(k_{\parallel} + p_{\parallel} + q_{\parallel}) \int_{\Delta_{k_{\perp}}} (1/\sin \alpha) dp_{\perp} dq_{\perp} \quad (17)$$

- In addition, assuming local interactions  $k_{\perp} \approx p_{\perp} \approx q_{\perp}$  and  $k_{\parallel} \approx p_{\parallel} \approx q_{\parallel}$

$$E^{\pm} = (E \pm V_{ph} E_C)/2$$

- Under these assumptions nonlinear diffusion equation for energy and cross-helicity has been derived<sup>[14]</sup>

$$\frac{\partial}{\partial t} \frac{E}{2} = \frac{\partial}{\partial k_{\perp}} \left\{ k_{\perp}^6 V_{ph} \sum_{r=\pm 1} E^{(-r)} \frac{\partial}{\partial k_{\perp}} \left( \frac{E^{(r)}}{k_{\perp}} \right) \right\} \quad (18)$$

$$\frac{\partial}{\partial t} \frac{E_C}{2} = \frac{\partial}{\partial k_{\perp}} \left\{ k_{\perp}^6 \sum_{r=\pm 1} (-1)^r E^{(-r)} \frac{\partial}{\partial k_{\perp}} \left( \frac{E^{(r)}}{k_{\perp}} \right) \right\} \quad (19)$$

[14] T. Passot and P. L. Sulem, JPP **85**, 905850301 (2019)

# Nonlinear diffusion equations (NDE) for weak turbulence

- Assuming isotropy in the transverse plane we can collapse

$$\int dp dq \delta(\mathbf{k} + \mathbf{p} + \mathbf{q}) \rightarrow 2\pi \int dp_{\parallel} dq_{\parallel} \delta(k_{\parallel} + p_{\parallel} + q_{\parallel}) \int_{\Delta_{k_{\perp}}} (1/\sin \alpha) dp_{\perp} dq_{\perp} \quad (17)$$

- In addition, assuming local interactions  $k_{\perp} \approx p_{\perp} \approx q_{\perp}$  and  $k_{\parallel} \approx p_{\parallel} \approx q_{\parallel}$

$$E^{\pm} = (E \pm V_{ph} E_C)/2$$

- Under these assumptions nonlinear diffusion equation for energy and cross-helicity has been derived<sup>[14]</sup>

$$\frac{\partial}{\partial t} \frac{E}{2} = \frac{\partial}{\partial k_{\perp}} \left\{ k_{\perp}^6 V_{ph} \sum_{r=\pm 1} E^{(-r)} \frac{\partial}{\partial k_{\perp}} \left( \frac{E^{(r)}}{k_{\perp}} \right) \right\} \quad (18)$$

$$\frac{\partial}{\partial t} \frac{E_C}{2} = \frac{\partial}{\partial k_{\perp}} \left\{ k_{\perp}^6 \sum_{r=\pm 1} (-1)^r E^{(-r)} \frac{\partial}{\partial k_{\perp}} \left( \frac{E^{(r)}}{k_{\perp}} \right) \right\} \quad (19)$$

[14] T. Passot and P. L. Sulem, JPP **85**, 905850301 (2019)

# Nonlinear diffusion equations (NDE) for weak turbulence

- Assuming isotropy in the transverse plane we can collapse

$$\int dp dq \delta(\mathbf{k} + \mathbf{p} + \mathbf{q}) \rightarrow 2\pi \int dp_{\parallel} dq_{\parallel} \delta(k_{\parallel} + p_{\parallel} + q_{\parallel}) \int_{\Delta_{k_{\perp}}} (1/\sin \alpha) dp_{\perp} dq_{\perp} \quad (17)$$

- In addition, assuming local interactions  $k_{\perp} \approx p_{\perp} \approx q_{\perp}$  and  $k_{\parallel} \approx p_{\parallel} \approx q_{\parallel}$

$$E^{\pm} = (E \pm V_{ph} E_C)/2$$

- Under these assumptions nonlinear diffusion equation for energy and cross-helicity has been derived<sup>[14]</sup>

$$\frac{\partial}{\partial t} \frac{E}{2} = \frac{\partial}{\partial k_{\perp}} \left\{ k_{\perp}^6 V_{ph} \sum_{r=\pm 1} E^{(-r)} \frac{\partial}{\partial k_{\perp}} \left( \frac{E^{(r)}}{k_{\perp}} \right) \right\} \quad (18)$$

$$\frac{\partial}{\partial t} \frac{E_C}{2} = \frac{\partial}{\partial k_{\perp}} \left\{ k_{\perp}^6 \sum_{r=\pm 1} (-1)^r E^{(-r)} \frac{\partial}{\partial k_{\perp}} \left( \frac{E^{(r)}}{k_{\perp}} \right) \right\} \quad (19)$$

[14] T. Passot and P. L. Sulem, JPP **85**, 905850301 (2019)

# Phenomenological extension to strong turbulence

- As turbulence proceeds to small scales it ceases to be weak:  $\tau_{NL} \sim \tau_L$
- In strong turbulence **parallel transfer** is non-negligible  $\tilde{k}_{\parallel} \neq \text{const}$
- In weak turbulence the transfer time is:  $\tau_{tr,w}^{\pm} = (k_{\perp}^3 V_{ph} E^{\mp})^{-1}$ ,  $\omega_L := V_{ph} \tilde{k}_{\parallel}$
- In strong turbulence transfer time is modified consistently with *critical balance*

$$\tau_{tr,w}^{\pm} = (\tau_{NL}^{\pm})^2 \omega_L \Rightarrow \tau_{NL}^{\pm} \sim (k_{\perp}^3 V_{ph}^2 E^{\mp})^{-1/2} \sim \tau_L = (V_{ph} \tilde{k}_{\parallel})^{-1} \quad (20)$$

- The + wave affects the correlation length of – so

$$\tilde{k}_{\parallel}^{(r)} = (k_{\perp}^3 E^{(r)})^{1/2} \left( \frac{E^{+}}{E^{-}} \right)^{(1-\chi)r/4}, \quad r = \pm 1 \quad (21)$$

- When  $\chi = 0$  both  $E_{+}$  and  $E_{-}$  cascades are strong.
- It is reasonable to treat  $E_{+}$  weakly <sup>[15]</sup>, where  $\chi = 1$
- DNS MHD simulations <sup>[16]</sup> are consistent with  $\chi \sim 1/4$

The effect of  $\chi$  in MHD

[15] B. D. G. Chandran, The Astrophysical Journal **685**, 646–658 (2008)

[16] A. Beresnyak and A. Lazarian, ApJ **682**, 1070–1075 (2008)

# Phenomenological extension to strong turbulence

- As turbulence proceeds to small scales it ceases to be weak:  $\tau_{NL} \sim \tau_L$
- In strong turbulence **parallel transfer** is non-negligible  $\tilde{k}_{\parallel} \neq \text{const}$
- In weak turbulence the transfer time is:  $\tau_{tr,w}^{\pm} = (k_{\perp}^3 V_{ph} E^{\mp})^{-1}$ ,  $\omega_L := V_{ph} \tilde{k}_{\parallel}$
- In strong turbulence transfer time is modified consistently with *critical balance*

$$\tau_{tr,w}^{\pm} = (\tau_{NL}^{\pm})^2 \omega_L \Rightarrow \tau_{NL}^{\pm} \sim (k_{\perp}^3 V_{ph}^2 E^{\mp})^{-1/2} \sim \tau_L = (V_{ph} \tilde{k}_{\parallel})^{-1} \quad (20)$$

- The + wave affects the correlation length of – so

$$\tilde{k}_{\parallel}^{(r)} = (k_{\perp}^3 E^{(r)})^{1/2} \left( \frac{E^{+}}{E^{-}} \right)^{(1-\chi)r/4}, \quad r = \pm 1 \quad (21)$$

- When  $\chi = 0$  both  $E_{+}$  and  $E_{-}$  cascades are strong.
- It is reasonable to treat  $E_{+}$  weakly <sup>[15]</sup>, where  $\chi = 1$
- DNS MHD simulations <sup>[16]</sup> are consistent with  $\chi \sim 1/4$

The effect of  $\chi$  in MHD

[15] B. D. G. Chandran, The Astrophysical Journal **685**, 646–658 (2008)

[16] A. Beresnyak and A. Lazarian, ApJ **682**, 1070–1075 (2008)



# Phenomenological extension to strong turbulence

- As turbulence proceeds to small scales it ceases to be weak:  $\tau_{NL} \sim \tau_L$
- In strong turbulence **parallel transfer** is non-negligible  $\tilde{k}_{\parallel} \neq \text{const}$
- In weak turbulence the transfer time is:  $\tau_{tr,w}^{\pm} = (k_{\perp}^3 V_{ph} E^{\mp})^{-1}$ ,  $\omega_L := V_{ph} \tilde{k}_{\parallel}$
- In strong turbulence transfer time is modified consistently with *critical balance*

$$\tau_{tr,w}^{\pm} = (\tau_{NL}^{\pm})^2 \omega_L \Rightarrow \tau_{NL}^{\pm} \sim (k_{\perp}^3 V_{ph}^2 E^{\mp})^{-1/2} \sim \tau_L = (V_{ph} \tilde{k}_{\parallel})^{-1} \quad (20)$$

- The + wave affects the correlation length of – so

$$\tilde{k}_{\parallel}^{(r)} = (k_{\perp}^3 E^{(r)})^{1/2} \left( \frac{E^{+}}{E^{-}} \right)^{(1-\chi)r/4}, \quad r = \pm 1 \quad (21)$$

- When  $\chi = 0$  both  $E_{+}$  and  $E_{-}$  cascades are strong.
- It is reasonable to treat  $E_{+}$  weakly <sup>[15]</sup>, where  $\chi = 1$
- DNS MHD simulations <sup>[16]</sup> are consistent with  $\chi \sim 1/4$

The effect of  $\chi$  in MHD

[15] B. D. G. Chandran, The Astrophysical Journal **685**, 646–658 (2008)

[16] A. Beresnyak and A. Lazarian, ApJ **682**, 1070–1075 (2008)

# Phenomenological extension to strong turbulence

- As turbulence proceeds to small scales it ceases to be weak:  $\tau_{NL} \sim \tau_L$
- In strong turbulence **parallel transfer** is non-negligible  $\tilde{k}_{\parallel} \neq \text{const}$
- In weak turbulence the transfer time is:  $\tau_{tr,w}^{\pm} = (k_{\perp}^3 V_{ph} E^{\mp})^{-1}$ ,  $\omega_L := V_{ph} \tilde{k}_{\parallel}$
- In strong turbulence transfer time is modified consistently with *critical balance*  

$$\tau_{tr,w}^{\pm} = (\tau_{NL}^{\pm})^2 \omega_L \Rightarrow \tau_{NL}^{\pm} \sim (k_{\perp}^3 V_{ph}^2 E^{\mp})^{-1/2} \sim \tau_L = (V_{ph} \tilde{k}_{\parallel})^{-1} \quad (20)$$
- The + wave affects the correlation length of – so

$$\tilde{k}_{\parallel}^{(r)} = (k_{\perp}^3 E^{(r)})^{1/2} \left( \frac{E^{+}}{E^{-}} \right)^{(1-\chi)r/4}, \quad r = \pm 1 \quad (21)$$

- When  $\chi = 0$  both  $E_{+}$  and  $E_{-}$  cascades are strong.
- It is reasonable to treat  $E_{+}$  weakly <sup>[15]</sup>, where  $\chi = 1$
- DNS MHD simulations <sup>[16]</sup> are consistent with  $\chi \sim 1/4$

The effect of  $\chi$  in MHD

[15] B. D. G. Chandran, The Astrophysical Journal **685**, 646–658 (2008)

[16] A. Beresnyak and A. Lazarian, ApJ **682**, 1070–1075 (2008)

# Phenomenological extension to strong turbulence

- As turbulence proceeds to small scales it ceases to be weak:  $\tau_{NL} \sim \tau_L$
- In strong turbulence **parallel transfer** is non-negligible  $\tilde{k}_{\parallel} \neq \text{const}$
- In weak turbulence the transfer time is:  $\tau_{tr,w}^{\pm} = (k_{\perp}^3 V_{ph} E^{\mp})^{-1}$ ,  $\omega_L := V_{ph} \tilde{k}_{\parallel}$
- In strong turbulence transfer time is modified consistently with *critical balance*  

$$\tau_{tr,w}^{\pm} = (\tau_{NL}^{\pm})^2 \omega_L \Rightarrow \tau_{NL}^{\pm} \sim (k_{\perp}^3 V_{ph}^2 E^{\mp})^{-1/2} \sim \tau_L = (V_{ph} \tilde{k}_{\parallel})^{-1} \quad (20)$$
- The + wave affects the correlation length of – so

$$\tilde{k}_{\parallel}^{(r)} = (k_{\perp}^3 E^{(r)})^{1/2} \left( \frac{E^{+}}{E^{-}} \right)^{(1-r)\chi/4}, \quad r = \pm 1 \quad (21)$$

- When  $\chi = 0$  both  $E_{+}$  and  $E_{-}$  cascades are strong.
- It is reasonable to treat  $E_{+}$  weakly [15], where  $\chi = 1$
- DNS MHD simulations [16] are consistent with  $\chi \sim 1/4$

The effect of  $\chi$  in MHD

[15] B. D. G. Chandran, The Astrophysical Journal **685**, 646–658 (2008)

[16] A. Beresnyak and A. Lazarian, ApJ **682**, 1070–1075 (2008)

# Phenomenological extension to strong turbulence

- As turbulence proceeds to small scales it ceases to be weak:  $\tau_{NL} \sim \tau_L$
- In strong turbulence **parallel transfer** is non-negligible  $\tilde{k}_{\parallel} \neq \text{const}$
- In weak turbulence the transfer time is:  $\tau_{tr,w}^{\pm} = (k_{\perp}^3 V_{ph} E^{\mp})^{-1}$ ,  $\omega_L := V_{ph} \tilde{k}_{\parallel}$
- In strong turbulence transfer time is modified consistently with *critical balance*  

$$\tau_{tr,w}^{\pm} = (\tau_{NL}^{\pm})^2 \omega_L \Rightarrow \tau_{NL}^{\pm} \sim (k_{\perp}^3 V_{ph}^2 E^{\mp})^{-1/2} \sim \tau_L = (V_{ph} \tilde{k}_{\parallel})^{-1} \quad (20)$$
- The + wave affects the correlation length of – so

$$\tilde{k}_{\parallel}^{(r)} = (k_{\perp}^3 E^{(r)})^{1/2} \left( \frac{E^{+}}{E^{-}} \right)^{(1-r)\chi/4}, \quad r = \pm 1 \quad (21)$$

- When  $\chi = 0$  both  $E_{+}$  and  $E_{-}$  cascades are strong.
- It is reasonable to treat  $E_{+}$  weakly <sup>[15]</sup>, where  $\chi = 1$
- DNS MHD simulations <sup>[16]</sup> are consistent with  $\chi \sim 1/4$

The effect of  $\chi$  in MHD


[15] B. D. G. Chandran, The Astrophysical Journal **685**, 646–658 (2008)

[16] A. Beresnyak and A. Lazarian, ApJ **682**, 1070–1075 (2008)

# Phenomenological extension to strong turbulence

- As turbulence proceeds to small scales it ceases to be weak:  $\tau_{NL} \sim \tau_L$
- In strong turbulence **parallel transfer** is non-negligible  $\tilde{k}_{\parallel} \neq \text{const}$
- In weak turbulence the transfer time is:  $\tau_{tr,w}^{\pm} = (k_{\perp}^3 V_{ph} E^{\pm})^{-1}$ ,  $\omega_L := V_{ph} \tilde{k}_{\parallel}$
- In strong turbulence transfer time is modified consistently with *critical balance*

$$\tau_{tr,w}^{\pm} = (\tau_{NL}^{\pm})^2 \omega_L \Rightarrow \tau_{NL}^{\pm} \sim (k_{\perp}^3 V_{ph}^2 E^{\pm})^{-1/2} \sim \tau_L = (V_{ph} \tilde{k}_{\parallel})^{-1} \quad (20)$$



- The + wave affects the correlation length of – so

$$\tilde{k}_{\parallel}^{(r)} = (k_{\perp}^3 E^{(r)})^{1/2} \left( \frac{E^{+}}{E^{-}} \right)^{(1-r)\chi/4}, \quad r = \pm 1 \quad (21)$$

- When  $\chi = 0$  both  $E_{+}$  and  $E_{-}$  cascades are strong.
- It is reasonable to treat  $E_{+}$  weakly <sup>[15]</sup>, where  $\chi = 1$
- DNS MHD simulations <sup>[16]</sup> are consistent with  $\chi \sim 1/4$

The effect of  $\chi$  in MHD


[15] B. D. G. Chandran, The Astrophysical Journal **685**, 646–658 (2008)

[16] A. Beresnyak and A. Lazarian, ApJ **682**, 1070–1075 (2008)

# Phenomenological extension to strong turbulence

- As turbulence proceeds to small scales it ceases to be weak:  $\tau_{NL} \sim \tau_L$
- In strong turbulence **parallel transfer** is non-negligible  $\tilde{k}_{\parallel} \neq \text{const}$
- In weak turbulence the transfer time is:  $\tau_{tr,w}^{\pm} = (k_{\perp}^3 V_{ph} E^{\pm})^{-1}$ ,  $\omega_L := V_{ph} \tilde{k}_{\parallel}$
- In strong turbulence transfer time is modified consistently with *critical balance*

$$\tau_{tr,w}^{\pm} = (\tau_{NL}^{\pm})^2 \omega_L \Rightarrow \tau_{NL}^{\pm} \sim (k_{\perp}^3 V_{ph}^2 E^{\pm})^{-1/2} \sim \tau_L = (V_{ph} \tilde{k}_{\parallel})^{-1} \quad (20)$$



The effect of  $\chi$  in MHD

- The + wave affects the correlation length of – so

$$\tilde{k}_{\parallel}^{(r)} = (k_{\perp}^3 E^{(r)})^{1/2} \left( \frac{E^{+}}{E^{-}} \right)^{(1-r)\chi/4}, \quad r = \pm 1 \quad (21)$$

- When  $\chi = 0$  both  $E_{+}$  and  $E_{-}$  cascades are strong.
- It is reasonable to treat  $E_{+}$  weakly <sup>[15]</sup>, where  $\chi = 1$
- DNS MHD simulations <sup>[16]</sup> are consistent with  $\chi \sim 1/4$

[15] B. D. G. Chandran, The Astrophysical Journal **685**, 646–658 (2008)


[16] A. Beresnyak and A. Lazarian, ApJ **682**, 1070–1075 (2008)

# Numerical scheme for NDE - Ask me later?

$$\frac{1}{2} \frac{\partial}{\partial t} \begin{pmatrix} E \\ E_C \end{pmatrix} = \frac{\partial}{\partial k_{\perp}} \left\{ k_{\perp}^6 \begin{pmatrix} V_{ph} \\ 1 \end{pmatrix} \sum_{r=\pm 1} \begin{pmatrix} 1 \\ (-1)^r \end{pmatrix} \frac{E^{(-r)}}{\tilde{k}_{\parallel}^{\pm}} \frac{\partial}{\partial k_{\perp}} \left( \frac{E^{(r)}}{k_{\perp}} \right) \right\} =: \frac{\partial}{\partial k_{\perp}} \begin{pmatrix} \varepsilon \\ \eta \end{pmatrix} \quad (22)$$


- We are modeling coupled nonlinear diffusion equations of **Leith** type.
  - Code for Gravitational wave turbulence <sup>[17]</sup> was used with modified the scheme
- Fields are evaluated on the expanding grid.

$$k_i = k_0 \cdot \lambda^i \quad (23)$$



Changes that were made:

- 1 Modification of the finite differencing
- 2 The addition for dispersive effects
- 3 Support for strong turbulence
- 4 Implementation of Landau damping



Coupling the grid using interpolated values

[17] S. Galtier et al., Phys. D: Nonl. Phen. **390**, 84 –88 (2019)


# Numerical scheme for NDE - Ask me later?

$$\frac{1}{2} \frac{\partial}{\partial t} \left( \frac{E}{E_C} \right) = \frac{\partial}{\partial k_{\perp}} \left\{ k_{\perp}^6 \begin{pmatrix} V_{ph} \\ 1 \end{pmatrix} \sum_{r=\pm 1} \begin{pmatrix} 1 \\ (-1)^r \end{pmatrix} \frac{E^{(-r)}}{\tilde{k}_{\parallel}^{\pm}} \frac{\partial}{\partial k_{\perp}} \left( \frac{E^{(r)}}{k_{\perp}} \right) \right\} =: \frac{\partial}{\partial k_{\perp}} \left( \frac{\varepsilon}{\eta} \right) \quad (22)$$

- We are modeling coupled nonlinear diffusion equations of **Leith** type.
- Code for Gravitational wave turbulence <sup>[17]</sup> was used with modified the scheme


Fields are evaluated on the expanding grid.

$$k_i = k_0 \cdot \lambda^i \quad (23)$$



**Changes** that were made:

- 1 Modification of the finite differencing
- 2 The addition for dispersive effects
- 3 Support for strong turbulence
- 4 Implementation of Landau damping



Coupling the grid using interpolated values

[17] S. Galtier et al., Phys. D: Nonl. Phen. **390**, 84 –88 (2019)




# Numerical scheme for NDE - Ask me later?

$$\frac{1}{2} \frac{\partial}{\partial t} \left( \frac{E}{E_C} \right) = \frac{\partial}{\partial k_{\perp}} \left\{ k_{\perp}^6 \begin{pmatrix} V_{ph} \\ 1 \end{pmatrix} \sum_{r=\pm 1} \begin{pmatrix} 1 \\ (-1)^r \end{pmatrix} \frac{E^{(-r)}}{\tilde{k}_{\parallel}^{\pm}} \frac{\partial}{\partial k_{\perp}} \left( \frac{E^{(r)}}{k_{\perp}} \right) \right\} =: \frac{\partial}{\partial k_{\perp}} \left( \frac{\varepsilon}{\eta} \right) \quad (22)$$

- We are modeling coupled nonlinear diffusion equations of **Leith** type.
- Code for Gravitational wave turbulence <sup>[17]</sup> was used with modified the scheme


Fields are evaluated on the expanding grid.

$$k_i = k_0 \cdot \lambda^i \quad (23)$$



**Changes** that were made:

- 1 Modification of the finite differencing
- 2 The addition for dispersive effects
- 3 Support for strong turbulence
- 4 Implementation of Landau damping



Coupling the grid using interpolated values

[17] S. Galtier et al., Phys. D: Nonl. Phen. **390**, 84 –88 (2019)

# Value of the correlation coefficient in the dispersive case?

We can rewrite the stationary weak diffusive equations using

$$E^{\pm}(k_{\perp}) =: \frac{1}{2} k_{\perp} \rho(k_{\perp}) e^{\pm \phi(k_{\perp})}. \quad (24)$$

so that  $|E_C(k_{\perp})| \leq E(k_{\perp})/v_{ph}(k_{\perp})$ . NDE rewrites

$$\frac{\partial}{\partial k_{\perp}} \rho^2(k_{\perp}) = -\frac{2\varepsilon}{C k_{\perp}^7 v_{ph}(k_{\perp})} \quad (25)$$

$$\rho^2(k_{\perp}) \frac{\partial}{\partial k_{\perp}} \phi(k_{\perp}) = -\frac{\eta}{C k_{\perp}^7}. \quad (26)$$

Weak turbulence  
analytics

- $v_{ph}(k_{\perp}) \rightarrow k_{\perp} \Rightarrow \rho^2(k_{\perp}) \sim \varepsilon k_{\perp}^{-7} \Rightarrow \phi(k_{\perp}) \sim a + b k_{\perp}$  where  $0 > b \propto \eta/\varepsilon \Rightarrow E^{\pm}(k_{\perp}) \sim k_{\perp}^{-5/2} e^{\pm b k_{\perp}}$

Strong turbulence  
simulation

- So one solutions exponentially diverge, unless we make modifications (confirmed in strong NDE simulation)
- When we choose  $\chi = 1$  this problem is cured.

# Value of the correlation coefficient in the dispersive case?

We can rewrite the stationary weak diffusive equations using

$$E^{\pm}(k_{\perp}) =: \frac{1}{2} k_{\perp} \rho(k_{\perp}) e^{\pm \phi(k_{\perp})}. \quad (24)$$

so that  $|E_C(k_{\perp})| \leq E(k_{\perp})/v_{ph}(k_{\perp})$ . NDE rewrites

$$\frac{\partial}{\partial k_{\perp}} \rho^2(k_{\perp}) = -\frac{2\varepsilon}{C k_{\perp}^7 v_{ph}(k_{\perp})} \quad (25)$$

$$\rho^2(k_{\perp}) \frac{\partial}{\partial k_{\perp}} \phi(k_{\perp}) = -\frac{\eta}{C k_{\perp}^7}. \quad (26)$$

Weak turbulence  
analytics

- $v_{ph}(k_{\perp}) \rightarrow k_{\perp} \Rightarrow \rho^2(k_{\perp}) \sim \varepsilon k_{\perp}^{-7} \Rightarrow \phi(k_{\perp}) \sim a + b k_{\perp}$  where  $0 > b \propto \eta/\varepsilon \Rightarrow E^{\pm}(k_{\perp}) \sim k_{\perp}^{-5/2} e^{\pm b k_{\perp}}$

Strong turbulence  
simulation

- So one solutions exponentially diverge, unless we make modifications (confirmed in strong NDE simulation)
- When we choose  $\chi = 1$  this problem is cured.

# Value of the correlation coefficient in the dispersive case?

We can rewrite the stationary weak diffusive equations using

$$E^{\pm}(k_{\perp}) =: \frac{1}{2} k_{\perp} \rho(k_{\perp}) e^{\pm \phi(k_{\perp})}. \quad (24)$$

so that  $|E_C(k_{\perp})| \leq E(k_{\perp})/v_{ph}(k_{\perp})$ . NDE rewrites

$$\frac{\partial}{\partial k_{\perp}} \rho^2(k_{\perp}) = -\frac{2\varepsilon}{C k_{\perp}^7 v_{ph}(k_{\perp})} \quad (25)$$

$$\rho^2(k_{\perp}) \frac{\partial}{\partial k_{\perp}} \phi(k_{\perp}) = -\frac{\eta}{C k_{\perp}^7}. \quad (26)$$


Weak turbulence  
analytics

- $v_{ph}(k_{\perp}) \rightarrow k_{\perp} \Rightarrow \rho^2(k_{\perp}) \sim \varepsilon k_{\perp}^{-7} \Rightarrow \phi(k_{\perp}) \sim a + b k_{\perp}$  where  $0 > b \propto \eta/\varepsilon \Rightarrow E^{\pm}(k_{\perp}) \sim k_{\perp}^{-5/2} e^{\pm b k_{\perp}}$

Strong turbulence  
simulation

- So one solutions exponentially diverge, unless we make modifications (confirmed in strong NDE simulation)
- When we choose  $\chi = 1$  this problem is cured.

# Value of the correlation coefficient in the dispersive case?



Weak turbulence  
analytics

Strong turbulence  
simulation

We can rewrite the stationary weak diffusive equations using

$$E^\pm(k_\perp) =: \frac{1}{2} k_\perp \rho(k_\perp) e^{\pm \phi(k_\perp)}. \quad (24)$$

so that  $|E_C(k_\perp)| \leq E(k_\perp)/v_{ph}(k_\perp)$ . NDE rewrites

$$\frac{\partial}{\partial k_\perp} \rho^2(k_\perp) = -\frac{2\varepsilon}{C k_\perp^7 v_{ph}(k_\perp)} \quad (25)$$

$$\rho^2(k_\perp) \frac{\partial}{\partial k_\perp} \phi(k_\perp) = -\frac{\eta}{C k_\perp^7}. \quad (26)$$

- $v_{ph}(k_\perp) \rightarrow k_\perp \Rightarrow \rho^2(k_\perp) \sim \varepsilon k_\perp^{-7} \Rightarrow \phi(k_\perp) \sim a + b k_\perp$  where  $0 > b \propto \eta/\varepsilon \Rightarrow E^\pm(k_\perp) \sim k_\perp^{-5/2} e^{\pm b k_\perp}$

- So one solutions exponentially diverge, unless we make modifications (confirmed in strong NDE simulation)

- When we choose  $\chi = 1$  this problem is cured.

# Value of the correlation coefficient in the dispersive case?

We can rewrite the stationary weak diffusive equations using

$$E^\pm(k_\perp) =: \frac{1}{2} k_\perp \rho(k_\perp) e^{\pm \phi(k_\perp)}. \quad (24)$$


so that  $|E_C(k_\perp)| \leq E(k_\perp)/v_{ph}(k_\perp)$ . NDE rewrites

$$\frac{\partial}{\partial k_\perp} \rho^2(k_\perp) = -\frac{2\varepsilon}{C k_\perp^7 v_{ph}(k_\perp)} \quad (25)$$


$$\rho^2(k_\perp) \frac{\partial}{\partial k_\perp} \phi(k_\perp) = -\frac{\eta}{C k_\perp^7}. \quad (26)$$

- $v_{ph}(k_\perp) \rightarrow k_\perp \Rightarrow \rho^2(k_\perp) \sim \varepsilon k_\perp^{-7} \Rightarrow \phi(k_\perp) \sim a + b k_\perp$  where  $0 > b \propto \eta/\varepsilon \Rightarrow E^\pm(k_\perp) \sim k_\perp^{-5/2} e^{\pm b k_\perp}$

- So one solutions exponentially diverge, unless we make modifications (confirmed in strong NDE simulation)
- When we choose  $\chi = 1$  this problem is cured.




Weak turbulence




Strong turbulence simulation

# Presence of dispersion range increases imbalance




Dispersive case,  $k_d = k_d(\nu)$




Imbalance vs. cross-helicity to energy flux ratio.

- The size of the dispersive range is determined by  $k_d$ , which is set by  $\nu$
- This size has significant impact on the size of imbalance:  $E_+/E_-$ !
- It appears that imbalance strongly depends on the cross-helicity flux  $\eta$
- The imbalance is also more pronounced for lower values of  $\beta_e$

# Presence of dispersion range increases imbalance



Dispersive case,  $k_d = k_d(\nu)$




Imbalance vs. cross-helicity to energy flux ratio.


- The size of the dispersive range is determined by  $k_d$ , which is set by  $\nu$
- This size has **significant impact** on the size of **imbalance**:  $E_+/E_-$ !
- It appears that **imbalance** strongly **depends on** the **cross-helicity flux**  $\eta$
- The imbalance is also more pronounced for lower values of  $\beta_e$



# Presence of dispersion range increases imbalance




Dispersive case,  $k_d = k_d(\nu)$




Imbalance vs. cross-helicity to energy flux ratio.

- The size of the dispersive range is determined by  $k_d$ , which is set by  $\nu$
- This size has **significant impact** on the size of **imbalance**:  $E_+/E_-$ !
- It appears that **imbalance** strongly **depends on** the **cross-helicity flux**  $\eta$
- The imbalance is also more pronounced for lower values of  $\beta_e$

# Presence of dispersion range increases imbalance



Dispersive case,  $k_d = k_d(\nu)$



Imbalance vs. cross-helicity to energy flux ratio.

- The size of the dispersive range is determined by  $k_d$ , which is set by  $\nu$
- This size has **significant impact** on the size of **imbalance**:  $E_+/E_-$ !
- It appears that **imbalance** strongly **depends on** the **cross-helicity flux**  $\eta$
- The imbalance is also more pronounced for lower values of  $\beta_e$

# Landau Damping as a turbulence sink

- In solar wind plasmas are collisionless so (hyper)dissipation is unphysical
- We confirm that in case of low beta  $\beta_e = 0.04$  electron contribution is stronger
- Terms  $-2\gamma E$  and  $-2\gamma E_C$  are introduced in spectral equations
- The expression for damping<sup>[18]</sup> is derived from **linearized of gyrokinetics**

$$\gamma = \sqrt{\frac{\pi}{2}} \frac{\delta}{\beta_e} \frac{1}{1 + 4(1 + \tau)^{-2} \beta_e^{-2}} \tilde{k}_{\parallel} k_{\perp}^2, \quad (27)$$

Wave-particle resonance

- Transfer time is also modified however the contribution is negligible.

[18] G. G. Howes et al., ApJ **651**, 590–614 (2006)

# Landau Damping as a turbulence sink

- In solar wind plasmas are collisionless so (hyper)dissipation is unphysical
- We confirm that in case of low beta  $\beta_e = 0.04$  electron contribution is stronger
- Terms  $-2\gamma E$  and  $-2\gamma E_C$  are introduced in spectral equations
- The expression for damping<sup>[18]</sup> is derived from linearized gyrokinetics

$$\gamma = \sqrt{\frac{\pi}{2}} \frac{\delta}{\beta_e} \frac{1}{1 + 4(1 + \tau)^{-2} \beta_e^{-2}} \tilde{k}_{\parallel} k_{\perp}^2, \quad (27)$$


Wave-particle resonance

- Transfer time is also modified however the contribution is negligible.

[18] G. G. Howes et al., ApJ **651**, 590–614 (2006)

# Landau Damping as a turbulence sink

- In solar wind plasmas are collisionless so (hyper)dissipation is unphysical
- We confirm that in case of low beta  $\beta_e = 0.04$  electron contribution is stronger
- Terms  $-2\gamma E$  and  $-2\gamma E_C$  are introduced in spectral equations
- The expression for damping<sup>[18]</sup> is derived from linearized gyrokinetics



$$\gamma = \sqrt{\frac{\pi}{2}} \frac{\delta}{\beta_e} \frac{1}{1 + 4(1 + \tau)^{-2} \beta_e^{-2}} \tilde{k}_{\parallel} k_{\perp}^2, \quad (27)$$


Wave-particle resonance

- Transfer time is also modified however the contribution is negligible.

[18] G. G. Howes et al., ApJ **651**, 590–614 (2006)

# Landau Damping as a turbulence sink

- In solar wind plasmas are collisionless so (hyper)dissipation is unphysical
- We confirm that in case of low beta  $\beta_e = 0.04$  electron contribution is stronger
- Terms  $-2\gamma E$  and  $-2\gamma E_C$  are introduced in spectral equations
- The expression for damping<sup>[18]</sup> is derived from **linearized of gyrokinetics**



$$\gamma = \sqrt{\frac{\pi}{2}} \frac{\delta}{\beta_e} \frac{1}{1 + 4(1 + \tau)^{-2} \beta_e^{-2}} \tilde{k}_{\parallel} k_{\perp}^2, \quad (27)$$


Wave-particle resonance

- Transfer time is also modified however the contribution is negligible.

[18] G. G. Howes et al., ApJ **651**, 590–614 (2006)

# Landau Damping as a turbulence sink

- In solar wind plasmas are collisionless so (hyper)dissipation is unphysical
- We confirm that in case of low beta  $\beta_e = 0.04$  electron contribution is stronger
- Terms  $-2\gamma E$  and  $-2\gamma E_C$  are introduced in spectral equations
- The expression for damping<sup>[18]</sup> is derived from **linearized of gyrokinetics**




$$\gamma = \sqrt{\frac{\pi}{2}} \frac{\delta}{\beta_e} \frac{1}{1 + 4(1 + \tau)^{-2} \beta_e^{-2}} \tilde{k}_{\parallel} k_{\perp}^2, \quad (27)$$

Wave-particle resonance


- Transfer time is also modified however the contribution is negligible.

[18] G. G. Howes et al., ApJ **651**, 590–614 (2006)

# Landau damping effects



Different models of dissipation




Steepening of the energy spectra vs  $\eta$ .

## Results


- Correct mechanism of dissipation has a profound effect on the dispersion range
- When  $\chi < 1$  we see the possibility of having a knee in the spectrum
- Non-universality: for larger values  $\eta$  injection the spectrum becomes steeper.



# Landau damping effects



Different models of dissipation




Steepening of the energy spectra vs  $\eta$ .


## Results

- Correct mechanism of dissipation has a profound effect on the dispersion range
- When  $\chi < 1$  we see the possibility of having a knee in the spectrum
- Non-universality: for larger values  $\eta$  injection the spectrum becomes steeper.

# Landau damping effects



Different models of dissipation




Steepening of the energy spectra vs  $\eta$ .


## Results

- Correct mechanism of dissipation has a profound effect on the dispersion range
- When  $\chi < 1$  we see the possibility of having a knee in the spectrum
- Non-universality: for larger values  $\eta$  injection the spectrum becomes steeper.

# Inverse cascade of cross-helicity and forward of energy



Dispersive case: Fluxes for  $k_f \sim 0.016$




Small scale to large scale cross-helicity injection ratio needed to drive the inverse cascade


- In presence of reservoir  $E_{\pm}$  it is possible to drive  $\eta_-$  from small scales
- Inverse cascade exists even in MHD but is stronger when dispersive range is present
- In sub-ion range  $E_C$  approaches magnetic-helicity so is perhaps related to Ref. [19]

[19] U. Frisch et al., J. of Fluid Mech. **68**, 769–778 (1975)

# Inverse cascade of cross-helicity and forward of energy



Dispersive case: Fluxes for  $k_f \sim 0.016$



Small scale to large scale cross-helicity injection ratio needed to drive the inverse cascade

- In presence of reservoir  $E_{\pm}$  it is possible to drive  $\eta_-$  from small scales
- Inverse cascade exists even in MHD but is stronger when dispersive range is present
- In sub-ion range  $E_C$  approaches magnetic-helicity so is perhaps related to Ref. [19]

[19] U. Frisch et al., J. of Fluid Mech. **68**, 769–778 (1975)

- 1 Introduction
  - MHD Turbulence
  - Hamiltonian gyrofluid model
- 2 Influence of the dispersive range
  - Nonlinear diffusion equation
  - Landau damping
  - Inverse Cascade
- 3 Conclusion
  - Comparisons with 3D gyrofluid simulations
  - Future Work

# Comparison with direct 3D gyrofluid simulations

- We have analysed consequences of NDEs that describe imbalanced KAW turbulence
- We conclude that **dispersive range** significantly affects the **imbalance**
- The dissipation mechanism is important, e.g. Landau damping

Preliminary results from 3D DNS are supporting some of the claims

- In particular the amount of imbalance strongly depends on the size of  $k_{\perp}$
- ToDo: Measure correlation lengths  $\tilde{\lambda}_{\parallel}^{\pm}$  and estimate  $\chi$

# Comparison with direct 3D gyrofluid simulations

- We have analysed consequences of NDEs that describe imbalanced KAW turbulence
- We conclude that **dispersive range** significantly affects the **imbalance**
- The dissipation mechanism is important, e.g. Landau damping

Preliminary results from 3D DNS are supporting some of the claims

- In particular the amount of imbalance strongly depends on the size of  $k_{\perp}$
- ToDo: Measure correlation lengths  $\tilde{\lambda}_{\parallel}^{\pm}$  and estimate  $\chi$

# Comparison with direct 3D gyrofluid simulations

- We have analysed consequences of NDEs that describe imbalanced KAW turbulence
- We conclude that **dispersive range** significantly affects the **imbalance**
- The dissipation mechanism is important, e.g. Landau damping

Preliminary results from 3D DNS are supporting some of the claims

- In particular the amount of imbalance strongly depends on the size of  $k_{\perp}$
- ToDo: Measure correlation lengths  $\tilde{\lambda}_{\parallel}^{\pm}$  and estimate  $\chi$




# Comparison with direct 3D gyrofluid simulations


- We have analysed consequences of NDEs that describe imbalanced KAW turbulence
- We conclude that **dispersive range** significantly affects the **imbalance**
- The dissipation mechanism is important, e.g. Landau damping

Preliminary results from 3D DNS are supporting some of the claims


- In particular the amount of imbalance strongly depends on the size of  $k_d$
- ToDo: Measure correlation lengths  $\tilde{k}_{\parallel}^{\pm}$  and estimate  $\chi$



Turbulent current  $j_z$



Spectra  $E_{\pm}$  for different box sizes




Growth of imbalance  $E_+/E_-$  vs. time

# Comparison with direct 3D gyrofluid simulations


- We have analysed consequences of NDEs that describe imbalanced KAW turbulence
- We conclude that **dispersive range** significantly affects the **imbalance**
- The dissipation mechanism is important, e.g. Landau damping

Preliminary results from 3D DNS are supporting some of the claims


- In particular the amount of imbalance strongly depends on the size of  $k_d$
- ToDo: Measure correlation lengths  $\tilde{k}_{||}^{\pm}$  and estimate  $\chi$



Turbulent current  $j_z$



Spectra  $E_{\pm}$  for different box sizes




Growth of imbalance  $E_+/E_-$  vs. time

# Comparison with direct 3D gyrofluid simulations


- We have analysed consequences of NDEs that describe imbalanced KAW turbulence
- We conclude that **dispersive range** significantly affects the **imbalance**
- The dissipation mechanism is important, e.g. Landau damping

Preliminary results from 3D DNS are supporting some of the claims


- In particular the amount of imbalance strongly depends on the size of  $k_d$
- ToDo: Measure correlation lengths  $\tilde{k}_{||}^{\pm}$  and estimate  $\chi$



Turbulent current  $j_z$



Spectra  $E_{\pm}$  for different box sizes



Growth of imbalance  $E_+/E_-$  vs. time