

## Two spins

The spin state  $|s\rangle$  for two spins is a unit vector in  $\mathbb{C}^4$ .

$$|s\rangle = \begin{pmatrix} c_{++} \\ c_{+-} \\ c_{-+} \\ c_{--} \end{pmatrix}, \quad |c_{++}|^2 + |c_{+-}|^2 + |c_{-+}|^2 + |c_{--}|^2 = 1$$

Spin measurement probabilities are the transition probabilities from  $|s\rangle$  to an eigenstate.

For spin measurements in the  $z$  direction we have

$$\begin{aligned} \Pr(S_{1z} = +\frac{\hbar}{2} \text{ and } S_{2z} = +\frac{\hbar}{2}) &= |\langle z_{++}|s\rangle|^2 = |c_{++}|^2 \\ \Pr(S_{1z} = +\frac{\hbar}{2} \text{ and } S_{2z} = -\frac{\hbar}{2}) &= |\langle z_{+-}|s\rangle|^2 = |c_{+-}|^2 \\ \Pr(S_{1z} = -\frac{\hbar}{2} \text{ and } S_{2z} = +\frac{\hbar}{2}) &= |\langle z_{-+}|s\rangle|^2 = |c_{-+}|^2 \\ \Pr(S_{1z} = -\frac{\hbar}{2} \text{ and } S_{2z} = -\frac{\hbar}{2}) &= |\langle z_{--}|s\rangle|^2 = |c_{--}|^2 \end{aligned}$$

where the eigenstates are

$$z_{++} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad z_{+-} = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \quad z_{-+} = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}, \quad z_{--} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}$$

Spin operators and expectation values for the first spin ( $\otimes$  is kronecker product).

$$\begin{aligned} S_{1x} &= \frac{\hbar}{2}\sigma_x \otimes I = \frac{\hbar}{2} \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, & \langle S_{1x} \rangle &= \langle s|S_{1x}|s \rangle \\ S_{1y} &= \frac{\hbar}{2}\sigma_y \otimes I = \frac{\hbar}{2} \begin{pmatrix} 0 & 0 & -i & 0 \\ 0 & 0 & 0 & -i \\ i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{pmatrix}, & \langle S_{1y} \rangle &= \langle s|S_{1y}|s \rangle \\ S_{1z} &= \frac{\hbar}{2}\sigma_z \otimes I = \frac{\hbar}{2} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, & \langle S_{1z} \rangle &= \langle s|S_{1z}|s \rangle \end{aligned}$$

Spin operators and expectation values for the second spin.

$$\begin{aligned}
S_{2x} &= \frac{\hbar}{2} I \otimes \sigma_x = \frac{\hbar}{2} \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, & \langle S_{2x} \rangle &= \langle s | S_{2x} | s \rangle \\
S_{2y} &= \frac{\hbar}{2} I \otimes \sigma_y = \frac{\hbar}{2} \begin{pmatrix} 0 & -i & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & -i \\ 0 & 0 & i & 0 \end{pmatrix}, & \langle S_{2y} \rangle &= \langle s | S_{2y} | s \rangle \\
S_{2z} &= \frac{\hbar}{2} I \otimes \sigma_z = \frac{\hbar}{2} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, & \langle S_{2z} \rangle &= \langle s | S_{2z} | s \rangle
\end{aligned}$$

By definition of expectation value we have for the  $z$  direction

$$\begin{aligned}
\langle S_{1z} \rangle &= \frac{\hbar}{2} (|c_{++}|^2 + |c_{+-}|^2 - |c_{-+}|^2 - |c_{--}|^2) \\
\langle S_{2z} \rangle &= \frac{\hbar}{2} (|c_{++}|^2 - |c_{+-}|^2 + |c_{-+}|^2 - |c_{--}|^2)
\end{aligned}$$

1. Verify spin operators for two spins.
---

```
sigmax = ((0,1),(1,0))
sigmay = ((0,-i),(i,0))
sigmaz = ((1,0),(0,-1))

I = ((1,0),(0,1))

S1x = 1/2 hbar kronecker(sigmax,I)
S1y = 1/2 hbar kronecker(sigmay,I)
S1z = 1/2 hbar kronecker(sigmaz,I)

S2x = 1/2 hbar kronecker(I,sigmax)
S2y = 1/2 hbar kronecker(I,sigmay)
S2z = 1/2 hbar kronecker(I,sigmaz)

check(S1x == 1/2 hbar ((0,0,1,0),(0,0,0,1),(1,0,0,0),(0,1,0,0)))
check(S1y == 1/2 hbar ((0,0,-i,0),(0,0,0,-i),(i,0,0,0),(0,i,0,0)))
check(S1z == 1/2 hbar ((1,0,0,0),(0,1,0,0),(0,0,-1,0),(0,0,0,-1)))

check(S2x == 1/2 hbar ((0,1,0,0),(1,0,0,0),(0,0,0,1),(0,0,1,0)))
check(S2y == 1/2 hbar ((0,-i,0,0),(i,0,0,0),(0,0,0,-i),(0,0,i,0)))
check(S2z == 1/2 hbar ((1,0,0,0),(0,-1,0,0),(0,0,1,0),(0,0,0,-1)))
```

2. Verify expectation values for the $z$ direction.
---

```
sigmax = ((0,1),(1,0))
sigmay = ((0,-i),(i,0))
sigmaz = ((1,0),(0,-1))

I = ((1,0),(0,1))

S1x = 1/2 hbar kronecker(sigmax,I)
S1y = 1/2 hbar kronecker(sigmay,I)
S1z = 1/2 hbar kronecker(sigmaz,I)

S2x = 1/2 hbar kronecker(I,sigmax)
S2y = 1/2 hbar kronecker(I,sigmay)
S2z = 1/2 hbar kronecker(I,sigmaz)

c1 = x1 + i y1
c2 = x2 + i y2
c3 = x3 + i y3
c4 = x4 + i y4

s = (c1,c2,c3,c4)

check(dot(conj(s),S1z,s) ==
1/2 hbar (conj(c1) c1 + conj(c2) c2 - conj(c3) c3 - conj(c4) c4))

check(dot(conj(s),S2z,s) ==
1/2 hbar (conj(c1) c1 - conj(c2) c2 + conj(c3) c3 - conj(c4) c4))
```