

Feynman and Hibbs problem 5-8

Note that equation (5.44) implies  $G_A^*(x, x') = G_A(x', x)$ . With this in mind show that for any two wave functions  $g(x)$  and  $f(x)$ , both of which approach 0 as  $x$  goes to  $\pm\infty$ ,

$$\int_{-\infty}^{\infty} g^*(x) \mathcal{A}f(x) dx = \int_{-\infty}^{\infty} (\mathcal{A}g(x))^* f(x) dx \quad (5.47)$$

Any operator, such as  $\mathcal{A}$ , for which equation (5.47) holds is called “hermitian” (see equation 4.30).

From equations (5.45) and (5.43)

$$\mathcal{A}f(x) = \int_{-\infty}^{\infty} G_A(x, x') f(x') dx' \quad (1)$$

Hence

$$\int_{-\infty}^{\infty} g^*(x) \mathcal{A}f(x) dx = \int_{-\infty}^{\infty} g^*(x) \left( \int_{-\infty}^{\infty} G_A(x, x') f(x') dx' \right) dx$$

Combine factors.

$$= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} G_A(x, x') g^*(x) f(x') dx' dx$$

By equation (5.44) substitute  $G_A^*(x', x)$  for  $G_A(x, x')$ .

$$= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} G_A^*(x', x) g^*(x) f(x') dx' dx$$

By complex conjugation of a product

$$= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (G_A(x', x) g(x))^* f(x') dx' dx$$

Factor out  $f(x') dx'$ .

$$= \int_{-\infty}^{\infty} \left( \int_{-\infty}^{\infty} (G_A(x', x) g(x))^* dx \right) f(x') dx'$$

Then by equation (1)

$$= \int_{-\infty}^{\infty} (\mathcal{A}g(x'))^* f(x') dx'$$