

(36.1) *An illustration of the reason for anticommutation and spin*

(a) Show that the Dirac equation can be recast in the form

$$i\frac{\partial\psi}{\partial t} = \hat{H}_D\psi \quad (36.33)$$

where  $\hat{H}_D = \boldsymbol{\alpha} \cdot \hat{\mathbf{p}} + \beta m$  and find  $\boldsymbol{\alpha}$  and  $\beta$  in terms of the  $\gamma$  matrices.

(b) Evaluate  $\hat{H}_D^2$  and show that for a Klein-Gordon dispersion to result we must have:

(i) that the  $\alpha^i$  and  $\beta$  objects all anticommute with each other; and

(ii)  $(\alpha^i)^2 = (\beta)^2 = 1$ .

*This provides some justification for the anticommutation relations we imposed on the  $\gamma$ s.*

---

(a) Consider the following form of the Dirac equation.

$$i\left(\gamma^0\frac{\partial}{\partial t} + \gamma^1\frac{\partial}{\partial x} + \gamma^2\frac{\partial}{\partial y} + \gamma^3\frac{\partial}{\partial z}\right)\psi = m\psi$$

Rewrite as

$$i\gamma^0\frac{\partial}{\partial t}\psi = -i\left(\gamma^1\frac{\partial}{\partial x} + \gamma^2\frac{\partial}{\partial y} + \gamma^3\frac{\partial}{\partial z}\right)\psi + m\psi$$

Noting that  $\gamma^0\gamma^0 = I$ , multiply both sides by  $\gamma^0$  to obtain

$$i\frac{\partial}{\partial t}\psi = -i\gamma^0\left(\gamma^1\frac{\partial}{\partial x} + \gamma^2\frac{\partial}{\partial y} + \gamma^3\frac{\partial}{\partial z}\right)\psi + m\gamma^0\psi$$

Hence for  $\hat{\mathbf{p}} = -i\nabla$  we have

$$\boldsymbol{\alpha} = \gamma^0 \begin{pmatrix} \gamma^1 \\ \gamma^2 \\ \gamma^3 \end{pmatrix}, \quad \beta = \gamma^0$$