

## Free-particle propagator

The propagator  $K(x_b, t_b, x_a, t_a)$  is the amplitude for a particle to go from  $x_a$  at time  $t_a$  to  $x_b$  at time  $t_b$ .

$$K(x_b, t_b, x_a, t_a) = \langle x_b | \exp \left( -\frac{i\hat{H}(t_b - t_a)}{\hbar} \right) | x_a \rangle$$

Let  $\hat{H}$  be the free-particle Hamiltonian

$$\hat{H} = \frac{\hat{p}^2}{2m}$$

Then

$$K(x_b, t_b, x_a, t_a) = \langle x_b | \exp \left( -\frac{i\hat{p}^2(t_b - t_a)}{2m\hbar} \right) | x_a \rangle$$

By the identity

$$\int_{-\infty}^{\infty} |p\rangle \langle p| dp = 1$$

we can write

$$K(x_b, t_b, x_a, t_a) = \int_{-\infty}^{\infty} \langle x_b | \exp \left( -\frac{i\hat{p}^2(t_b - t_a)}{2m\hbar} \right) | p \rangle \langle p | x_a \rangle dp$$

By the identity  $\hat{p}|p\rangle = p|p\rangle$  replace operator  $\hat{p}$  with its eigenvalue  $p$ .

$$K(x_b, t_b, x_a, t_a) = \int \langle x_b | \exp \left( -\frac{ip^2(t_b - t_a)}{2m\hbar} \right) | p \rangle \langle p | x_a \rangle dp$$

Rearrange as

$$K(x_b, t_b, x_a, t_a) = \int_{-\infty}^{\infty} \exp \left( -\frac{ip^2(t_b - t_a)}{2m\hbar} \right) \langle x_b | p \rangle \langle p | x_a \rangle dp$$

For a free particle

$$\langle x | p \rangle = \frac{1}{\sqrt{2\pi\hbar}} \exp \left( \frac{ipx}{\hbar} \right)$$

Hence

$$K(x_b, t_b, x_a, t_a) = \int_{-\infty}^{\infty} \exp \left( -\frac{ip^2(t_b - t_a)}{2m\hbar} \right) \frac{1}{\sqrt{2\pi\hbar}} \exp \left( \frac{ipx_b}{\hbar} \right) \frac{1}{\sqrt{2\pi\hbar}} \exp \left( -\frac{ipx_a}{\hbar} \right) dp$$

Simplify as

$$K(x_b, t_b, x_a, t_a) = \frac{1}{2m\hbar} \int_{-\infty}^{\infty} \exp \left( -\frac{ip^2(t_b - t_a)}{2m\hbar} + \frac{ip(x_b - x_a)}{\hbar} \right) dp$$

Let

$$a = \frac{i(t_b - t_a)}{2m\hbar}, \quad b = \frac{i(x_b - x_a)}{\hbar}$$

Then by the integral

$$\int_{-\infty}^{\infty} \exp(-ay^2 + by) \, dy = \left(\frac{\pi}{a}\right)^{1/2} \exp\left(\frac{b^2}{4a}\right)$$

we have

$$K(x_b, t_b, x_a, t_a) = \frac{1}{2\pi\hbar} \left(\frac{2\pi m\hbar}{i(t_b - t_a)}\right)^{1/2} \exp\left(-\frac{m(x_b - x_a)^2}{2i\hbar(t_b - t_a)}\right)$$

Rearrange as

$$K(x_b, t_b, x_a, t_a) = \left(\frac{m}{2\pi i\hbar(t_b - t_a)}\right)^{1/2} \exp\left(\frac{im(x_b - x_a)^2}{2\hbar(t_b - t_a)}\right)$$

[Click here to verify.](#)