

Feynman and Hibbs problem 4-12

Carry out the integral in equation (4.64) by completing the square. Show that the correct free particle kernel (i.e., the three dimensional version of equation 3.3) results.

$$K_0(\mathbf{x}_b, t_b, \mathbf{x}_a, t_a) = \int \exp\left(\frac{i}{\hbar} \mathbf{p} \cdot (\mathbf{x}_b - \mathbf{x}_a)\right) \exp\left(-\frac{i}{\hbar} \frac{\mathbf{p}^2}{2m} (t_b - t_a)\right) \frac{d^3 \mathbf{p}}{(2\pi\hbar)^3} \quad (4.64)$$

Rewrite equation (4.64) in component notation.

$$K_0(\mathbf{x}_b, t_b, \mathbf{x}_a, t_a) = \frac{1}{(2\pi\hbar)^3} \int_{\mathbb{R}^3} \exp\left(\frac{i}{\hbar} p_1 X_1 - \frac{i}{\hbar} \frac{p_1^2}{2m} T\right) \times \exp\left(\frac{i}{\hbar} p_2 X_2 - \frac{i}{\hbar} \frac{p_2^2}{2m} T\right) \exp\left(\frac{i}{\hbar} p_3 X_3 - \frac{i}{\hbar} \frac{p_3^2}{2m} T\right) dp_1 dp_2 dp_3$$

where

$$X_k = (\mathbf{x}_b - \mathbf{x}_a)_k \\ T = t_b - t_a$$

Let

$$A = -\frac{i}{2m\hbar} T \\ B = \frac{i}{\hbar} X_k$$

Then

$$\int_{-\infty}^{\infty} \exp(Ap^2 + Bp) dp = \left(-\frac{\pi}{A}\right)^{1/2} \exp\left(-\frac{B^2}{4A}\right) \\ = \left(-\frac{2\pi im\hbar}{T}\right)^{1/2} \exp\left(\frac{imX_k^2}{2\hbar T}\right) \quad (1)$$

Hence

$$K_0(\mathbf{x}_b, t_b, \mathbf{x}_a, t_a) = \frac{1}{(2\pi\hbar)^3} \prod_{k=1}^3 \left(-\frac{2\pi im\hbar}{T}\right)^{1/2} \exp\left(\frac{imX_k^2}{2\hbar T}\right)$$

Combine coefficients and convert to vector notation.

$$K_0(\mathbf{x}_b, t_b, \mathbf{x}_a, t_a) = \left(-\frac{im}{2\pi\hbar(t_b - t_a)}\right)^{3/2} \exp\left(\frac{im(\mathbf{x}_b - \mathbf{x}_a)^2}{2\hbar(t_b - t_a)}\right) \quad (2)$$