



Rutherford Scattering

Consider an electron scattered by an atomic nucleus.



Here is the same diagram with momentum and spinor labels.



The incident electron can be modeled as coincident with the z axis, resulting in the following momentum vectors.

$$p_1 = \begin{pmatrix} E \\ 0 \\ 0 \\ p \end{pmatrix} \quad p_2 = \begin{pmatrix} E \\ p \sin \theta \cos \phi \\ p \sin \theta \sin \phi \\ p \cos \theta \end{pmatrix}$$

inbound electron outbound electron

Symbol $E = \sqrt{p^2 + m^2}$ is total energy, p is electron momentum, and m is electron mass.

The spinors are

$$u_{11} = \begin{pmatrix} E + m \\ 0 \\ p \\ 0 \end{pmatrix} \quad u_{12} = \begin{pmatrix} 0 \\ E + m \\ 0 \\ -p \end{pmatrix}$$

inbound electron, spin up inbound electron, spin down

$$u_{21} = \begin{pmatrix} E + m \\ 0 \\ p_{2z} \\ p_{2x} + ip_{2y} \end{pmatrix} \quad u_{22} = \begin{pmatrix} 0 \\ E + m \\ p_{2x} - ip_{2y} \\ -p_{2z} \end{pmatrix}$$

outbound electron, spin up outbound electron, spin down

The second digit in each spinor subscript indicates whether the spin state is up or down. It should be noted that the spinors shown above are not individually normalized. Instead, a combined spinor normalization constant $N = (E + m)^2$ is used.

The following formula computes a probability density $|\mathcal{M}_{jk}|^2$ for Rutherford scattering where j is the spin state of the inbound electron and k is the spin state of the outbound electron.

$$|\mathcal{M}_{jk}|^2 = \frac{Z^2 e^4}{q^4} \frac{1}{N} |\bar{u}_{2k} \gamma^0 u_{1j}|^2$$

Symbol Z is the atomic number of the nucleus, e is electron charge, and $q = p_1 - p_2$ is momentum transfer.

The expected probability density $\langle |\mathcal{M}|^2 \rangle$ is computed by summing $|\mathcal{M}_{jk}|^2$ over all four spin states and then dividing by the number of inbound states. There are two inbound states.

$$\begin{aligned} \langle |\mathcal{M}|^2 \rangle &= \frac{1}{2} \sum_{j=1}^2 \sum_{k=1}^2 |\mathcal{M}_{jk}|^2 \\ &= \frac{Z^2 e^4}{2q^4} \frac{1}{N} \sum_{j=1}^2 \sum_{k=1}^2 |\bar{u}_{2k} \gamma^0 u_{1j}|^2 \\ &= \frac{Z^2 e^4}{2q^4} \text{Tr} \left((\not{p}_1 + m) \gamma^0 (\not{p}_2 + m) \gamma^0 \right) \\ &= \frac{2Z^2 e^4}{q^4} (E^2 + m^2 + p^2 \cos \theta) \end{aligned}$$

Run “rutherford-scattering-1.txt” to verify the following formulas.

$$\begin{aligned} \frac{1}{N} \sum_{j=1}^2 \sum_{k=1}^2 |\bar{u}_{2k} \gamma^0 u_{1j}|^2 &= \text{Tr} \left((\not{p}_1 + m) \gamma^0 (\not{p}_2 + m) \gamma^0 \right) = 4(E^2 + m^2 + p^2 \cos \theta) \\ q^4 &= (p_1 - p_2)^4 = 16p^4 \sin^4(\theta/2) = 4p^4(\cos \theta - 1)^2 \end{aligned}$$

Low energy approximation

For low energy electrons such that $p \ll m$ we can use the following approximation.

$$E^2 + m^2 + p^2 \cos \theta \approx 2m^2$$

Hence

$$\langle |\mathcal{M}|^2 \rangle = \frac{4m^2 Z^2 e^4}{q^4}$$

Substituting $e^2 = 4\pi\alpha$ and $q^4 = 4p^4(\cos \theta - 1)^2$ we have

$$\langle |\mathcal{M}|^2 \rangle = \frac{16\pi^2 m^2 Z^2 \alpha^2}{p^4 (\cos \theta - 1)^2}$$

The differential cross section is

$$\frac{d\sigma}{d\Omega} = \frac{\langle |\mathcal{M}|^2 \rangle}{16\pi^2} = \frac{m^2 Z^2 \alpha^2}{p^4 (\cos \theta - 1)^2}$$

We can integrate $d\sigma$ to obtain a cumulative distribution function. Recall that $d\Omega = \sin \theta \, d\theta \, d\phi$, hence

$$d\sigma = \frac{m^2 Z^2 \alpha^2}{p^4 (\cos \theta - 1)^2} \sin \theta \, d\theta \, d\phi$$

Let $I(\xi)$ be the following definite integral.

$$\begin{aligned} I(\xi) &= \frac{p^4}{m^2 Z^2 \alpha^2} \int_0^{2\pi} \int_a^\xi d\sigma \\ &= 2\pi \int_a^\xi \frac{1}{(\cos \theta - 1)^2} \sin \theta \, d\theta \\ &= 2\pi \left(\frac{1}{\cos \theta - 1} \right) \Big|_a^\xi \\ &= 2\pi \left(\frac{1}{\cos \xi - 1} - \frac{1}{\cos a - 1} \right), \quad a \leq \xi \leq \pi \end{aligned}$$

A lower bound of $a > 0$ is required because $I(0)$ is undefined.

Let C be the normalization constant $C = I(\pi)$. Then the cumulative distribution function $F(\theta)$ is

$$F(\theta) = C^{-1} I(\theta), \quad a \leq \theta \leq \pi$$


The probability of observing scattering events in the interval θ_1 to θ_2 can now be computed.

$$P(\theta_1 \leq \theta \leq \theta_2) = F(\theta_2) - F(\theta_1)$$

Probability density function $f(\theta)$ is the derivative of $F(\theta)$.

$$f(\theta) = \frac{dF(\theta)}{d\theta} = C^{-1} \frac{dI(\theta)}{d\theta} = C^{-1} \frac{2\pi \sin \theta}{(\cos \theta - 1)^2}$$

Run “rutherford-scattering-3.txt” to draw a graph of $f(\theta)$ for $a = \pi/4 = 45^\circ$.



The following table shows the corresponding probability distribution for three bins.

θ_1	θ_2	$P(\theta_1 \leq \theta \leq \theta_2)$
0°	45°	—
45°	90°	0.83
90°	135°	0.14
135°	180°	0.03