

Exercise 6.6. Assume Charlie has prepared the two spins in the singlet state. This time, Bob measures  $\tau_y$  and Alice measures  $\sigma_x$ . What is the expectation value of  $\sigma_x \tau_y$ ?

What does this say about the correlation between the two measurements?

---

Recall that

$$|sing\rangle = \frac{|ud\rangle - |du\rangle}{\sqrt{2}}$$

The expectation value of  $\sigma_x \tau_y$  is

$$\langle \sigma_x \tau_y \rangle = \langle sing | \sigma_x \tau_y | sing \rangle$$

Applying the  $\tau_y$  operator first we have from Table 1 the following result.

$$\langle \sigma_x \tau_y \rangle = \langle sing | \sigma_x \left( \frac{-i|uu\rangle - i|dd\rangle}{\sqrt{2}} \right)$$

Then applying the  $\sigma_x$  operator we have

$$\langle \sigma_x \tau_y \rangle = \langle sing | \left( \frac{-i|du\rangle - i|ud\rangle}{\sqrt{2}} \right)$$

Hence

$$\begin{aligned} \langle \sigma_x \tau_y \rangle &= \frac{1}{2} (\langle ud | - \langle du |) (-i|du\rangle - i|ud\rangle) \\ &= \frac{1}{2} (-i\langle ud | ud \rangle + i\langle du | du \rangle) \\ &= 0 \end{aligned} \tag{1}$$

From (1) and the result  $\langle \sigma_x \rangle = \langle \tau_y \rangle = 0$  given on page 174 we have

$$\langle \sigma_x \tau_y \rangle - \langle \sigma_x \rangle \langle \tau_y \rangle = 0$$

Hence the measurements  $\sigma_x$  and  $\tau_y$  are uncorrelated.