

15.7.1. Consider a set of modes of the electromagnetic field in which the electric field is polarized along the  $x$  direction and the magnetic field is polarized in the  $y$  direction. Restricting consideration only to those modes, find the simplest expression you can for the commutation relation  $[\hat{E}_x, \hat{B}_y]$  for this multimode field.

---

Consider the following formulas.

$$\hat{\mathbf{E}}(\mathbf{r}, t) = i \sum_{\lambda} \left( \hat{a}_{\lambda} - \hat{a}_{\lambda}^{\dagger} \right) \sqrt{\frac{\hbar \omega_{\lambda}}{2 \epsilon_0}} \mathbf{u}_{\lambda}(\mathbf{r}) \quad (15.132)$$

$$\hat{\mathbf{B}}(\mathbf{r}, t) = \sum_{\lambda} \left( \hat{a}_{\lambda} + \hat{a}_{\lambda}^{\dagger} \right) \sqrt{\frac{\hbar \omega_{\lambda} \mu_0}{2}} \mathbf{v}_{\lambda}(\mathbf{r}) \quad (15.135)$$

By hypothesis for the polarization we have

$$\mathbf{u}_{\lambda}(\mathbf{r}) = \begin{pmatrix} u_{x\lambda}(x, y, \xi) \\ 0 \\ 0 \end{pmatrix}, \quad \mathbf{v}_{\lambda}(\mathbf{r}) = \begin{pmatrix} 0 \\ v_{y\lambda}(x, y, \xi) \\ 0 \end{pmatrix}$$

Hence

$$\begin{aligned} \hat{E}_x(\mathbf{r}, t) &= i \sum_{\lambda} \left( \hat{a}_{\lambda} - \hat{a}_{\lambda}^{\dagger} \right) \sqrt{\frac{\hbar \omega_{\lambda}}{2 \epsilon_0}} u_{x\lambda} \\ \hat{B}_y(\mathbf{r}, t) &= \sum_{\lambda} \left( \hat{a}_{\lambda} + \hat{a}_{\lambda}^{\dagger} \right) \sqrt{\frac{\hbar \omega_{\lambda} \mu_0}{2}} v_{y\lambda} \end{aligned}$$

Using Eigenmath we obtain

$$[\hat{E}_x, \hat{B}_y] = i \hbar \sqrt{\frac{\mu_0}{\epsilon_0}} \sum_{\lambda} \omega_{\lambda} \left( u_{x\lambda} v_{y\lambda} + \xi u_{x\lambda} \frac{d}{d\xi} v_{y\lambda} \right)$$