Spin operator table

Verify the following spin operator table.¹

Table elements are final states. For example,

$$\sigma_z |dd\rangle = -|dd\rangle$$

For single spins we have

$$|u\rangle = \begin{pmatrix} 1\\0 \end{pmatrix}, \quad |d\rangle = \begin{pmatrix} 0\\1 \end{pmatrix}$$

For a system of two spins we use the Kronecker product of single spins.

$$|uu\rangle = |u\rangle \otimes |u\rangle$$
$$|ud\rangle = |u\rangle \otimes |d\rangle$$
$$|du\rangle = |d\rangle \otimes |u\rangle$$
$$|dd\rangle = |d\rangle \otimes |d\rangle$$

Hence

$$|uu\rangle = \begin{pmatrix} 1\\0\\0\\0 \end{pmatrix}, \quad |ud\rangle = \begin{pmatrix} 0\\1\\0\\0 \end{pmatrix}, \quad |du\rangle = \begin{pmatrix} 0\\0\\1\\0 \end{pmatrix}, \quad |dd\rangle = \begin{pmatrix} 0\\0\\0\\1 \end{pmatrix}$$

The spin operators for single spins are

$$\sigma_z = \tau_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \sigma_x = \tau_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \tau_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$$

For a system of two spins, the σ operators operate on the first spin and the τ operators operate on the second spin.

$$\sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \otimes I = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

$$\tau_z = I \otimes \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

¹See Quantum Mechanics by Susskind and Friedman, p. 350.