


# Klein-Nishina formula

The Klein-Nishina formula is the differential cross section for photon-electron scattering.



It is easy to derive the Klein-Nishina formula from Dirac's equation by starting out in the center-of-mass frame and then boosting to the lab frame. In the center-of-mass frame we have the following momentum vectors where  $E = \sqrt{\omega^2 + m^2}$ .

$$\begin{aligned}
 p_1 &= \begin{pmatrix} \omega \\ 0 \\ 0 \\ \omega \end{pmatrix} & p_2 &= \begin{pmatrix} E \\ 0 \\ 0 \\ -\omega \end{pmatrix} & p_3 &= \begin{pmatrix} \omega \\ \omega \sin \theta \cos \phi \\ \omega \sin \theta \sin \phi \\ \omega \cos \theta \end{pmatrix} & p_4 &= \begin{pmatrix} E \\ -\omega \sin \theta \cos \phi \\ -\omega \sin \theta \sin \phi \\ -\omega \cos \theta \end{pmatrix} \\
 &\text{inbound photon} & &\text{inbound electron} & &\text{outbound photon} & &\text{outbound electron}
 \end{aligned}$$

Spinors for the inbound electron.

$$\begin{aligned}
 u_{21} &= \frac{1}{\sqrt{E+m}} \begin{pmatrix} E+m \\ 0 \\ -\omega \\ 0 \end{pmatrix} & u_{22} &= \frac{1}{\sqrt{E+m}} \begin{pmatrix} 0 \\ E+m \\ 0 \\ \omega \end{pmatrix} \\
 &\text{inbound electron spin up} & &\text{inbound electron spin down}
 \end{aligned}$$

Spinors for the outbound electron.

$$\begin{aligned}
 u_{41} &= \frac{1}{\sqrt{E+m}} \begin{pmatrix} E+m \\ 0 \\ p_{4z} \\ p_{4x} + ip_{4y} \end{pmatrix} & u_{42} &= \frac{1}{\sqrt{E+m}} \begin{pmatrix} 0 \\ E+m \\ p_{4x} - ip_{4y} \\ -p_{4z} \end{pmatrix} \\
 &\text{outbound electron spin up} & &\text{outbound electron spin down}
 \end{aligned}$$

The scattering probability amplitude  $\mathcal{M}_{ab}$  for spin state  $ab$  is

$$\mathcal{M}_{ba} = \mathcal{M}_{1ab} + \mathcal{M}_{2ab}$$

where

$$\mathcal{M}_{1ab} = \frac{\bar{u}_{4b}(-ie\gamma^\mu)(\not{p}_1 + m)(-ie\gamma^\nu)u_{2a}}{s - m^2}, \quad \mathcal{M}_{2ab} = \frac{\bar{u}_{4b}(-ie\gamma^\nu)(\not{p}_2 + m)(-ie\gamma^\mu)u_{2a}}{u - m^2}$$

Matrices  $\not{q}_1$  and  $\not{q}_2$  represent momentum transfer.

$$\begin{aligned}\not{q}_1 &= (p_1 + p_2)^\alpha g_{\alpha\beta} \gamma^\beta \\ \not{q}_2 &= (p_4 - p_1)^\alpha g_{\alpha\beta} \gamma^\beta\end{aligned}$$

Scalars  $s$  and  $u$  are Mandelstam variables.

$$\begin{aligned}s &= (p_1 + p_2)^2 \\ u &= (p_1 - p_4)^2\end{aligned}$$

In component form (note that indices  $\mu$  and  $\nu$  are interchanged for  $\mathcal{M}_{2ab}$ )

$$\begin{aligned}(\mathcal{M}_{1ab})^{\mu\nu} &= \frac{(\bar{u}_{4b})_\alpha (-ie\gamma^{\mu\alpha}_\beta)(\not{q}_1 + m)^\beta_\rho (-ie\gamma^{\nu\rho}_\sigma)(u_{2a})^\sigma}{s - m^2} \\ (\mathcal{M}_{2ab})^{\nu\mu} &= \frac{(\bar{u}_{4b})_\alpha (-ie\gamma^{\nu\alpha}_\beta)(\not{q}_2 + m)^\beta_\rho (-ie\gamma^{\mu\rho}_\sigma)(u_{2a})^\sigma}{u - m^2}\end{aligned}$$

The expected probability density  $\langle |\mathcal{M}|^2 \rangle$  is the average of spin states.

$$\langle |\mathcal{M}|^2 \rangle = \frac{1}{4} \sum_{a=1}^2 \sum_{b=1}^2 |\mathcal{M}_{ab}|^2$$

Hence

$$\langle |\mathcal{M}|^2 \rangle = \frac{1}{4} \sum_{a=1}^2 \sum_{b=1}^2 (\mathcal{M}_{1ab} \mathcal{M}_{1ab}^* + \mathcal{M}_{1ab} \mathcal{M}_{2ab}^* + \mathcal{M}_{2ab} \mathcal{M}_{1ab}^* + \mathcal{M}_{2ab} \mathcal{M}_{2ab}^*)$$

Metric tensor  $g_{\mu\nu}$  is required to sum over indices  $\mu$  and  $\nu$ .

$$\begin{aligned}\mathcal{M}_{1ab} \mathcal{M}_{1ab}^* &= (\mathcal{M}_{1ab})^{\mu\nu} (\mathcal{M}_{1ab}^*)_{\mu\nu} = (\mathcal{M}_{1ab})^{\mu\nu} [g_{\mu\alpha} (\mathcal{M}_{1ab}^*)^{\alpha\beta} g_{\beta\nu}] \\ \mathcal{M}_{1ab} \mathcal{M}_{2ab}^* &= (\mathcal{M}_{1ab})^{\mu\nu} (\mathcal{M}_{2ab}^*)_{\nu\mu} = (\mathcal{M}_{1ab})^{\mu\nu} [g_{\nu\alpha} (\mathcal{M}_{2ab}^*)^{\alpha\beta} g_{\beta\mu}] \\ \mathcal{M}_{2ab} \mathcal{M}_{2ab}^* &= (\mathcal{M}_{2ab})^{\nu\mu} (\mathcal{M}_{2ab}^*)_{\nu\mu} = (\mathcal{M}_{2ab})^{\nu\mu} [g_{\nu\alpha} (\mathcal{M}_{2ab}^*)^{\alpha\beta} g_{\beta\mu}]\end{aligned}$$

The Casimir trick uses matrix arithmetic to sum over spin states.

$$\langle |\mathcal{M}|^2 \rangle = \frac{e^4}{4} \left( \frac{f_{11}}{(s - m^2)^2} + \frac{2f_{12}}{(s - m^2)(u - m^2)} + \frac{f_{22}}{(u - m^2)^2} \right) \quad (1)$$

where

$$\begin{aligned}f_{11} &= \text{Tr} \left( (\not{p}_2 + m) \gamma^\mu (\not{q}_1 + m) \gamma^\nu (\not{p}_4 + m) \gamma_\nu (\not{q}_1 + m) \gamma_\mu \right) \\ f_{12} &= \text{Tr} \left( (\not{p}_2 + m) \gamma^\mu (\not{q}_2 + m) \gamma^\nu (\not{p}_4 + m) \gamma_\mu (\not{q}_1 + m) \gamma_\nu \right) \\ f_{22} &= \text{Tr} \left( (\not{p}_2 + m) \gamma^\mu (\not{q}_2 + m) \gamma^\nu (\not{p}_4 + m) \gamma_\nu (\not{q}_2 + m) \gamma_\mu \right)\end{aligned}$$

The trace operator  $\text{Tr}$  is the sum of diagonal matrix elements.

The following formulas are equivalent to the Casimir trick. (Recall that  $a \cdot b = a^\mu g_{\mu\nu} b^\nu$ .)

$$\begin{aligned} f_{11} &= 32(p_1 \cdot p_2)(p_1 \cdot p_4) + 64m^2(p_1 \cdot p_2) - 32m^2(p_1 \cdot p_3) - 32m^2(p_1 \cdot p_4) + 32m^4 \\ f_{12} &= 16m^2(p_1 \cdot p_2) - 16m^2(p_1 \cdot p_4) + 32m^4 \\ f_{22} &= 32(p_1 \cdot p_2)(p_1 \cdot p_4) + 32m^2(p_1 \cdot p_2) - 32m^2(p_1 \cdot p_3) - 64m^2(p_1 \cdot p_4) + 32m^4 \end{aligned} \quad (2)$$

In Mandelstam variables

$$\begin{aligned} f_{11} &= -8su + 24sm^2 + 8um^2 + 8m^4 \\ f_{12} &= 8sm^2 + 8um^2 + 16m^4 \\ f_{22} &= -8su + 8sm^2 + 24um^2 + 8m^4 \end{aligned} \quad (3)$$

Scattering experiments are typically done in the lab frame. Define Lorentz boost  $\Lambda$  for transforming momentum vectors to the lab frame.

$$\Lambda = \begin{pmatrix} E/m & 0 & 0 & \omega/m \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ \omega/m & 0 & 0 & E/m \end{pmatrix}$$

The electron is at rest in the lab frame.

$$\Lambda p_2 = \begin{pmatrix} m \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

Mandelstam variables are invariant under a boost.

$$\begin{aligned} s &= (p_1 + p_2)^2 = (\Lambda p_1 + \Lambda p_2)^2 \\ t &= (p_1 - p_3)^2 = (\Lambda p_1 - \Lambda p_3)^2 \\ u &= (p_1 - p_4)^2 = (\Lambda p_1 - \Lambda p_4)^2 \end{aligned} \quad (4)$$

In the lab frame, let  $\omega_L$  be the angular frequency of the incident photon and let  $\omega'_L$  be the angular frequency of the scattered photon.

$$\begin{aligned} \omega_L &= \Lambda p_1 \cdot \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = \frac{\omega^2}{m} + \frac{\omega E}{m} \\ \omega'_L &= \Lambda p_3 \cdot \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = \frac{\omega^2 \cos \theta}{m} + \frac{\omega E}{m} \end{aligned}$$

It can be shown that

$$\begin{aligned} s &= m^2 + 2m\omega_L \\ t &= 2m(\omega'_L - \omega_L) \\ u &= m^2 - 2m\omega'_L \end{aligned} \quad (5)$$

Then by (1), (3), and (5) we have

$$\langle |\mathcal{M}|^2 \rangle = 2e^4 \left( \frac{\omega_L}{\omega'_L} + \frac{\omega'_L}{\omega_L} + \left( \frac{m}{\omega_L} - \frac{m}{\omega'_L} + 1 \right)^2 - 1 \right) \quad (6)$$

Lab scattering angle  $\theta_L$  is given by the Compton equation

$$\cos \theta_L = \frac{m}{\omega_L} - \frac{m}{\omega'_L} + 1$$

Hence

$$\begin{aligned} \langle |\mathcal{M}|^2 \rangle &= 2e^4 \left( \frac{\omega_L}{\omega'_L} + \frac{\omega'_L}{\omega_L} + \cos^2 \theta_L - 1 \right) \\ &= 2e^4 \left( \frac{\omega_L}{\omega'_L} + \frac{\omega'_L}{\omega_L} - \sin^2 \theta_L \right) \end{aligned}$$

Now that we have derived  $\langle |\mathcal{M}|^2 \rangle$  we can investigate the angular distribution of scattered photons. For simplicity let us drop the  $L$  subscript from lab variables. From now on the symbols  $\omega$ ,  $\omega'$ , and  $\theta$  will be lab frame variables.

The differential cross section is

$$\frac{d\sigma}{d\Omega} = \frac{1}{4(4\pi\epsilon_0)^2 s} \left( \frac{\omega'}{\omega} \right)^2 \langle |\mathcal{M}|^2 \rangle$$

where

$$s = m^2 + 2m\omega = (mc^2)^2 + 2(mc^2)(\hbar\omega)$$

and  $\omega'$  is given by the Compton equation

$$\omega' = \frac{\omega}{1 + \frac{\hbar\omega}{mc^2}(1 - \cos \theta)}$$

For the lab frame we have

$$\langle |\mathcal{M}|^2 \rangle = 2e^4 \left( \frac{\omega}{\omega'} + \frac{\omega'}{\omega} - \sin^2 \theta \right)$$

Hence in the lab frame

$$\frac{d\sigma}{d\Omega} = \frac{e^4}{2(4\pi\epsilon_0)^2 s} \left( \frac{\omega'}{\omega} \right)^2 \left( \frac{\omega}{\omega'} + \frac{\omega'}{\omega} - \sin^2 \theta \right)$$

Substituting

$$e^2 = 4\pi\epsilon_0\alpha\hbar c$$

we have

$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2(\hbar c)^2}{2s} \left( \frac{\omega'}{\omega} \right)^2 \left( \frac{\omega}{\omega'} + \frac{\omega'}{\omega} - \sin^2 \theta \right)$$

which is the Klein-Nishina formula.