

## Alpha decay 2

Let  $\nu$  be the frequency of tunneling attempts. Then the decay rate  $\lambda$  is

$$\lambda = \nu e^{-2\gamma}$$

Gamow theorized that

$$\nu = \frac{v}{2r_1}$$

Velocity  $v$  is unknown but we can guess that  $v$  is proportional to  $\sqrt{E/m}$  and write

$$\lambda = \frac{\beta \sqrt{E/m}}{2r_1} e^{-2\gamma}$$

Coefficient  $\beta$  is estimated to be

$$\beta = 843$$

The half-life  $\tau_{1/2}$  is

$$\tau_{1/2} = \frac{\log 2}{\lambda}$$

Hence

$$\tau_{1/2} = \frac{2 \log 2}{\beta} \frac{r_1}{\sqrt{E/m}} \exp \left( 2C_1 \frac{Z}{\sqrt{E}} - 2C_2 \sqrt{Zr_1} \right)$$

For the alpha decay process  ${}^{238}_{92}\text{U} \rightarrow {}^{234}_{90}\text{Th}$  we have

$$A = 238$$

$$Z = 90$$

$$E = 4.27 \text{ MeV}$$

Hence

$$\tau_{1/2} = 3.9 \times 10^{17} \text{ s} = 12 \times 10^9 \text{ yr}$$

The actual half-life of  ${}^{238}\text{U}$  is

$$4.5 \times 10^9 \text{ yr}$$

Eigenmath script