

# Alpha decay 1

Let  $V(r)$  be an ansatz potential for alpha decay. The alpha particle has 2 protons and the daughter nucleus has  $Z$  protons.

$$V(r) = \begin{cases} -V_0 & 0 < r < r_1 \\ \frac{2Z\alpha\hbar c}{r} & r > r_1 \end{cases}$$

Let  $V(r_2)$  have energy  $E$ .

$$V(r_2) = \frac{2Z\alpha\hbar c}{r_2} = E$$

Hence the region from  $r_1$  to  $r_2$  is a barrier potential for an alpha particle with energy  $E$ .

Using the WKB method we have for tunneling probability  $T$

$$T \approx e^{-2\gamma}$$

where

$$\gamma = \frac{\sqrt{2m}}{\hbar} \int_{r_1}^{r_2} \sqrt{V(r) - E} dr$$

Factor out  $\sqrt{E}$  to obtain

$$\begin{aligned} \gamma &= \frac{\sqrt{2mE}}{\hbar} \int_{r_1}^{r_2} \sqrt{\frac{V(r)}{E} - 1} dr \\ &= \frac{\sqrt{2mE}}{\hbar} \int_{r_1}^{r_2} \sqrt{\frac{r_2}{r} - 1} dr \end{aligned}$$

Use the substitution

$$r = r_2 \sin^2 u \quad dr = 2r_2 \sin u \cos u du$$

to obtain

$$\begin{aligned} \gamma &= \frac{\sqrt{2mE}}{\hbar} 2r_2 \int_{u_1}^{u_2} \sqrt{\frac{1}{\sin^2 u} - 1} \sin u \cos u du \\ &= \frac{\sqrt{2mE}}{\hbar} 2r_2 \int_{u_1}^{u_2} \sqrt{1 - \sin^2 u} \cos u du \\ &= \frac{\sqrt{2mE}}{\hbar} 2r_2 \int_{u_1}^{u_2} \cos^2 u du \end{aligned}$$

where

$$u_1 = \arcsin \sqrt{r_1/r_2} \quad u_2 = \frac{\pi}{2}$$

Solve the integral to obtain

$$\gamma = \frac{\sqrt{2mE}}{\hbar} 2r_2 \left[ \frac{u}{2} + \frac{\sin(2u)}{4} \right]_{u_1}^{u_2} \quad (1)$$

Evaluate the limits.

$$\gamma = \frac{\sqrt{2mE}}{\hbar} 2r_2 \left[ \frac{\pi}{4} - \frac{1}{2} \arcsin \sqrt{r_1/r_2} - \frac{1}{4} \sin \left( 2 \arcsin \sqrt{r_1/r_2} \right) \right] \quad (2)$$

From the identities

$$\sin(2x) = 2 \sin x \cos x$$

and

$$\cos(\arcsin x) = \sqrt{1 - x^2}$$

the final term simplifies as

$$\frac{1}{4} \sin \left( 2 \arcsin \sqrt{r_1/r_2} \right) = \frac{1}{2} \sqrt{r_1/r_2} \sqrt{1 - r_1/r_2}$$

Hence

$$\gamma = \frac{\sqrt{2mE}}{\hbar} 2r_2 \left( \frac{\pi}{4} - \frac{1}{2} \arcsin \sqrt{r_1/r_2} - \frac{1}{2} \sqrt{r_1/r_2} \sqrt{1 - r_1/r_2} \right)$$

By equation (4) below we have  $r_1 \ll r_2$  hence we use the approximations

$$\begin{aligned} \arcsin \sqrt{r_1/r_2} &\approx \sqrt{r_1/r_2} \\ \sqrt{r_1/r_2} \sqrt{1 - r_1/r_2} &\approx \sqrt{r_1/r_2} \end{aligned}$$

and simplify  $\gamma$  as

$$\gamma = \frac{\sqrt{2mE}}{\hbar} 2r_2 \left( \frac{\pi}{4} - \sqrt{r_1/r_2} \right)$$

Distribute the  $r_2$ .

$$\gamma = \frac{2\sqrt{2mE}}{\hbar} \left( \frac{\pi r_2}{4} - \sqrt{r_1 r_2} \right)$$

Substitute

$$r_2 = \frac{2Z\alpha\hbar c}{E}$$

to obtain

$$\gamma = C_1 \frac{Z}{\sqrt{E}} - C_2 \sqrt{Z r_1} \quad (3)$$

where

$$C_1 = \sqrt{2m} \pi \alpha c \quad C_2 = 4 \sqrt{\frac{m \alpha c}{\hbar}}$$

For the alpha decay process  ${}^{238}_{92}\text{U} \rightarrow {}^{234}_{90}\text{Th}$  we have

$$A = 238$$

$$Z = 90$$

$$E = 4.27 \text{ MeV}$$

$$m = 4.0015 \text{ u}$$

Hence

$$\begin{aligned} r_1 &= A^{1/3} \times 1.2 \times 10^{-15} \text{ m} &= 7.44 \times 10^{-15} \text{ m} \\ r_2 &= \frac{2Z\alpha\hbar c}{E} &= 60.7 \times 10^{-15} \text{ m} \end{aligned} \tag{4}$$

Then for  $\gamma$  we obtain

$$\gamma = 47.8$$

For tunneling probability  $T$  we have

$$T = e^{-2\gamma} = 3.17 \times 10^{-42}$$

Eigenmath script