

Dirac equation 2

From Fitzpatrick problem 11.11. Let

$$\psi^r = \exp\left(-\frac{i}{\hbar}\epsilon_r p_\mu x^\mu\right) w^r$$

where $\epsilon = (1, 1, -1, -1)$ and

$$\begin{aligned} w^1 &= \sqrt{\frac{E + mc^2}{2mc^2}} \begin{pmatrix} 1 \\ 0 \\ \frac{p_z c}{E + mc^2} \\ \frac{(p_x + ip_y)c}{E + mc^2} \end{pmatrix} & w^2 &= \sqrt{\frac{E + mc^2}{2mc^2}} \begin{pmatrix} 0 \\ 1 \\ \frac{(p_x - ip_y)c}{E + mc^2} \\ \frac{-p_z c}{E + mc^2} \end{pmatrix} \\ w^3 &= \sqrt{\frac{E + mc^2}{2mc^2}} \begin{pmatrix} \frac{p_z c}{E + mc^2} \\ \frac{(p_x + ip_y)c}{E + mc^2} \\ 1 \\ 0 \end{pmatrix} & w^4 &= \sqrt{\frac{E + mc^2}{2mc^2}} \begin{pmatrix} \frac{(p_x - ip_y)c}{E + mc^2} \\ \frac{-p_z c}{E + mc^2} \\ 0 \\ 1 \end{pmatrix} \end{aligned}$$

Verify ψ^r are solutions to the Dirac equation

$$i\hbar\gamma^\mu\partial_\mu\psi = mc^2\psi$$

where

$$\gamma^\mu\partial_\mu = \gamma^0\frac{\partial}{\partial t} + c\gamma^1\frac{\partial}{\partial x} + c\gamma^2\frac{\partial}{\partial y} + c\gamma^3\frac{\partial}{\partial z}$$

Verify ψ^r are normalized as

$$|\psi|^2 = \frac{E}{mc^2}$$

Eigenmath script