

Born approximation 1

The Born approximation takes potential energy $V(\mathbf{x})$ and gives back $\psi(\mathbf{x})$ as

$$\psi(\mathbf{x}) = -\frac{m}{2\pi\hbar^2} \frac{\exp(ik|\mathbf{x}|)}{|\mathbf{x}|} \int \exp\left(\frac{i\mathbf{p} \cdot \mathbf{x}}{\hbar}\right) V(\mathbf{x}) d\mathbf{x}$$

where

$$k = \frac{\sqrt{2mE}}{\hbar}$$

Vector $\mathbf{p}$ is momentum transfer

$$\mathbf{p} = \mathbf{p}_i - \mathbf{p}_f$$

Let us derive the Born approximation. Start with the Schrodinger equation

$$-\frac{\hbar^2}{2m} \nabla^2 \psi(\mathbf{x}) + V(\mathbf{x})\psi(\mathbf{x}) = E\psi(\mathbf{x})$$

Multiply through by $-2m/\hbar^2$ to obtain

$$\nabla^2 \psi(\mathbf{x}) - \frac{2m}{\hbar^2} V(\mathbf{x})\psi(\mathbf{x}) = -\frac{2mE}{\hbar^2} \psi(\mathbf{x})$$

Interchange E and V terms.

$$\nabla^2 \psi(\mathbf{x}) + \frac{2mE}{\hbar^2} \psi(\mathbf{x}) = \frac{2m}{\hbar^2} V(\mathbf{x})\psi(\mathbf{x})$$

Let

$$k = \frac{\sqrt{2mE}}{\hbar}$$

and rewrite as

$$(\nabla^2 + k^2) \psi(\mathbf{x}) = \frac{2m}{\hbar^2} V(\mathbf{x})\psi(\mathbf{x})$$

By the sifting property of $\delta(\mathbf{x} - \mathbf{y})$ we can make the right-hand side an integral over $\mathbf{y}$.

$$(\nabla^2 + k^2) \psi(\mathbf{x}) = \frac{2m}{\hbar^2} \int \delta(\mathbf{x} - \mathbf{y}) V(\mathbf{y})\psi(\mathbf{y}) d\mathbf{y}$$

Let $G(\mathbf{x})$ be a Green's function such that

$$(\nabla^2 + k^2) G(\mathbf{x}) = \delta(\mathbf{x})$$

Then by substitution

$$(\nabla^2 + k^2) \psi(\mathbf{x}) = \frac{2m}{\hbar^2} \int (\nabla^2 + k^2) G(\mathbf{x} - \mathbf{y}) V(\mathbf{y})\psi(\mathbf{y}) d\mathbf{y}$$

By linearity the integral can be factored as

$$(\nabla^2 + k^2) \psi(\mathbf{x}) = (\nabla^2 + k^2) \frac{2m}{\hbar^2} \int G(\mathbf{x} - \mathbf{y}) V(\mathbf{y})\psi(\mathbf{y}) d\mathbf{y}$$

Hence

$$\psi(\mathbf{x}) = \frac{2m}{\hbar^2} \int G(\mathbf{x} - \mathbf{y}) V(\mathbf{y}) \psi(\mathbf{y}) d\mathbf{y}$$

The solution for $G(\mathbf{x})$ is

$$G(\mathbf{x}) = -\frac{\exp(ik|\mathbf{x}|)}{4\pi|\mathbf{x}|}$$

Hence

$$\psi(\mathbf{x}) = -\frac{m}{2\pi\hbar^2} \int \frac{\exp(ik|\mathbf{x} - \mathbf{y}|)}{|\mathbf{x} - \mathbf{y}|} V(\mathbf{y}) \psi(\mathbf{y}) d\mathbf{y}$$

For scattering experiments we can use the approximation

$$\frac{\exp(ik|\mathbf{x} - \mathbf{y}|)}{|\mathbf{x} - \mathbf{y}|} V(\mathbf{y}) \approx \frac{1}{|\mathbf{x}|} \exp\left(ik|\mathbf{x}| - \frac{ik\mathbf{x} \cdot \mathbf{y}}{|\mathbf{x}|}\right) V(\mathbf{y})$$

Hence

$$\psi(\mathbf{x}) = -\frac{m}{2\pi\hbar^2} \frac{\exp(ik|\mathbf{x}|)}{|\mathbf{x}|} \int \exp\left(-\frac{ik\mathbf{x} \cdot \mathbf{y}}{|\mathbf{x}|}\right) V(\mathbf{y}) \psi(\mathbf{y}) d\mathbf{y}$$

For a first-order approximation we substitute a plane wave for $\psi(\mathbf{y})$.

$$\psi(\mathbf{y}) = \exp\left(\frac{ik\mathbf{x}' \cdot \mathbf{y}}{|\mathbf{x}'|}\right)$$

Hence

$$\psi(\mathbf{x}) = -\frac{m}{2\pi\hbar^2} \frac{\exp(ik|\mathbf{x}|)}{|\mathbf{x}|} \int \exp\left(\frac{ik\mathbf{x}' \cdot \mathbf{y}}{|\mathbf{x}'|} - \frac{ik\mathbf{x} \cdot \mathbf{y}}{|\mathbf{x}|}\right) V(\mathbf{y}) d\mathbf{y}$$

Let $\mathbf{p}$ be momentum transfer

$$\mathbf{p} = \mathbf{p}_i - \mathbf{p}_f = \hbar k \left(\frac{\mathbf{x}'}{|\mathbf{x}'|} - \frac{\mathbf{x}}{|\mathbf{x}|} \right)$$

Then by substitution

$$\psi(\mathbf{x}) = -\frac{m}{2\pi\hbar^2} \frac{\exp(ik|\mathbf{x}|)}{|\mathbf{x}|} \int \exp\left(\frac{i\mathbf{p} \cdot \mathbf{y}}{\hbar}\right) V(\mathbf{y}) d\mathbf{y}$$

In polar coordinates

$$\psi(\mathbf{r}) = -\frac{m}{2\pi\hbar^2} \frac{\exp(ikr)}{r} \int \exp\left(\frac{i\mathbf{p} \cdot \mathbf{y}}{\hbar}\right) V(\mathbf{y}) d\mathbf{y}$$

We now consider the differential cross section $d\sigma$ such that

$$d\sigma = |\psi(\mathbf{r})|^2 r^2 d\Omega$$

where

$$|\psi(\mathbf{r})|^2 = \left| -\frac{m}{2\pi\hbar^2} \frac{\exp(ikr)}{r} \int \exp\left(\frac{i\mathbf{p} \cdot \mathbf{x}}{\hbar}\right) V(\mathbf{x}) d\mathbf{x} \right|^2$$

The factor $|\exp(ikr)/r|^2 = 1/r^2$ cancels with r^2 in $d\sigma$ hence

$$d\sigma = \left| -\frac{m}{2\pi\hbar^2} \int \exp\left(\frac{i\mathbf{p} \cdot \mathbf{x}}{\hbar}\right) V(\mathbf{x}) d\mathbf{x} \right|^2 d\Omega$$

Let $f(\theta, \phi)$ be the scattering amplitude

$$f(\theta, \phi) = -\frac{m}{2\pi\hbar^2} \int \exp\left(\frac{i\mathbf{p} \cdot \mathbf{x}}{\hbar}\right) V(\mathbf{x}) d\mathbf{x}$$

so that for the Born approximation

$$\frac{d\sigma}{d\Omega} = |f(\theta, \phi)|^2$$

For example, for Coulomb potentials we have

$$f(\theta, \phi) = \frac{Z\alpha\hbar c}{2E(1 - \cos\theta)}$$

Hence the Rutherford cross section

$$\frac{d\sigma}{d\Omega} = |f(\theta, \phi)|^2 = \left[\frac{Z\alpha\hbar c}{2E(1 - \cos\theta)} \right]^2$$

Eigenmath script