
Compton scattering 1

Compton scattering is the process $e^- + \gamma \to e^- + \gamma$.



Momentum vectors where $E = \sqrt{\omega^2 + m^2}$.

$$p_1 = \underset{e^- \longrightarrow}{\begin{pmatrix} E \\ 0 \\ 0 \\ \omega \end{pmatrix}} \qquad p_2 = \underset{\longleftarrow \gamma}{\begin{pmatrix} \omega \\ 0 \\ 0 \\ -\omega \end{pmatrix}} \qquad p_3 = \underset{\nearrow}{\overset{e^-}{\begin{pmatrix} E \\ \omega \sin\theta \cos\phi \\ \omega \sin\theta \sin\phi \\ \omega \cos\theta \end{pmatrix}}} \qquad p_4 = \underset{\gamma \swarrow}{\begin{pmatrix} \omega \\ -\omega \sin\theta \cos\phi \\ -\omega \sin\theta \sin\phi \\ -\omega \cos\theta \end{pmatrix}}$$

Spinors for p_1.

$$u_{11} = \frac{1}{\sqrt{E+m}} \underset{\text{spin up}}{\begin{pmatrix} E+m \\ 0 \\ \omega \\ 0 \end{pmatrix}} \qquad u_{12} = \frac{1}{\sqrt{E+m}} \underset{\text{spin down}}{\begin{pmatrix} 0 \\ E+m \\ 0 \\ -\omega \end{pmatrix}}$$

Spinors for p_3.

$$u_{31} = \frac{1}{\sqrt{E+m}} \underset{\text{spin up}}{\begin{pmatrix} E+m \\ 0 \\ p_{3z} \\ p_{3x} + ip_{3y} \end{pmatrix}} \qquad u_{32} = \frac{1}{\sqrt{E+m}} \underset{\text{spin down}}{\begin{pmatrix} 0 \\ E+m \\ p_{3x} - ip_{3y} \\ -p_{3z} \end{pmatrix}}$$

The scattering amplitude $\mathcal{M}$ is

$$\mathcal{M} = \mathcal{M}_1 + \mathcal{M}_2$$

where $\mathcal{M}_1$ is the s-channel amplitude

$$\mathcal{M}_1 = \frac{\bar{u}_{3b}(-ie\gamma^\mu)(\not{q}_1 + m)(-ie\gamma^\nu)u_{1a}}{s - m^2}$$

and $\mathcal{M}_2$ is the u-channel amplitude

$$\mathcal{M}_2 = \frac{\bar{u}_{3b}(-ie\gamma^\nu)(\not{q}_2 + m)(-ie\gamma^\mu)u_{1a}}{u - m^2}$$

Matrices $\not{q}_1$ and $\not{q}_2$ are momentum transfer.

$$\not{q}_1 = (p_1 + p_2)^\mu g_{\mu\nu}\gamma^\nu$$
$$\not{q}_2 = (p_1 - p_4)^\mu g_{\mu\nu}\gamma^\nu$$

Expected density $\langle|\mathcal{M}|^2\rangle$ is the average density.

$$\langle|\mathcal{M}|^2\rangle = \frac{1}{4}\sum_{ab}|\mathcal{M}|^2 = \frac{1}{4}\sum_{ab}(\mathcal{M}_1 + \mathcal{M}_2)(\mathcal{M}_1 + \mathcal{M}_2)^*$$

Expand the summand. (Note indices μ and ν of $\mathcal{M}_2$ are interchanged relative to $\mathcal{M}_1$.)

$$\langle|\mathcal{M}|^2\rangle = \frac{1}{4}\sum_{ab}\left(\mathcal{M}_1{}^{\mu\nu}\mathcal{M}^*_{1\mu\nu} + \mathcal{M}_1{}^{\mu\nu}\mathcal{M}^*_{2\nu\mu} + \mathcal{M}_2{}^{\nu\mu}\mathcal{M}^*_{1\mu\nu} + \mathcal{M}_2{}^{\nu\mu}\mathcal{M}^*_{2\nu\mu}\right)$$

The summation results are

$$\sum_{ab}\mathcal{M}_1{}^{\mu\nu}\mathcal{M}^*_{1\mu\nu} = \frac{e^4}{(s-m^2)^2}\left(-8su + 24m^2s + 8m^2u + 8m^4\right)$$
$$\sum_{ab}\mathcal{M}_1{}^{\mu\nu}\mathcal{M}^*_{2\nu\mu} = \frac{e^4}{(s-m^2)(u-m^2)}\left(8m^2s + 8m^2u + 16m^4\right)$$
$$\sum_{ab}\mathcal{M}_2{}^{\nu\mu}\mathcal{M}^*_{2\nu\mu} = \frac{e^4}{(u-m^2)^2}\left(-8su + 8m^2s + 24m^2u + 8m^4\right)$$

where s, t, and u are Mandelstam variables

$$s = (p_1 + p_2)^2$$
$$t = (p_1 - p_3)^2$$
$$u = (p_1 - p_4)^2$$

For clarity define

$$f_{11} = -8su + 24m^2s + 8m^2u + 8m^4$$
$$f_{12} = 8m^2s + 8m^2u + 16m^4$$
$$f_{22} = -8su + 8m^2s + 24m^2u + 8m^4$$

and write $\langle|\mathcal{M}|^2\rangle$ more compactly as

$$\langle|\mathcal{M}|^2\rangle = \frac{e^4}{4}\left(\frac{f_{11}}{(s-m^2)^2} + \frac{2f_{12}}{(s-m^2)(u-m^2)} + \frac{f_{22}}{(u-m^2)^2}\right)$$

For $E \gg m$ a useful approximation is to set $m = 0$ and obtain

$$f_{11} = -8su$$
$$f_{12} = 0$$
$$f_{22} = -8su$$

Then $\langle|\mathcal{M}|^2\rangle$ simplifies as

$$\langle|\mathcal{M}|^2\rangle = \frac{e^4}{4}\left(-\frac{8su}{s^2} - \frac{8su}{u^2}\right) = -2e^4\left(\frac{u}{s} + \frac{s}{u}\right)$$

For $m = 0$ the Mandelstam variables are

$$\begin{aligned} s &= 4E^2 \\ t &= -2E^2(1 - \cos\theta) \\ u &= -2E^2(1 + \cos\theta) \end{aligned}$$

Hence

$$\langle|\mathcal{M}|^2\rangle = 2e^4\left(\frac{1 + \cos\theta}{2} + \frac{2}{1 + \cos\theta}\right)$$

The differential cross section is

$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{e^4}\frac{\langle|\mathcal{M}|^2\rangle}{4s} = \frac{\alpha^2}{8E^2}\left(\frac{1 + \cos\theta}{2} + \frac{2}{1 + \cos\theta}\right)$$

Multiply by $(\hbar c)^2$ to convert from natural units to physical units.

$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{8E^2}\left(\frac{1 + \cos\theta}{2} + \frac{2}{1 + \cos\theta}\right)(\hbar c)^2$$

Alternate form of $d\sigma$.

$$\frac{d\sigma}{d\cos\theta} = 2\pi\frac{d\sigma}{d\Omega} = \frac{\pi\alpha^2}{s}\left(\frac{\cos\theta + 1}{2} + \frac{2}{\cos\theta + 1}\right)(\hbar c)^2$$

See equation (1) of arxiv.org/pdf/hep-ex/0504012.

Eigenmath script