

Green's function

We seek a Green's function $G(\mathbf{x})$ such that

$$(\nabla^2 + k^2) G(\mathbf{x}) = \delta^3(\mathbf{x}) \quad (1)$$

Let $g(\mathbf{y})$ be the Fourier transform of $G(\mathbf{x})$.

$$G(\mathbf{x}) = \frac{1}{(2\pi)^{\frac{3}{2}}} \int \exp(i\mathbf{x} \cdot \mathbf{y}) g(\mathbf{y}) d\mathbf{y} \quad (2)$$

Substitute (2) into (1).

$$(\nabla^2 + k^2) \left(\frac{1}{(2\pi)^{\frac{3}{2}}} \int \exp(i\mathbf{x} \cdot \mathbf{y}) g(\mathbf{y}) d\mathbf{y} \right) = \delta^3(\mathbf{x})$$

By linearity of differentiation, $(\nabla^2 + k^2)$ can be moved inside the integral.

$$\frac{1}{(2\pi)^{\frac{3}{2}}} \int (\nabla^2 + k^2) \exp(i\mathbf{x} \cdot \mathbf{y}) g(\mathbf{y}) d\mathbf{y} = \delta^3(\mathbf{x}) \quad (3)$$

Noting that

$$\nabla^2 \exp(i\mathbf{x} \cdot \mathbf{y}) = -y^2 \exp(i\mathbf{x} \cdot \mathbf{y})$$

and

$$\delta^3(\mathbf{x}) = \frac{1}{(2\pi)^3} \int \exp(i\mathbf{x} \cdot \mathbf{y}) d\mathbf{y}$$

we have by substitution into (3)

$$\frac{1}{(2\pi)^{\frac{3}{2}}} \int (-y^2 + k^2) \exp(i\mathbf{x} \cdot \mathbf{y}) g(\mathbf{y}) d\mathbf{y} = \frac{1}{(2\pi)^3} \int \exp(i\mathbf{x} \cdot \mathbf{y}) d\mathbf{y}$$

Solve for $g(\mathbf{y})$.

$$g(\mathbf{y}) = \frac{1}{(2\pi)^{\frac{3}{2}} (k^2 - y^2)}$$

Substitute for $g(\mathbf{y})$ in (2) to obtain

$$G(\mathbf{x}) = \frac{1}{(2\pi)^3} \int \frac{\exp(i\mathbf{x} \cdot \mathbf{y})}{k^2 - y^2} d\mathbf{y}$$

Change to polar coordinates where $x = |\mathbf{x}|$, $y = |\mathbf{y}|$, and θ and ϕ are the angular distance from $\mathbf{x}$ to $\mathbf{y}$.

$$G(\mathbf{x}) = \frac{1}{(2\pi)^3} \int_0^\infty \int_0^\pi \int_0^{2\pi} \frac{\exp(ixy \cos \theta)}{k^2 - y^2} y^2 \sin \theta dy d\theta d\phi$$

For the integrals over θ and ϕ we have

$$\int_0^\pi \int_0^{2\pi} \exp(ixy \cos \theta) \sin \theta d\theta d\phi = \frac{4\pi \sin(xy)}{xy}$$

Hence

$$G(\mathbf{x}) = \frac{1}{2\pi^2 x} \int_0^\infty \frac{y \sin(xy)}{k^2 - y^2} dy$$

Noting that $y \sin(xy)$ is an even function of y we can change the integral limits as follows.

$$G(\mathbf{x}) = \frac{1}{4\pi^2 x} \int_{-\infty}^\infty \frac{y \sin(xy)}{k^2 - y^2} dy$$

Negate the denominator.

$$G(\mathbf{x}) = \frac{1}{4\pi^2 x} \int_{-\infty}^\infty -\frac{y \sin(xy)}{y^2 - k^2} dy$$

Change the sine function to exponential form and factor the denominator.

$$G(\mathbf{x}) = \frac{i}{8\pi^2 x} \left[\int_{-\infty}^\infty \frac{y \exp(ixy)}{(y - k)(y + k)} dy - \int_{-\infty}^\infty \frac{y \exp(-ixy)}{(y - k)(y + k)} dy \right]$$

By Cauchy's integral formula we have

$$\int_{-\infty}^\infty \frac{y \exp(ixy)}{y + k} \frac{1}{y - k} dy = i\pi \exp(ikx)$$

and

$$\int_{-\infty}^\infty \frac{y \exp(-ixy)}{y - k} \frac{1}{y + k} dy = -i\pi \exp(ikx)$$

Hence

$$G(\mathbf{x}) = -\frac{e^{ikx}}{4\pi x}$$

Substitute $\mathbf{r}$ for $\mathbf{x}$.

$$G(\mathbf{r}) = -\frac{e^{ikr}}{4\pi r} \tag{4}$$

Let us verify $G(\mathbf{r})$ in (4) satisfies equation (1). We will need the following formula from Griffiths and Schroeter problem 10.8.

$$\nabla^2(1/r) = -4\pi\delta^3(\mathbf{r}) \tag{5}$$

Recall that $\nabla^2 = \nabla \cdot \nabla$ and for scalar functions f and g we have

$$\nabla \cdot (f \nabla g) = \nabla f \cdot \nabla g + f \nabla^2 g$$

Hence we have for the Laplacian of $G(\mathbf{r})$

$$\begin{aligned} \nabla^2 G(\mathbf{r}) &= \nabla \cdot \nabla \left(-\frac{e^{ikr}}{4\pi r} \right) \\ &= -\frac{1}{4\pi} \nabla \cdot \left(\frac{1}{r} \nabla e^{ikr} + e^{ikr} \nabla \frac{1}{r} \right) \\ &= -\frac{1}{4\pi} \left(\nabla \frac{1}{r} \cdot \nabla e^{ikr} + \frac{1}{r} \nabla^2 e^{ikr} + \nabla e^{ikr} \cdot \nabla \frac{1}{r} + e^{ikr} \nabla^2 \frac{1}{r} \right) \end{aligned} \tag{6}$$

In spherical coordinates the gradient of f is

$$\nabla f = \left(\frac{\partial f}{\partial r} \right) \mathbf{e}_r + \left(\frac{1}{r} \frac{\partial f}{\partial \theta} \right) \mathbf{e}_\theta + \left(\frac{1}{r \sin \theta} \frac{\partial f}{\partial \phi} \right) \mathbf{e}_\phi$$

hence

$$\nabla \frac{1}{r} \cdot \nabla e^{ikr} = \left(-\frac{1}{r^2} \mathbf{e}_r + 0 \mathbf{e}_\theta + 0 \mathbf{e}_\phi \right) \cdot \left(ik e^{ikr} \mathbf{e}_r + 0 \mathbf{e}_\theta + 0 \mathbf{e}_\phi \right) = -\frac{ik e^{ikr}}{r^2} \quad (7)$$

In spherical coordinates the Laplacian of f is

$$\nabla^2 f = \frac{1}{r} \frac{\partial^2}{\partial r^2} (rf) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial f}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 f}{\partial \phi^2}$$

hence

$$\begin{aligned} \frac{1}{r} \nabla^2 e^{ikr} &= \frac{1}{r^2} \frac{\partial^2}{\partial r^2} (r e^{ikr}) \\ &= \frac{1}{r^2} \frac{\partial}{\partial r} (e^{ikr} + ikr e^{ikr}) \\ &= \frac{1}{r^2} (2ike^{ikr} - k^2 r e^{ikr}) \\ &= \frac{2ike^{ikr}}{r^2} - \frac{k^2 e^{ikr}}{r} \end{aligned} \quad (8)$$

Substitute (5), (7), and (8) into (6) to obtain

$$\nabla^2 G(\mathbf{r}) = \frac{k^2 e^{ikr}}{4\pi r} + \delta^3(\mathbf{r}) e^{ikr}$$

Then by equation (4)

$$\nabla^2 G(\mathbf{r}) = -k^2 G(\mathbf{r}) + \delta^3(\mathbf{r}) e^{ikr}$$

Noting that $e^{ikr} = 1$ at $r = 0$, the e^{ikr} term can be discarded leaving

$$\nabla^2 G(\mathbf{r}) = -k^2 G(\mathbf{r}) + \delta^3(\mathbf{r}) \quad (9)$$

Hence

$$(\nabla^2 + k^2) G(\mathbf{r}) = \delta^3(\mathbf{r})$$

Eigenmath script