

Exercise sheet

Lecturer: Georgios P. Karagiannis

georgios.karagiannis@durham.ac.uk

Part 1. Types of spatial data

Exercise 1. (★)(Columbus Columbus OH data set) Figure 2a shows the Property crime (number per thousand households) in 49 districts in Columbus in 1980, as well as the average value of the house in USD. Figure 2b presents the corresponding average house value. This is the R dataset `columbus{spdep}`. Interest may lie to find whether high rates of crime are clustered in a particular areas, and if yes, perhaps what is the association of it with the value of the houses in the area. To which principal spatial statistical are would you associate this problem?



FIGURE 1. Columbus Columbus OH spatial analysis dataset

Solution. Aerial unit data / spatial data on lattices

Exercise 2. (★)(Columbus Columbus OH data set) Figure 2a shows the Property crime (number per thousand households) in 49 districts in Columbus in 1980, as well as the average value of the house in USD. Figure 2b presents the corresponding average house value. This is the R dataset

`columbus{spdep}`. Interest may lie to find whether high rates of crime are clustered in a particular areas, and if yes, perhaps what is the association of it with the value of the houses in the area. To which principal spatial statistical are would you associate this problem?



FIGURE 2. Columbus Columbus OH spatial analysis dataset

Solution. Aerial unit data / spatial data on lattices

Exercise 3. (★)(Soil chemistry properties data set.) It contains measurements of various chemical properties of soil samples collected at different locations in a field. These properties include: the acidity or alkalinity of the soil (PH), the salt concentration in the soil (Salinity), and others. It is the R dataset `soil250{geoR}`. Figure 3 presents the locations these measurements are taken. The data (measurements) are in fixed locations at a regular grid of points. The domain scientist would be interested in the nutrient levels and pH to assess soil fertility and make recommendations for agricultural practices. The statistician could (i.) estimate/predict values of soil properties at unsampled locations based on measurements at sampled locations; and (ii.) assess the spatial variability of soil properties (nutrient levels and pH) to identify regions with high or low variability. To which principal spatial statistical are would you associate this problem?



FIGURE 3. Soil chemistry data set

Solution. Point referenced data, or geostatistical data

Exercise 4. (★)(Scallop abundance data) Figure 4 presents 148 locations (degrees of longitude & latitude) in the Atlantic waters off the coasts of New Jersey and Long Island New York as coordinates and the size of scallop catch at the corresponding location as the dot size. The sites are at fixed locations within an irregular grid of points. Sustainable scallop abundance is critical for the long-term economic viability of the fishing industry. A healthy and stable scallop population supports a consistent source of income for fishermen and related businesses. To which principal spatial statistical are would you associate this problem?



FIGURE 4. Scallop abundance data

Solution. Point referenced data, or geostatistical data

Exercise 5. (★)(Wolfcamp-aquifer data) Figure 5 presents locations and levels (in feet above sea level) of piezometric head for the aquifer; they are obtained by drilling a narrow pipe into the aquifer

and letting the water find its own level in the pipe. After rigorous screening of unsuitable wells, 85 remained. There is interest to find where the radionuclide contamination would flow from the site in Deaf Smith County, Texas. Beneath Deaf Smith County is a deep brine aquifer known as the Wolfcamp aquifer, a potential pathway for any radionuclides leaking from the repository. The predicted direction of flow can be used to determine locations of downgradient and upgradient wells for a groundwater monitoring system. A first direction in analyzing this spatial data set is to draw a map of a predicted surface based on the (irregularly located) 85 data. To which principal statistical are would you associate this problem?



FIGURE 5. Wolfcamp-aquifer data. Piezometric-head levels (feet above sea level) vs coordinates.

Solution. Point referenced data, or geostatistical data

Exercise 6. (★)(Swiss rainfall data) Figure 6 presents the locations of the 100 locations in Switzerland as dots whose size and color indicates the amount of the corresponding rainfall measurements (in 10th of mm) taken on May 8, 1986. This is the R data set `SIC{geoR}`. Observation sites are irregularly spaced, and fixed. A scientific objective may be to analyzing rainfall patterns with purpose to optimize crop planting and irrigation schedules. A statistician is able to estimate rainfall values at unsampled locations based on available measurements, create maps that represent the spatial distribution of rainfall, or quantify the uncertainty associated with rainfall estimates and predictions, which are important for risk assessment and decision-making. To which principal spatial statistical are would you associate this problem?




FIGURE 6. Swiss rainfall data

Solution. Point referenced data, or geostatistical data

Part 2. INLA

Exercise 7. (★) Consider the model

$$\begin{cases} z_i | \eta_i \sim \text{Poisson}(\exp(\eta_i)) & i = 1, \dots, n \\ \eta_i = \beta_0 + \beta_1 w_i + u_{j(i)} \\ u \sim N_m(0, I\tau^{-1}) \end{cases}$$

where $\{w_i\}$ are covariates, $j(i)$ is a known mapping from $1 : n$ to $1 : m$ (given below in the dataset as `idx`).

For training use the following data set $\{(z_i, w_i)\}_{i=1}^n$ by running

```
rm(list=ls())
# generate the dataset
set.seed(123456L)
n = 50;
m = 10
w = rnorm(n, sd = 1/3)
u = rnorm(m, sd = 1/4)
intercept = 0;
beta = 1
idx = sample(1:m, n, replace = TRUE)
z = rpois(n, lambda = exp(intercept + beta * w + u[idx]))
table(z, dnn=NULL)
```

Do the following, by using R-INLA

- (1) Run `inla{INLA}` in order to train the above model, and generate an `inla` object (that you will call it `out.inla`). For the function `inla{INLA}` specify the formula, data, and family

arguments. To approximate the conditional pdf of latent variables of the GMRF use the Gaussian approximation. For the rest parameters just use the default R-INLA options.

- (2) Print a summary of the marginal posteriors
- (3) Produce and print the marginal posterior pdf of $\Pr(\beta_1|z)$.

Solution.

- (1)

```
my.data = data.frame(z, w, idx)
formula = z ~ 1 + w + f(idx, model="iid")
out.inla = inla(formula, data = my.data,
                family = "poisson",
                control.inla = list(strategy = "gaussian")
                )
```
- (2)

```
summary(out.inla)
```

```
> res.predict$summary.linear.predictor[7,]
              mean          sd 0.025quant 0.5quant 0.975quant      mode      kld
Predictor.07 3.021652 0.1847223   2.639797 3.029161   3.362469 3.045581 1.224947e-07
```

Call:

```
c("inla.core(formula = formula, family = family, contrasts = contrasts, ", " data =
data, quantiles = quantiles, E = E, offset = offset, ", " scale = scale, weights =
weights, Ntrials = Ntrials, strata = strata, ", " lp.scale = lp.scale, link.covariates
= link.covariates, verbose = verbose, ", " lincomb = lincomb, selection = selection,
control.compute = control.compute, ", " control.predictor = control.predictor,
control.family = control.family, ", " control.inla = control.inla, control.fixed =
control.fixed, ", " control.mode = control.mode, control.expert = control.expert, ", "
control.hazard = control.hazard, control.lincomb = control.lincomb, ", " control.update
= control.update, control.lp.scale = control.lp.scale, ", " control.pardiso =
control.pardiso, only.hyperparam = only.hyperparam, ", " inla.call = inla.call,
inla.arg = inla.arg, num.threads = num.threads, ", " blas.num.threads =
blas.num.threads, keep = keep, working.directory = working.directory, ", " silent =
silent, inla.mode = inla.mode, safe = FALSE, debug = debug, ", " .parent.frame =
.parent.frame)")
```

Time used:

```
Pre = 0.763, Running = 0.224, Post = 0.0168, Total = 1
```

Fixed effects:

```
              mean          sd 0.025quant 0.5quant 0.975quant      mode kld
(Intercept) -0.069 0.153      -0.370   -0.069      0.232 -0.069  0
w            1.178 0.401        0.391    1.178      1.964  1.178  0
```

Random effects:

```
Name      Model
idx IID model
```

Model hyperparameters:

```
              mean          sd 0.025quant 0.5quant 0.975quant      mode
Precision for idx 19980.67 19912.46      599.82 13939.51   74289.61 214.74
```


Marginal log-Likelihood: -69.62

is computed

Posterior summaries for the linear predictor and the fitted values are computed

(Posterior marginals needs also 'control.compute=list(return.marginals.predictor=TRUE)')

```
(3) b_1_pdf = out.inla$marginals.fixed$w
plot(b_1_pdf[,1], b_1_pdf[,2], type="l")
```



Part 3. Point referenced data / Geostatistics

Exercise 8. (★) If $c : \mathbb{R}^d \rightarrow \mathbb{R}$ is the covariogram of a weakly stationary random field $Z = (Z_s)_{s \in \mathbb{R}^d}$ then $c(\cdot)$ is semi-positive definite; i.e. for all $n \in \mathbb{N}$, $a \in \mathbb{R}^n$, and $\{s_1, \dots, s_n\} \subseteq S$

$$\sum_{i=1}^n \sum_{j=1}^n a_i a_j c(s_i - s_j) \geq 0$$

Solution. To show that $c(\cdot)$ is semi-positive definite, I need to show that $\forall a \in \mathbb{R}^n - \{0\}$ it is

$$\sum_{i=1}^n \sum_{j=1}^n a_i a_j c(s_i - s_j) \geq 0$$

Well it is

$$\begin{aligned} 0 &\leq \text{Var} \left(\sum_{i=1}^n a_i Z(s_i) \right) = \text{Cov} \left(\sum_{i=1}^n a_i Z(s_i), \sum_{j=1}^n a_j Z(s_j) \right) \\ &= \sum_{i=1}^n a_i \sum_{j=1}^n a_j \text{Cov}(Z(s_i), Z(s_j)) \\ &= \sum_{i=1}^n a_i \sum_{j=1}^n a_j c(s_i - s_j) = \sum_{i=1}^n a_i \sum_{j=1}^n a_j c(s_i - s_j) \end{aligned}$$

Exercise 9. (★) Show that if $c_1(\cdot, \cdot)$ and $c_2(\cdot, \cdot)$ are covariance functions (are non-negative definite) then so are $c_3(\cdot, \cdot) = bc_1(\cdot, \cdot) + dc_2(\cdot, \cdot)$ and $c_4(\cdot, \cdot) = c_1(\cdot, \cdot) c_2(\cdot, \cdot)$.

Solution. For all $n \in \mathbb{N}$ and a_1, \dots, a_n

$$\begin{aligned}
\sum_{i=1}^n a_i \sum_{j=1}^n a_j c_3(s_i, s_j) &= \sum_{i=1}^n a_i \sum_{j=1}^n a_j (bc_1(s_i, s_j) + dc_2(s_i, s_j)) \\
&= \underbrace{\sum_{i=1}^n a_i \sum_{j=1}^n a_j bc_1(s_i, s_j)}_{\geq 0} + \underbrace{\sum_{i=1}^n a_i \sum_{j=1}^n a_j dc_2(s_i, s_j)}_{\geq 0} \\
&\geq 0
\end{aligned}$$

Regarding c_4 , assume independent stochastic processes $(Y_s)_{s \in S}$ and $(X_s)_{s \in S}$ with mean zero and covariance functions $c_1(\cdot, \cdot)$ and $c_2(\cdot, \cdot)$ correspondingly. Let stochastic processes $(Z_s)_{s \in S}$ with $Z_s = Y_s X_s$. Then

$$\begin{aligned}
\text{Cov}(Z_s, Z_t) &= \text{Cov}(Y_s X_s, Y_t X_t) \\
&= \text{E}(Y_s X_s Y_t X_t) \\
&= \text{E}(Y_s Y_t X_s X_t), \text{ but } Y_s \perp X_s \\
&= \text{E}(X_s X_t) \text{E}(Y_s Y_t) \\
&= \text{Cov}(X_s, X_t) \text{Cov}(Y_s, Y_t) \\
&= c_1(s, t) c_2(s, t) = c_4(s, t)
\end{aligned}$$

that is $c_4(\cdot, \cdot)$ is a covariance function of a stochastic processes.

Exercise 10. (★) Consider the Gaussian c.f. $c(h) = \sigma^2 \exp(-\beta \|h\|_2^2)$ for $\sigma^2, \beta > 0$ and $h \in \mathbb{R}^d$. Compute the spectral density from Bochner's theorem

Solution. It is

$$\begin{aligned}
f(\omega) &= \left(\frac{1}{2\pi}\right)^d \int_{\mathbb{R}^d} \exp(-i\omega^\top h) \sigma^2 \exp(-\beta \|h\|_2^2) dh \\
&= \sigma^2 \left(\frac{1}{2\pi}\right)^d \prod_{j=1}^d \int_{\mathbb{R}} \exp(-i\omega_j h_j - \beta h_j^2) dh \\
&= \sigma^2 \left(\frac{1}{2\pi}\right)^d \prod_{j=1}^d \int_{\mathbb{R}} \exp(-\beta (h_j - (-i\omega_j / (2\beta)))^2) dh_j \\
&= \sigma^2 \left(\frac{1}{4\pi\beta}\right)^{d/2} \exp(-\|\omega\|_2^2 / (4\beta))
\end{aligned}$$

Exercise 11. (★) Consider the Exponential c.f. $c(h) = \sigma^2 \exp(-\beta \|h\|_1)$ for $\sigma^2, \beta > 0$ and $h \in \mathbb{R}^d$. Compute the spectral density from Bochner's theorem

Solution. It is

$$\begin{aligned} f(\omega) &= \left(\frac{1}{2\pi}\right)^d \int_{\mathbb{R}^d} \exp(-i\omega^\top h) \sigma^2 \exp(-\beta \|h\|_1) dh \\ &= \sigma^2 \left(\frac{1}{2\pi}\right)^d \prod_{j=1}^d \int_{\mathbb{R}} \exp(-i\omega_j h_j - \beta |h_j|) dh_j \end{aligned}$$

where

$$\begin{aligned} \int_{\mathbb{R}} \exp(-i\omega_j h_j - \beta |h_j|) dh_j &= \int_{-\infty}^0 \exp(-i\omega_j h_j - \beta |h_j|) dh_j + \int_0^{\infty} \exp(-i\omega_j h_j - \beta |h_j|) dh_j \\ &= \int_{-\infty}^0 \exp(-i\omega_j h_j + \beta h_j) dh_j + \int_0^{\infty} \exp(-i\omega_j h_j - \beta h_j) dh_j \\ &= \int_{-\infty}^0 \exp(-(i\omega_j - \beta) h_j) dh_j + \int_0^{\infty} \exp(-(i\omega_j + \beta) h_j) dh_j \\ &= \int_0^{\infty} \exp(-(\beta - i\omega_j) h_j) dh_j + \int_0^{\infty} \exp(-(i\omega_j + \beta) h_j) dh_j \\ &= \frac{1}{(\beta - i\omega_j)} + \frac{1}{(\beta + i\omega_j)} = \frac{2\beta}{\beta^2 + \omega_j^2} \end{aligned}$$

hence

$$f(\omega) = \sigma^2 \left(\frac{\beta}{\pi}\right)^d \prod_{j=1}^d \frac{1}{\beta^2 + \omega_j^2}$$

(Given as Formative assessment 1)

Exercise 12. (★) Let $Z = (Z_s)_{s \in \mathbb{R}^d}$ be an intrinsically stationary stochastic process, and let $\gamma : \mathbb{R}^d \rightarrow \mathbb{R}$ be its semivariogram. Assume $a \in \mathbb{R}^n$ s.t. $\sum_{i=1}^n a_i = 0$.

(1) Let $a \in \mathbb{R}^n$ be a vector of constants. Show that

$$\text{Var} \left(\sum_{i=1}^n a_i Z(s_i) \right) = \sum_{i=1}^n \sum_{j=1}^n a_i a_j c_Y(s_i, s_j)$$

where $c_Y(s, t) = E(Y(s)Y(t))$, and $Y_s = Z_s - Z_0$.

(2) Show that

$$c_Y(s, t) = \gamma(s) + \gamma(t) - \gamma(s - t)$$

(3) Show that for all $\forall \{s_1, \dots, s_n\} \subseteq S$ it is

$$\sum_{i=1}^n \sum_{j=1}^n a_i a_j \gamma(s_i - s_j) \leq 0$$

Solution. Assume $0 \in S$, and a random variable $Z(0)$. Let $Y_s = Z_s - Z_0$.

(1) It is

$$\begin{aligned}\text{Var} \left(\sum_{i=1}^n a_i Z(s_i) \right) &= \text{Var} \left(\sum_{i=1}^n a_i Z(s_i) - \overbrace{\sum_{i=1}^n a_i Z(0)}^{0=} \right) = \text{Var} \left(\sum_{i=1}^n a_i Y(s_i) \right) \\ &= \sum_{i=1}^n \sum_{j=1}^n a_i a_j \text{E}(Y(s_i) Y(s_j)) = c_Y(s, t)\end{aligned}$$

(2) For $\text{E}(Y(s_i)) = 0$ it is

$$\begin{aligned}\gamma(s-t) &= \frac{1}{2} \text{E}(Z(s) - Z(0) + Z(t) - Z(0))^2 \\ &= \frac{1}{2} (2\gamma(s) + 2\gamma(t) - 2c_Y(s, t)) \\ \implies c_Y(s, t) &= \gamma(s) + \gamma(t) - \gamma(s-t)\end{aligned}$$

(3) It is

$$\begin{aligned}0 \leq \text{Var} \left(\sum_{i=1}^n a_i Z(s_i) \right) &= \sum_{i=1}^n \sum_{j=1}^n a_i a_j c_Y(s_i, s_j) \\ &= \sum_{i=1}^n \sum_{j=1}^n a_i a_j (\gamma(s_i) + \gamma(s_j) - \gamma(s_i - s_j)) \\ &= \sum_{i=1}^n a_i \gamma(s_i) \sum_{j=1}^n a_j + \sum_{j=1}^n a_j \gamma(s_j) \sum_{i=1}^n a_i - \sum_{i=1}^n \sum_{j=1}^n a_i a_j \gamma(s_i - s_j)\end{aligned}$$

hence

$$\sum_{i=1}^n \sum_{j=1}^n a_i a_j \gamma(s_i - s_j) \leq 0$$

(Given as Formative assessment 1)

Exercise 13. (★) Consider the zero-mean geostatistical process $Z = (Z_s)_{s \in \mathbb{R}^d}$ with a weakly stationary and isotropic covariance function given by

$$c(h) = \begin{cases} \xi^2 (1 + \rho \|h\|) \exp(-\rho \|h\|), & h > 0 \\ \nu^2 + \xi^2, & h = 0 \end{cases}$$

- (1) Compute the semi-variogram for the geostatistical process (Z_s)
- (2) What are the nugget, sill and partial sill for this covariance model? Justify your answer.
- (3) Would the slightly altered covariance function defined below be a good model for spatial data for $\phi > 0$? Justify your answer.

$$c(h) = \begin{cases} \xi^2 (1 + \rho \|h\|) \exp(-\rho \|h\|) + \phi, & h > 0 \\ \nu^2 + \xi^2 + \phi, & h = 0 \end{cases}$$

Solution.

(1) For all $h \neq 0$, it is

$$\begin{aligned} \gamma(h) &= c(0) - c(h), \\ &= \nu^2 + \xi^2 - \xi^2 (1 + \rho \|h\|) \exp(-\rho \|h\|) \\ &= \nu^2 + \xi^2 (1 - (1 + \rho \|h\|) \exp(-\rho \|h\|)) \end{aligned}$$

then

$$\gamma(h) = \begin{cases} \nu^2 + \xi^2 (1 - (1 + \rho \|h\|) \exp(-\rho \|h\|)) & h > 0 \\ 0 & h = 0 \end{cases}$$

(2)

- The sill is the covariance function at distance 0, that is $c(0) = \nu^2 + \xi^2$. Or since analogously, it is $\lim_{\|h\| \rightarrow \infty} \gamma(h)$. So,

$$\begin{aligned} \lim_{\|h\| \rightarrow \infty} (\|h\| \exp(-\rho \|h\|)) &= \lim_{\|h\| \rightarrow \infty} (\|h\| / \exp(\rho \|h\|)) \\ &= \lim_{\|h\| \rightarrow \infty} (\|h\| / \exp(\rho \|h\|)) = \lim_{\|h\| \rightarrow \infty} (\exp(-\rho \|h\|)) = 0 \end{aligned}$$

then

$$\lim_{\|h\| \rightarrow \infty} \gamma(h) = \nu^2 + \xi^2$$

- The nugget effect is the limiting value of the semi-variogram as $h \rightarrow 0$ from above, hence it is $\gamma(h) \rightarrow \nu^2$ as $h \rightarrow 0^+$.
 - The partial sill is the sill minus the nugget and is hence ξ^2 .
- (3) No, it would be unrealistic because if $\phi > 0$ then the covariance is always positive for infinitely large distances h . In practical terms this means that two points will always be correlated however far apart they are, it would be unrealistic.

(Given as Formative assessment 2)

Exercise 14. (★) Consider we the geostatistical model $(Z_s)_{s \in \mathcal{S}}$ with

$$Z(s) = \mu(s) + w(s) + \varepsilon(s)$$

where $w(s)$ is a weakly stationary process with mean zero and covariogram $c_w(h; \sigma^2, \phi) = \sigma^2 \exp\left(-\frac{1}{\phi} \|h\|\right)$, $\mu(s; \beta)$ is a deterministic function

$$\mu(s; \beta) = \sum_{j=0}^p \psi_j(s) \beta_j = (\psi(s))^\top \beta$$

with unknown coefficients $\beta = (\beta_0, \dots, \beta_p)^\top$ and known basis functions $\psi(s) = (\psi_0(s), \dots, \psi_p(s))^\top$, $\varepsilon(s)$ is a nugget effect process whose covariogram has sill τ^2 , and assume that $w(s)$ and $\varepsilon(s)$ are independent Gaussian Processes.

- (1) Write down the formula of the covariogram $c(h; (\sigma^2, \phi, \tau))$ of (Z_s) .
- (2) Consider a re-parametrization $\theta = (\sigma^2, \phi, \xi)$ where $\xi^2 = \frac{\tau^2}{\sigma^2}$ is called signal to noise ratio. Assume there is available a dataset $\{(s_i, Z_i)\}_{i=1}^n$ where $Z_i := Z(s_i)$ is a realization of $(Z_s)_{s \in \mathcal{S}}$ at site s_i .
 - (a) Let Ψ be a matrix with $[\Psi]_{i,j} = \psi_j(s_i)$. Let D be a matrix such as $[D]_{i,j} = \|s_i - s_j\|$. Consider that you can use convenient notation such as $\exp(D)$ meaning $[\exp(D)]_{i,j} = \exp(D_{i,j})$. Write down the covariance matrix $C(\theta)$ of $Z = (Z_1, \dots, Z_n)^\top$ as a function of D and θ .
 - (b) Write down the log likelihood function $\log(L(Z; \theta))$ of $Z = (Z_1, \dots, Z_n)^\top$ given $\theta = (\sigma^2, \phi, \xi)$.
- (3) Let $r(\cdot)$ (called correlogram) such as $c(\cdot) = \sigma^2 r(\cdot)$. Assume that (ϕ, ξ) as known constants.
 - (a) Compute the likelihood equations¹ w.r.t. (β, σ^2) , and for given (ϕ, ξ) .
 - (b) Compute the MLE $\hat{\beta}_{(\phi, \xi)}$ of β as a function of (ϕ, ξ) .
 - (c) Compute the MLE $\hat{\sigma}_{(\phi, \xi)}^2$ of σ^2 as a function of (ϕ, ξ) .
 - (d) Compute the unbiased estimator of $\tilde{\sigma}^2$ of σ^2 .

Hint: Consider the fitted values $e = (e_1, \dots, e_n)^\top$ as $e = [I - H]Z$ where $H = (\Psi^\top R^{-1} \Psi)^{-1} \Psi^\top R^{-1}$, and write $\hat{\sigma}_{(\phi, \xi)}^2$ w.r.t. e .

Hint: It is given that $E(Z^\top A Z) = E(Z)^\top A E(Z) + \text{tr}(A \text{Var}(Z))$ when $Z \sim \text{Normal}$

- (4) Compute the so-called log “profiled likelihood” $\log(L(Z; (\phi, \xi)))$ resulting as

$$L(Z; (\phi, \xi)) = L\left(Z; \beta = \hat{\beta}_{(\phi, \xi)}, \sigma^2 = \hat{\sigma}_{(\hat{\beta}_{(\phi, \xi)}, \phi, \xi)}^2, \phi, \xi\right)$$

by replacing the β with $\hat{\beta}_{(\phi, \xi)}$ and σ^2 with $\hat{\sigma}_{(\hat{\beta}_{(\phi, \xi)}, \phi, \xi)}^2$ in the actual likelihood $L(Z; \beta, \theta = (\sigma^2, \phi, \xi))$.

Describe how you would compute suitable values $(\hat{\phi}, \hat{\xi})$ for the MLE of (ϕ, ξ)

Solution. It is

- (1) It is

$$\begin{aligned} c(h; (\sigma^2, \phi, \tau)) &= c_\delta(h; \sigma^2, \phi) + c_\varepsilon(h; \tau) \\ &= \sigma^2 \exp\left(-\frac{1}{\phi} \|h\|\right) + \tau 1_{\{0\}}(h) \end{aligned}$$

- (2) It is

- (a)

$$C(\sigma^2, \phi, \xi) = \sigma^2 \exp\left(-\frac{1}{\phi} D\right) + \sigma^2 \xi^2 I$$

¹that is, the gradient of the log-likelihood

(b)

$$\begin{aligned} 2 \log (L (Z ; \theta)) &= 2 \log (\mathcal{N} (Z | \Psi \beta, C (\theta))) \\ &= -n \log (\sigma^2) - \log \left(\left| \exp \left(-\frac{1}{\phi} D \right) + \xi^2 I \right| \right) \\ &\quad - \frac{1}{\sigma^2} (Z - \Psi \beta)^\top \left(\exp \left(-\frac{1}{\phi} D \right) + \xi^2 I \right)^{-1} (Z - \Psi \beta) \end{aligned}$$

(3) It is

$$\begin{aligned} 2 \log (L (Z ; \theta)) &= -n \log (\sigma^2) - \log \left(\left| \exp \left(-\frac{1}{\phi} D \right) + \xi^2 I \right| \right) \\ &= -\frac{1}{\sigma^2} (Z - \Psi \beta)^\top \left(\exp \left(-\frac{1}{\phi} D \right) + \xi^2 I \right)^{-1} (Z - \Psi \beta) \end{aligned}$$

Let $R_{(\phi, \xi)}$ matrix with $[R_{(\phi, \xi)}]_{i, j} = r (s_i - s_j | \phi, \xi)$

(a) So the likelihood equations are $0 = \nabla_{(\beta, \sigma^2)} \log (L (Z ; \theta))$

$$\begin{cases} 0 = \Psi^\top (R_{(\phi, \xi)})^{-1} Z - \Psi^\top (R_{(\phi, \xi)})^{-1} \Psi \beta \\ 0 = \frac{n}{2} \frac{1}{\sigma^2} + \frac{1}{2(\sigma^2)^2} (Z - \Psi \beta)^\top (R_{(\phi, \xi)})^{-1} (Z - \Psi \beta) \end{cases}$$

(b) It is

$$\hat{\beta}_{(\phi, \xi)} = \left(\Psi^\top (R_{(\phi, \xi)})^{-1} \Psi \right)^{-1} \Psi^\top (R_{(\phi, \xi)})^{-1} Z$$

(c) It is

$$\hat{\sigma}_{(\beta, \phi, \xi)} = \frac{1}{n} (Z - \Psi \beta)^\top (R_{(\phi, \xi)})^{-1} (Z - \Psi \beta)$$

and by substituting I get

$$\begin{aligned} \hat{\sigma}_{(\phi, \xi)} &= \hat{\sigma}_{(\hat{\beta}_{(\phi, \xi)}, \phi, \xi)} = \frac{1}{n} \left(Z - \Psi \hat{\beta}_{(\phi, \xi)} \right)^\top (R_{(\phi, \xi)})^{-1} \left(Z - \Psi \hat{\beta}_{(\phi, \xi)} \right) \\ &= \frac{1}{n} \left(Z - \Psi \hat{\beta}_{(\phi, \xi)} \right)^\top (R_{(\phi, \xi)})^{-1} \left(Z - \Psi \hat{\beta}_{(\phi, \xi)} \right) \end{aligned}$$

(d) It is

$$e = Z - \Psi \hat{\beta}_{(\phi, \xi)} = (I - H) Z$$

So

$$\begin{aligned} n \hat{\sigma}_{(\phi, \xi)} &= Z^\top (I - H) (R_{(\phi, \xi)})^{-1} (I - H) Z \\ &= [(I - H) Z]^\top (R_{(\phi, \xi)})^{-1} [(I - H) Z] \\ &= e^\top R_{(\phi, \xi)} e \end{aligned}$$

where

$$\mathbb{E} [e] = 0$$

then

$$\begin{aligned}
\mathbb{E}(n\hat{\sigma}(\phi, \xi)) &= \mathbb{E}\left(Z^\top (I - H) (R_{(\phi, \xi)})^{-1} (I - H) Z\right) \\
&= \mathbb{E}\left[Z^\top (R_{(\phi, \xi)})^{-1} Z\right] = \text{tr}\left((R_{(\phi, \xi)})^{-1} \text{Var}(e)\right) \\
&= \text{tr}\left((R_{(\phi, \xi)})^{-1} \text{Var}((I - H) Z)\right) \\
&= \text{tr}\left((R_{(\phi, \xi)})^{-1} (I - H) \sigma^2 R_{(\phi, \xi)} (I - H)\right) = \sigma^2 \text{tr}\left((R_{(\phi, \xi)})^{-1} (I - H) R_{(\phi, \xi)} (I - H)\right) \\
&= \text{tr}((I - H)) = \sigma^2 (n - p)
\end{aligned}$$

So it is

$$\tilde{\sigma}(\beta, \phi, \xi) = \frac{1}{n - p} (Z - \Psi\beta)^\top (R_{(\phi, \xi)})^{-1} (Z - \Psi\beta)$$

because

$$\mathbb{E}(\tilde{\sigma}(\beta, \phi, \xi)) = \sigma^2$$

(e) It is

$$\hat{\beta}(\phi, \xi) = \left(\Psi^\top (R_{(\phi, \xi)})^{-1} \Psi\right)^{-1} \Psi^\top (R_{(\phi, \xi)})^{-1} Z$$

so it is Normal as a linear combination of normal random variables, with mean

$$\mathbb{E}(\hat{\beta}(\phi, \xi)) = \left(\Psi^\top (R_{(\phi, \xi)})^{-1} \Psi\right)^{-1} \Psi^\top (R_{(\phi, \xi)})^{-1} \mathbb{E}(Z) = \beta$$

and Variance

$$\begin{aligned}
\text{Var}(\hat{\beta}(\phi, \xi)) &= \left(\Psi^\top (R_{(\phi, \xi)})^{-1} \Psi\right)^{-1} \Psi^\top (R_{(\phi, \xi)})^{-1} \text{Var}(Z) \Psi (R_{(\phi, \xi)})^{-1} \Psi \\
&= \left(\Psi^\top (R_{(\phi, \xi)})^{-1} \Psi\right)^{-1} \Psi^\top (R_{(\phi, \xi)})^{-1} \sigma^2 R_{(\phi, \xi)} \Psi (R_{(\phi, \xi)})^{-1} \Psi \\
&= \left(\Psi^\top (R_{(\phi, \xi)})^{-1} \Psi\right)^{-1}
\end{aligned}$$

(4) It is

$$\begin{aligned}
\log(L(Z; (\phi, \xi))) &= L\left(Z; \beta = \hat{\beta}_{(\phi, \xi)}, \sigma^2 = \hat{\sigma}_{(\hat{\beta}_{(\phi, \xi)}, \phi, \xi)}^2, \phi, \xi\right) \\
&\quad - \frac{n}{2} \log\left(\hat{\sigma}_{(\hat{\beta}_{(\phi, \xi)}, \phi, \xi)}^2\right) - \frac{1}{2} \log(|R_{(\phi, \xi)}|)
\end{aligned}$$

where obviously

$$0 = \nabla_{(\phi, \xi)} \log(L(Z; (\phi, \xi)))|_{(\phi, \xi) = (\hat{\phi}, \hat{\xi})}$$

cannot be solved numerically. The Newton method or the gradient descent method can be used to maximize $\log(L(Z; (\phi, \xi)))$.

Exercise 15. (★) Let $(Z_s)_{s \in \mathcal{S}}$ be a specified statistical model. Assume that $(Z_s)_{s \in \mathcal{S}}$ is weakly stationary with unknown constant mean $\mu = \mathbb{E}(Z(s))$ and known covariogram $c(\cdot)$. Assume there is available a dataset $\{(s_i, Z_i := Z(s_i))\}_{i=1}^n$ and assume they are realizations of $(Z_s)_{s \in \mathcal{S}}$. Assume that the matrix C such as $[C]_{i,j} = c(\|s_i - s_j\|)$ has an inverse. Consider the “Kriging” estimator

μ_{KM} of μ as the BLUE (Best Linear Unbiased Estimator)

$$\mu_{\text{KM}} = \sum_{i=1}^n w_i Z(s_i) = w^\top Z,$$

for some unknown $\{w_i\}$ that we need to learn.

- (1) Find sufficient conditions on $w = (w_1, \dots, w_n)$ so that the Kriging estimator μ_{KM} to be unbiased.
- (2) Assume C is invertable. Compute the MSE of μ_{KM} as a function of $w = (w_1, \dots, w_n)$ and C
- (3) Derive the Kriging estimator μ_{KM} of μ as a function of C
- (4) Derive the Kriging standard error as $\sigma_{\text{KM}} = \sqrt{E(\mu_{\text{KM}} - \mu)^2}$ as a function of C

Solution 16. The method is called Kriging the Mean, and hence we denote it as KM.

- (1) It is

$$\mu_{\text{KM}} = \sum_{i=1}^n w_i Z(s_i) = w^\top Z,$$

where $\{w_i\}$ is a set of unknown weights to be learned.

We assume that assume zero systematic error (unbiasness), hence

$$E(\mu_{\text{KM}} - \mu) = E\left(\sum_{i=1}^n w_i Z(s_i) - \mu\right) = \sum_{i=1}^n w_i \overset{=1}{\cancel{E(Z(s_i))}} - \mu$$

which is satisfied given the assumption

$$\sum_{i=1}^n w_i = 1 \iff 1^\top w = 1 \quad (\text{ASSUMPTION})$$

- (2) It is

$$\begin{aligned} E(\mu_{\text{KM}} - \mu)^2 &= E(\mu_{\text{KM}}^2 + \mu^2 - 2\mu_{\text{KM}}\mu) = \sum_{i=1}^n \sum_{j=1}^n w_i w_j E(Z(s_i) Z(s_j)) - \sum_{i=1}^n w_i \overset{=1}{\cancel{\sum_{j=1}^n w_j \mu}} \\ &= \sum_{i=1}^n \sum_{j=1}^n w_i w_j (c(s_i - s_j) - \mu) = \sum_{i=1}^n \sum_{j=1}^n w_i w_j c(s_i - s_j) = w^\top C w \end{aligned}$$

- (3) To learn the unknown weights $\{w_i\}$ we need to solve

$$w^{\text{KM}} = \arg \min_w E(\mu_{\text{KM}} - \mu)^2, \quad \text{subject to } \sum_{i=1}^n w_i = 1$$

The Lagrange function is

$$\begin{aligned} \mathcal{L}(w, \lambda) &= \sum_{i=1}^n \sum_{j=1}^n w_i w_j c(s_i - s_j) - 2\lambda \left(\sum_{i=1}^n w_i - 1 \right) \\ &= w^\top C w - 2\lambda (1^\top w - 1) \end{aligned}$$

The Kriging to mean equations are $0 = \nabla_{w,\lambda} \mathfrak{L}(w, \lambda)$ producing

$$\begin{cases} 0 = 2 \sum_{j=1}^n w_j^{\text{KM}} c(s_i - s_j) - 2\lambda & \forall i = 1, \dots, n \\ 1 = \sum_{i=1}^n w_i^{\text{KM}} \end{cases}$$

$$\begin{cases} 2Cw^{\text{KM}} - 2\lambda 1 = 0 \\ 1^\top w^{\text{KM}} = 1 \end{cases}$$

Given that C^{-1} exists, I multiply by $1^\top C^{-1}$ and I get

$$21^\top C^{-1}Cw - 21^\top C^{-1}\lambda 1 = 0$$

so

$$\lambda = \frac{1}{1^\top C^{-1}1}$$

I substitute and I get

$$w^{\text{KM}} = \frac{C^{-1}1}{1^\top C^{-1}1}$$

So

$$\mu_{\text{KM}} = \left(\frac{C^{-1}1}{1^\top C^{-1}1} \right)^\top Z$$

(4) It is

$$\sigma_{\text{KM}} = \sqrt{\mathbb{E}(\mu_{\text{KM}} - \mu)^2} = \sqrt{\left(\frac{C^{-1}1}{1^\top C^{-1}1} \right)^\top C \frac{C^{-1}1}{1^\top C^{-1}1}} = \frac{1}{\sqrt{1^\top C^{-1}1}}$$

(Given as Formative assessment 2)

Exercise 17. (★) Let $(Z_s)_{s \in \mathcal{S}}$ be a specified statistical model. Assume that process $(Z_s)_{s \in \mathcal{S}}$ has known mean $\mu(s) = \mathbb{E}(Z(s))$ and known covariance function $c(\cdot, \cdot)$. Assume there is available a dataset $\{(s_i, Z_i := Z(s_i))\}_{i=1}^n$. Assume that the matrix C such as $[C]_{i,j} = c(s_i, s_j)$ has an inverse. Consider the “Kriging” estimator μ_{SK} . Consider the “Kriging” estimator $Z_{\text{SK}}(s_0)$ of $Z(s_0)$ at an unseen spatial location s_0 as the BLUE (Best Linear Unbiased Estimator)

$$Z_{\text{SK}}(s_0) = w_{n+1} + \sum_{i=1}^n w_i Z(s_i) = w_{n+1} + w^\top Z,$$

for some unknown $\{w_i\}$ that we need to learn, and $Z = (Z_1, \dots, Z_n)^\top$. Let $w = (w_1, \dots, w_n)^\top$.

- (1) Find sufficient conditions on $w = (w_1, \dots, w_n)^\top$ so that the Kriging estimator $Z_{\text{SK}}(s_0)$ to be unbiased.
- (2) Derive the MSE of $Z_{\text{SK}}(s_0)$ as

$$\mathbb{E}(Z_{\text{SK}}(s_0) - Z(s_0))^2 = w^\top C w + c(s_0, s_0) - 2w^\top C_0$$

where C_0 is a vector such as $[C_0]_i = c(s_0, s_i)$.

- (3) Derive the Kriging estimator of $Z(s_0)$ as

$$Z_{\text{SK}}(s_0) = \mu(s_0) + C_0^\top C^{-1} [Z - \mu(s_{1:n})]$$

where $\mu(s_{1:n})$ is a vector such as $[\mu(s_{1:n})]_i = \mu(s_i)$.

(4) Compute the Kriging standard error $\sigma_{\text{SK}} = \sqrt{\text{E}(Z_{\text{SK}}(s_0) - Z(s_0))^2}$.

Solution. The method is called Simple Kriging, and hence we denote it as SK.

(1) It is

$$Z_{\text{SK}}(s_0) = w_{n+1} + \sum_{i=1}^n w_i Z(s_i) = w_{n+1} + w^\top Z,$$

where $\{w_i\}$ is a set of unknown weights to be learned.

We assume that assume zero systematic error (unbiasness), hence

$$\text{E}(Z_{\text{SK}}(s_0) - Z(s_0)) = \text{E}\left(w_{n+1} + \sum_{i=1}^n w_i Z(s_i) - Z(s_0)\right) = w_{n+1} + \sum_{i=1}^n w_i \mu(s_i) - \mu(s_0)$$

which is satisfied given the assumption

$$w_{n+1} = \mu(s_0) - \sum_{i=1}^n w_i \mu(s_i) \iff w_{n+1} = \mu(s_0) - w^\top \mu(s_{1:n})$$

where $w = (w_1, \dots, w_n)^\top$.

(2) It is

$$\begin{aligned} \text{E}(Z_{\text{SK}}(s_0) - Z(s_0))^2 &= \text{Var}(Z_{\text{SK}}(s_0) - Z(s_0)) = \text{Var}(w_{n+1} + w^\top Z - Z(s_0)) \\ &= \text{Var}(w_{n+1} + w^\top Z) + \text{Var}(Z(s_0)) - 2\text{Cov}(w_{n+1} + w^\top Z, Z(s_0)) \\ &= w^\top C w + c(s_0, s_0) - 2w^\top \text{Cov}(Z, Z(s_0)) \\ &= w^\top C w + c(s_0, s_0) - 2w^\top C_0 \end{aligned}$$

where $C_0 = \text{Cov}(Z, Z(s_0))$, i.e. $[C_0]_j = c(s_j, s_0)$.

(a) To learn the unknown weights $\{w_i\}$ we need to solve

$$w^{\text{SK}} = \arg \min_w \text{E}(Z_{\text{SK}}(s_0) - Z(s_0))^2, \text{ subject to } w_{n+1} = \mu(s_0) - w^\top \mu(s_{1:n})$$

As $\text{E}(\mu_{\text{SK}} - Z(s_0))^2$ does not depend on w_{n+1} we minimize

$$\begin{aligned} 0 &= \nabla_w \text{E}(Z_{\text{SK}}(s_0) - Z(s_0))^2 = \nabla_w \text{Var}(Z_{\text{SK}}(s_0) - Z(s_0)) \\ &= 2Cw - 2C_0 \end{aligned}$$

So I get

$$w_{\text{SK}} = C^{-1}C_0$$

So

$$\begin{aligned} Z_{\text{SK}}(s_0) &= w_{n+1} + C^{-1}C_0 Z \\ &= \mu(s_0) - (C^{-1}C_0)^\top \mu(s_{1:n}) + (C^{-1}C_0)^\top Z \\ &= \mu(s_0) + C_0^\top C^{-1} [Z - \mu(s_{1:n})] \end{aligned}$$

(3) It is

$$\begin{aligned}\sigma_{\text{SK}} &= \sqrt{\text{E}(Z_{\text{SK}}(s_0) - Z(s_0))^2} \\ &= \sqrt{w_{\text{SK}}^\top C w_{\text{SK}} + c(s_0, s_0) - 2w_{\text{SK}}^\top C_0} \\ &= \sqrt{c(s_0, s_0) - C_0^\top C^{-1} C_0}\end{aligned}$$

Exercise 18. (★) Assume a spatial model

$$(1) \quad Z(s) = \mu + \delta(s), \quad s \in \mathcal{S}$$

with unknown mean $\mu \in \mathbb{R}$. Assume a set of n observed realizations $Z_i := Z(s_i)$ of (1) at sites s_i for $i = 1, \dots, n$. Assume that $Z(s)$ is a weak stationary stochastic process with known covariogram $c(\cdot)$. Derive the formula for the Ordinary Kriging predictor $Z_0 := Z(s_0)$ at spatial location s_0 and its kriging variance as function of the covariogram $c(h)$ and not the semi-variogram.

Exercise 19. (★) Let $(Z_s)_{s \in \mathcal{S}}$ be a specified statistical model. Assume that $(Z_s)_{s \in \mathcal{S}}$ is an intrinsic stationary process with unknown constant mean $\mu(s) = \text{E}(Z(s))$ and known semi-variogram $\gamma(\cdot)$. Assume there is available a dataset $\{(s_i, Z_i := Z(s_i))\}_{i=1}^n$. Consider the “Kriging” estimator $Z_{\text{OK}}(s_0)$ of $Z(s_0)$ at any unseen spatial location s_0 as the BLUE (Best Linear Unbiased Estimator)

$$Z_{\text{OK}}(s_0) = w_{n+1} + \sum_{i=1}^n w_i Z(s_i) = w_{n+1} + w^\top Z$$

for some unknown $\{w_i\}$ that we need to learn, and $Z = (Z_1, \dots, Z_n)^\top$. Let $w = (w_1, \dots, w_n)^\top$.

- (1) Find sufficient conditions on $w = (w_1, \dots, w_n)$ so that the Kriging estimator $Z_{\text{OK}}(s_0)$ to be unbiased.
- (2) Derive the MSE of $Z_{\text{OK}}(s_0)$ as

$$\text{E}(Z_{\text{OK}}(s_0) - Z(s_0))^2 = -w^\top \mathbf{\Gamma} w + 2w^\top \boldsymbol{\gamma}_0$$

where $\boldsymbol{\gamma}_0 = (\gamma(s_0 - s_1), \dots, \gamma(s_0 - s_n))^\top$ and $\mathbf{\Gamma}$ with $[\mathbf{\Gamma}]_{i,j} = \gamma(s_i - s_j)$

- (3) Assume $\mathbf{\Gamma}$ is invertible matrix. Derive the Kriging estimator of $Z(s_0)$ as

$$Z_{\text{OK}}(s_0) = \mathbf{\Gamma}^{-1} \left(\boldsymbol{\gamma}_0 + \frac{1 - \mathbf{1}^\top \mathbf{\Gamma}^{-1} \boldsymbol{\gamma}_0}{\mathbf{1}^\top \mathbf{\Gamma}^{-1} \mathbf{1}} \mathbf{1} \right)^\top Z$$

- (4) Derive the Kriging standard error of $Z_{\text{OK}}(s_0)$ as

$$\sigma_{\text{SK}} = \sqrt{\boldsymbol{\gamma}_0^\top \mathbf{\Gamma}^{-1} \boldsymbol{\gamma}_0 - \frac{(1 - \mathbf{1}^\top \mathbf{\Gamma}^{-1} \boldsymbol{\gamma}_0)^2}{\mathbf{1}^\top \mathbf{\Gamma}^{-1} \mathbf{1}}}$$

Solution. The method is called Ordinary Kriging, and hence we denote it as OK.

(1) It is

$$Z_{\text{OK}}(s_0) = w_{n+1} + \sum_{i=1}^n w_i Z(s_i) = w_{n+1} + w^\top Z,$$

where $\{w_i\}$ is a set of unknown weights to be learned.

$$\mathbb{E}(Z_{\text{OK}}(s_0)) = w_{n+1} + \sum_{i=1}^n w_i \mathbb{E}(Z(s_i)) \Leftrightarrow \mu = w_{n+1} + \mu \sum_{i=1}^n w_i$$

Unbiasness is satisfied given the assumption $w_{n+1} = 0$, and

$$\sum_{i=1}^n w_i = 1 \iff 1^\top w = 1 \quad (\text{ASSUMPTION})$$

(2) The MSE of $Z_{\text{OK}}(s_0)$ is

$$\begin{aligned} \text{MSE}(Z_{\text{OK}}(s_0)) &= \mathbb{E}(Z_{\text{OK}}(s_0) - Z(s_0))^2 = \mathbb{E} \left(\sum_{i=1}^n w_i Z(s_i) - \underbrace{\sum_{i=1}^n w_i Z(s_0)}_{=1} \right)^2 \\ &= \mathbb{E} \left(\sum_{i=1}^n w_i (Z(s_i) - Z(s_0)) \right)^2 \\ &= -\mathbb{E} \left(\frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n w_i w_j (Z(s_i) - Z(s_j))^2 - 2 \sum_{i=1}^n \frac{1}{2} w_i (Z(s_i) - Z(s_0))^2 \right) \\ &= -\sum_{i=1}^n w_i \sum_{j=1}^n w_j \frac{1}{2} \mathbb{E} \left((Z(s_i) - Z(s_j))^2 \right) + 2 \sum_{i=1}^n w_i \frac{1}{2} \mathbb{E} \left((Z(s_i) - Z(s_0))^2 \right) \\ &= -\sum_{i=1}^n w_i \sum_{j=1}^n w_j \gamma(s_i - s_j) + 2 \sum_{i=1}^n w_i \gamma(s_i - s_0) \\ &= -w^\top \mathbf{\Gamma} w + 2w^\top \gamma_0 \end{aligned}$$

where $w = (w_1, \dots, w_n)^\top$, $\gamma_0 = (\gamma(s_0 - s_1), \dots, \gamma(s_0 - s_n))^\top$, and $[\mathbf{\Gamma}]_{i,j} = \gamma(s_i - s_j)$.

(3) The Lagrange multiplier function to minimize the MSE under the assumption is

$$\begin{aligned} \mathcal{L}(w, \lambda) &= -\sum_{i=1}^n w_i w_j \gamma(s_i - s_j) + 2 \sum_{i=1}^n w_i \gamma(s_0 - s_i) - \lambda \left(\sum_{i=1}^n w_i - 1 \right) \\ &= -w^\top \mathbf{\Gamma} w + 2w^\top \gamma_0 - \lambda (1^\top w - 1) \end{aligned}$$

The OK system of equations is $0 = \nabla_{(\{w_i\}, \lambda)} L(w, \lambda)|_{(w, \lambda)}$ producing

$$\begin{cases} 0 = -2 \sum_{j=1}^n w_j^{\text{OK}} \gamma(s_i - s_j) + 2\gamma(s_0 - s_i) - \lambda, & i = 1, \dots, n \\ 1 = \sum_{i=1}^n w_i^{\text{OK}} \end{cases} \iff \begin{cases} 0 = -2\mathbf{\Gamma}w_{\text{OK}} + 2\gamma_0 - \lambda_{\text{OK}}1 \\ 1 = 1^\top w_{\text{OK}} \end{cases}$$

Assuming $\mathbf{\Gamma}$ is invertable and multiplying by $1^\top \mathbf{\Gamma}^{-1}$ it is

$$0 = -2\mathbf{\Gamma}w_{\text{OK}} + 2\gamma_0 - \lambda_{\text{OK}}1 \iff$$

$$0 = -2 \cancel{1^\top \mathbf{\Gamma}^{-1} \mathbf{\Gamma} w_{\text{OK}}} \overset{=1}{+} 2 1^\top \mathbf{\Gamma}^{-1} \gamma_0 - 1^\top \mathbf{\Gamma}^{-1} \lambda_{\text{OK}} 1 \iff$$

$$\lambda_{\text{OK}} = 2 \frac{1^\top \mathbf{\Gamma}^{-1} \gamma_0 - 1}{1^\top \mathbf{\Gamma}^{-1} 1}$$

By substitution I get

$$w_{\text{OK}} = \mathbf{\Gamma}^{-1} \left(\gamma_0 + \frac{1 - 1^\top \mathbf{\Gamma}^{-1} \gamma_0}{1^\top \mathbf{\Gamma}^{-1} 1} 1 \right)$$

Hence

$$Z_{\text{OK}}(s_0) = w_{\text{OK}} Z = \mathbf{\Gamma}^{-1} \left(\gamma_0 + \frac{1 - 1^\top \mathbf{\Gamma}^{-1} \gamma_0}{1^\top \mathbf{\Gamma}^{-1} 1} 1 \right) Z$$

(4) It is

$$\begin{aligned} \sigma_{\text{OK}}(s_0) &= \sqrt{\text{MSE}(Z_{\text{OK}}(s_0))} \\ &= \sqrt{-w^\top \mathbf{\Gamma} w + w^\top \gamma_0} \\ &= \sqrt{\gamma_0^\top \mathbf{\Gamma}^{-1} \gamma_0 - \frac{(1 - 1^\top \mathbf{\Gamma}^{-1} \gamma_0)^2}{1^\top \mathbf{\Gamma}^{-1} 1}} \end{aligned}$$

Note regarding the calculations in MSE:

$$\begin{aligned}
\left(\sum_{i=1}^n w_i (Z(s_i) - Z(s_0)) \right)^2 &= \left(\sum_{i=1}^n w_i (Z_i - Z_0) \right)^2 \\
&= \sum_{i=1}^n w_i^2 (Z_i - Z_0)^2 + 2 \sum_{1 \leq i < j \leq n} w_i (Z_i - Z_0) w_j (Z_j - Z_0) \\
&= \sum_{i=1}^n \sum_{j=1}^n w_i w_j (Z_i - Z_0) (Z_j - Z_0) \\
&= 2 \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n w_i w_j (Z_i - Z_0) (Z_j - Z_0) \\
&\quad - \frac{1}{2} \sum_{i=1}^n w_i (Z_i - Z_0)^2 - \frac{1}{2} \sum_{j=1}^n w_j (Z_j - Z_0)^2 \\
&\quad + 2 \frac{1}{2} \sum_{i=1}^n w_i (Z_i - Z_0)^2 \\
&= - \frac{1}{2} \left(\sum_{i=1}^n w_i \sum_{j=1}^n w_j [(Z_i - Z_0)^2 + (Z_j - Z_0)^2 - 2w_i w_j (Z_i - Z_0) (Z_j - Z_0)] \right) \\
&\quad + 2 \frac{1}{2} \sum_{i=1}^n w_i (Z_i - Z_0)^2 \\
&= - \frac{1}{2} \left(\sum_{i=1}^n w_i \sum_{j=1}^n w_j [(Z_i - Z_0) - (Z_j - Z_0)]^2 \right) + 2 \frac{1}{2} \sum_{i=1}^n w_i (Z_i - Z_0)^2 \\
&= - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n w_i w_j (Z_i - Z_j)^2 + 2 \frac{1}{2} \sum_{i=1}^n w_i (Z_i - Z_0)^2 \\
&= - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n w_i w_j (Z(s_i) - Z(s_j))^2 + 2 \frac{1}{2} \sum_{i=1}^n w_i (Z(s_i) - Z(s_0))^2
\end{aligned}$$

(Given as Problem class 3 material)

Exercise 20. (★)

Inventory of useful formulas.

[Normal distr. conditioning] Let $x_1 \in \mathbb{R}^{d_1}$, and $x_2 \in \mathbb{R}^{d_2}$. If

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \sim N_{d_1+d_2} \left(\begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix}, \begin{bmatrix} \Sigma_1 & \Sigma_{21}^\top \\ \Sigma_{21} & \Sigma_2 \end{bmatrix} \right)$$

then it is

$$x_2|x_1 \sim N_{d_2}(\mu_{2|1}, \Sigma_{2|1})$$

where

$$\mu_{2|1} = \mu_2 + \Sigma_{21}\Sigma_1^{-1}(x_1 - \mu_1) \quad \text{and} \quad \Sigma_{2|1} = \Sigma_2 - \Sigma_{21}\Sigma_1^{-1}\Sigma_{21}^\top$$

Consider the Bayesian Kriging from your lecture notes:

$$Z(s) = Y(s) + \varepsilon(s), \quad s \in \mathcal{S}$$

where

$$\varepsilon(\cdot) \sim \text{GP}(0, c_\varepsilon(\cdot, \cdot|\tau))$$

with $c_\varepsilon(s, s'|\tau) = \tau^2 1_{\{0\}}(\|s - s'\|)$ and

$$Y(\cdot)|\beta, \theta \sim \text{GP}(\mu(\cdot|\beta), c_Y(\cdot, \cdot|\sigma^2, \phi))$$

with mean function $\mu(\cdot|\beta)$ (to be specified later) labeled by unknown parameter β and covariance function $c_Y(\cdot, \cdot|\sigma^2, \phi) = \sigma^2 r(\cdot, \cdot|\phi)$.

Assume there is available a dataset $\{(s_i, Z_i)\}_{i=1}^n$ where $Z_i = Z(s_i)$ is a realization of a stochastic process (Z_s) .

- (1) Write the hierarchical spatial model $Z(\cdot)|Y(\cdot), \beta, \varphi$ and $Y(\cdot)|\beta, \varphi$ where $\varphi = (\sigma^2, \phi, \tau)^\top$.
- (2) Write the marginal process $Z(\cdot)|\beta, \varphi$ where $\varphi = (\sigma^2, \phi, \tau)^\top$, its mean function denoted as $\mu(\cdot|\cdot)$, and its covariance function denoted as $c(\cdot|\cdot)$.

Hint:: Let Y and X be independent random variables with $X \sim N(\mu_X, \Sigma_X)$, $Y \sim N(\mu_Y, \Sigma_Y)$. Let A and B be fixed matrices. Let c be a fixed vector. Then

$$AX + BY + c \sim N(A\mu_X + B\mu_Y + c, A\Sigma_X A^\top + B\Sigma_Y B^\top)$$

- (3) Compute the predictive process $Z(\cdot)|Z, \beta, \varphi$ as

$$Z(\cdot)|Z, \beta, \varphi \sim \text{GP}(\mu_1(\cdot|\beta, \varphi), c_1(\cdot, \cdot|\varphi))$$

with

$$\begin{aligned} c_1(s, s'|\varphi) &= c(s, s|\varphi) + (C(S, s|\varphi))^\top (C(S, S|\varphi))^{-1} C(S, s'|\varphi) \\ \mu_1(s|\beta, \varphi) &= \mu(s|\beta) - (C(S, s|\varphi))^\top (C(S, S|\varphi))^{-1} (\mu(S|\beta) - Z) \end{aligned}$$

Hint: See the Conditional Normal formula above.

- (4) Assume $\mu(s|\beta) = \psi(s)^\top \beta$. Consider a conjugate prior on β as $\beta \sim N(b, B)$ where $B > 0$.
 - (a) Write down the Bayesian statistical model involving $[Z|\beta, \varphi]$, and $[\beta|\varphi]$.

(b) Compute the posterior distribution as

$$\beta|Z, \varphi \sim N(b_n(\varphi), B_n(\varphi))$$

with

$$B_n = \left(B^{-1} + \Psi^\top (C(S, S|\varphi))^{-1} \Psi \right)^{-1}$$

$$b_n = B_n \left(B^{-1}b + \Psi^\top (C(S, S|\varphi))^{-1} Z \right)$$

where $R(S, S|\varphi)$ is a matrix with $[R(S, S|\varphi)]_{i,j} = r(s_i, s_j|\varphi)$.

Hint: Use the following identity

$$(y - \Phi\beta)^\top \Sigma^{-1}(y - \Phi\beta) + (\beta - \mu)^\top V^{-1}(\beta - \mu) = (\beta - \mu^*)^\top (V^*)^{-1}(\beta - \mu^*) + S^*;$$

$$V^* = \left(V^{-1} + \Phi^\top \Sigma^{-1} \Phi \right)^{-1}; \quad \mu^* = V^* \left(V^{-1}\mu + \Phi^\top \Sigma^{-1}y \right)$$

$$S^* = \mu^\top V^{-1}\mu - (\mu^*)^\top (V^*)^{-1}(\mu^*) + y^\top \Sigma^{-1}y;$$

(c) Compute the (posterior) predictive process $Z(\cdot)|Z, \varphi$ given the data Z and given the parameters φ as

$$Z(\cdot)|Z, \varphi \sim \text{GP}(\mu_2(\cdot|\varphi), c_2(\cdot, \cdot|\varphi))$$

with

$$\mu_2(s|\varphi) = \left(\Psi C^{-1} (C(s))^\top - \psi(s) \right)^\top \left(B^{-1} + \Psi^\top C^{-1} \Psi \right)^{-1} B^{-1}b$$

$$+ \left[(C(s))^\top + \left(\Psi C^{-1} (C(s))^\top - \psi(s) \right)^\top \left(B^{-1} + \Psi^\top C^{-1} \Psi \right)^{-1} \Psi \right] C^{-1}Z$$

$$c_2(s, s'|\varphi) = c(s, s'|\varphi) - (C(s))^\top C^{-1}C(s')$$

$$+ \left(\Psi C^{-1} (C(s))^\top - \psi(s) \right)^\top \left(B^{-1} + \Psi^\top C^{-1} \Psi \right)^{-1} \left(\Psi C^{-1} (C(s'))^\top - \psi(s') \right)$$

with column vector $C(s) := (c(s, s_1|\varphi), \dots, c(s, s_n|\varphi))^\top$, and matrix $C := C(S, S|\varphi)$.

(d) Compute the marginal likelihood $\Pr(Z|\varphi)$ in the form

$$\Pr(Z|\sigma^2, \varphi) = N \left(Z|\Psi b, \left(C^{-1} - C^{-1}\Psi \left(B^{-1} + \Psi^\top B^{-1}\Psi \right)^{-1} \Psi^\top C^{-1} \right)^{-1} \right)$$

where Ψ is a matrix with $[\Psi]_{i,j} = \psi_j(s_i)$, and R is a matrix with $[C]_{i,j} = c(s_i, s_j|\varphi)$.

Hint-2:: It is

$$\int N(Z|\Psi\beta, C) N(\beta|b, B) d\beta = N(Z|\Psi b, C + \Psi B \Psi^\top)$$

Hint 3:: [Woodbury matrix identity]

$$(A + UCV)^{-1} = A^{-1} - A^{-1}U(C^{-1} + VA^{-1}U)^{-1}VA^{-1}$$

(5) Consider non-informative prior $\Pr(\beta) \propto 1$ for β by specifying $b \rightarrow 0$ and letting $B^{-1} \rightarrow 0$. Argue whether such a prior can be used. Recompute the (asymptotic) quantities $\Pr(Z|\varphi)$, $[Z(\cdot)|Z, \varphi]$ under this new prior.

Solution.

(1) The hierarchical model was

$$\begin{aligned} Z(\cdot) | Y(\cdot), \tau &\sim \text{GP}(Y(\cdot), c_\varepsilon(\cdot, \cdot | \sigma^2, \xi)) \\ Y(\cdot) | \beta, &\sim \text{GP}(\mu(\cdot | \beta), c_Y(\cdot, \cdot | \sigma^2, \phi)) \end{aligned}$$

with $c_\varepsilon(\cdot, \cdot | \sigma^2, \xi) = \sigma^2 \xi^2 1_{\{0\}}(\|s - s'\|)$

(2) We use the additive property of the Gaussian distribution (Hint-1) it is

$$Z(\cdot) | \beta, \xi, \sigma^2, \phi \sim \text{GP}(\mu(\cdot | \beta), c(\cdot, \cdot | \xi, \sigma^2, \phi))$$

where

$$c(s, s' | \xi, \sigma^2, \phi) = c_Y(s, s' | \sigma^2, \phi) + c_\varepsilon(s, s' | \sigma^2, \xi)$$

(3) Assume a vector of “unseen” sites $S_* = (s_{*,1}, \dots, s_{*,q})^\top$ for any $q \in \mathbb{N}_0$. Let convenient notation $Z := Z(S)$, and $Z_* := Z(S_*)$. The joint marginal distribution of $(Z_*, Z)^\top$ given β , $\theta = (\sigma^2, \phi, \tau)^\top$ is

$$(2) \quad \begin{pmatrix} Z_* \\ Z \end{pmatrix} | \beta, \varphi \sim \text{N} \left(\begin{pmatrix} \mu(S_*; \beta) \\ \mu(S; \beta) \end{pmatrix}, \begin{pmatrix} C(S_*, S_* | \varphi) & (C(S_*, S | \varphi))^\top \\ C(S_*, S | \varphi) & C(S, S | \varphi) \end{pmatrix} \right)$$

by using convenient notation $[C(S_*, S | \varphi)]_{i,j} = s(s_{*,i}, s_j | \varphi)$ and $[\mu(S; \beta)]_i = \mu(s_i; \beta)$.

Using the Normal distribution conditioning, I get

$$Z_* | Z, \beta, \varphi \sim \text{N}(\mu_*(S_* | \beta, \varphi), C_*(S_*, S_* | \varphi))$$

where

$$\begin{aligned} C_*(S_*, S_* | \varphi) &= C(S_*, S_* | \varphi) + (C(S, S_* | \varphi))^\top (C(S, S | \varphi))^{-1} C(S, S_* | \varphi) \\ \mu_*(S_* | \beta, \varphi) &= \mu(S_* | \beta) - (C(S, S_* | \varphi))^\top (C(S, S | \varphi))^{-1} (\mu(S | \beta) - Z) \end{aligned}$$

As it is for any length of of any vector S_* , then it is a Gaussian process

$$Z(\cdot) | Z, \beta, \varphi \sim \text{GP}(\mu_1(\cdot | \beta, \varphi), c_1(\cdot, \cdot | \varphi))$$

with

$$\begin{aligned} c_1(s, s' | \varphi) &= c(s, s' | \varphi) + (C(S, s | \varphi))^\top (C(S, S | \varphi))^{-1} C(S, s' | \varphi) \\ \mu_1(s | \beta, \varphi) &= \mu(s | \beta) - (C(S, s | \varphi))^\top (C(S, S | \varphi))^{-1} (\mu(S | \beta) - Z) \end{aligned}$$

(4)

(a) the Bayesian model is

$$(3) \quad \begin{cases} Z | \beta, \varphi \stackrel{\text{ind}}{\sim} \text{N}(\Psi \beta, C(S, S | \varphi)) \\ \beta | \varphi \sim \text{N}(b, B) \end{cases}$$

(b) Let $C := C(S, S|\varphi)$. The posterior distribution is

$$\begin{aligned}
\Pr(\beta|Z, \varphi) &\propto \Pr(Z|\beta, \varphi) \Pr(\beta|\varphi) \\
&= N(Z|\Psi\beta, C) N(\beta|b, B) \\
&\propto \exp\left(-\frac{1}{2}(Z - \Psi\beta)^\top C^{-1}(Z - \Psi\beta)\right) \exp\left(-\frac{1}{2}(\beta - b)^\top B^{-1}(\beta - b)\right) \\
&= \exp\left(-\frac{1}{2}\left[(Z - \Psi\beta)^\top C^{-1}(Z - \Psi\beta) + (\beta - b)^\top B^{-1}(\beta - b)\right]\right)
\end{aligned}$$

By using the hint I have

$$(Z - \Psi\beta)^\top C^{-1}(Z - \Psi\beta) + (\beta - b)^\top B^{-1}(\beta - b) = (\beta - b_n)^\top (B_n)^{-1}(\beta - b_n) + R_n$$

where by denoting $B_n := B_n(\varphi)$, and $b_n := b_n(\varphi)$ I get

$$\begin{aligned}
B_n &= (B^{-1} + \Psi^\top C^{-1} \Psi)^{-1} \\
b_n &= B_n (B^{-1} b + \Psi^\top C^{-1} Z)
\end{aligned}$$

and R_n is a “constant” quantity that does not contain any β . Hence

$$\begin{aligned}
\Pr(\beta|Z, \varphi) &\propto \exp\left(-\frac{1}{2}(\beta - b_n)^\top (B_n)^{-1}(\beta - b_n) - \frac{1}{2}R_n\right) \\
&\propto \exp\left(-\frac{1}{2}(\beta - b_n)^\top (B_n)^{-1}(\beta - b_n)\right)
\end{aligned}$$

Well, from the above, I recognize the kernel of the Multivariate Normal distribution, as

$$\beta|Z, \varphi \sim N(b_n(\varphi), B_n(\varphi))$$

(c) Assume a vector of “unseen” sites $S_* = (s_{*,1}, \dots, s_{*,q})^\top$ for any $q \in \mathbb{N}_0$. Let convenient notation $Z := Z(S)$, and $Z_* := Z(S_*)$. It is

$$\begin{aligned}
\Pr(Z_*|Z, \varphi) &= \int \Pr(Z_*|Z, \varphi) \Pr(\beta|Z, \varphi) d\beta \\
&= \int N(Z_*|\mu_1(S_*), C_1(S_*, S_*)) N(\beta|b_n, B_n) d\beta
\end{aligned}$$

Let $\Psi_* = \Psi(S_*)$, $C_* = C(S_*, S|\varphi)$, and $C_{**} = C(S_*, S_*|\varphi)$. Notice that

$$\begin{aligned}
\mu_1(S_*) &= \Psi_*\beta - C_*C^{-1}(\Psi\beta - Z) \\
&= [\Psi_* - C_*C^{-1}] \beta + C_*C^{-1}Z
\end{aligned}$$

Hence, for given/fixed Z, φ , it is

$$Z_* = C_*C^{-1}Z + [\Psi_* - C_*C^{-1}] \beta + \zeta, \quad \zeta \sim N(0, C_1(S_*, S_*))$$

Hence, because $\beta \sim N(b_n, B_n)$, and because $Z_*|Z, \varphi$ is a linear combination of the Normally distributed random vector $\beta \sim N(b_n, B_n)$, $Z_*|Z, \varphi$ follows a Normal distribution,

with mean

$$\begin{aligned}
\mu_2(S_*) &= \mathbb{E}_{\beta \sim \mathcal{N}(b_n, B_n)} (Z_* | \mu_1(S_*), C_1(S_*, S_*)) \\
&= (\Psi_* - C_* C^{-1}) \mathbb{E}_{\beta \sim \mathcal{N}(b_n, B_n)} (\beta) + C_* C^{-1} Z \\
&= (\Psi_* - C_* C^{-1}) b_n + C_* C^{-1} Z \\
&= (\Psi_* - C_* C^{-1}) \left(B^{-1} + \Psi^\top C^{-1} \Psi \right)^{-1} \left(B^{-1} b + \Psi^\top C^{-1} Z \right) + C_* C^{-1} Z \\
&= (\Psi_* - C_* C^{-1}) \left(B^{-1} + \Psi^\top C^{-1} \Psi \right)^{-1} B^{-1} b \\
&\quad + \left[(\Psi_* - C_* C^{-1}) \left(B^{-1} + \Psi^\top C^{-1} \Psi \right)^{-1} \Psi^\top + C_* \right] C^{-1} Z
\end{aligned}$$

and with covariance matrix

$$\begin{aligned}
C_2(S_*, S_*) &= \text{Var}_{\beta \sim \mathcal{N}(b_n, B_n)} (Z_* | \mu_1(S_*), C_1(S_*, S_*)) \\
&= \text{Var}_{\beta \sim \mathcal{N}(b_n, B_n)} ([\Psi_* - C_* C^{-1}] \beta) + \text{Var}_{\zeta \sim \mathcal{N}(0, C_1(S_*, S_*))} (\zeta) \\
&= [\Psi_* - C_* C^{-1}] B_n [\Psi_* - C_* C^{-1}]^\top + C_1(S_*, S_*) \\
&= [\Psi_* - C_* C^{-1}] \left(B^{-1} + \Psi^\top C^{-1} \Psi \right)^{-1} [\Psi_* - C_* C^{-1}]^\top \\
&\quad + C_{**} + (C_*)^\top C^{-1} C_*
\end{aligned}$$

Since this is for any vector S_* of any length, then

$$Z(\cdot) | Z, \varphi \sim \text{GP}(\mu_2(\cdot | \varphi), c_2(\cdot, \cdot | \varphi))$$

with mean function at s

$$\begin{aligned}
\mu_2(s | \varphi) &= (\psi(s) - C(s) C^{-1}) \left(B^{-1} + \Psi^\top C^{-1} \Psi \right)^{-1} B^{-1} b \\
&\quad + \left[(\psi(s) - C(s) C^{-1}) \left(B^{-1} + \Psi^\top C^{-1} \Psi \right)^{-1} \Psi^\top + C(s) \right] C^{-1} Z
\end{aligned}$$

and covariance function as s, s'

$$\begin{aligned}
c_2(s, s' | \varphi) &= [\psi(s) - C(s) C^{-1}] \left(B^{-1} + \Psi^\top C^{-1} \Psi \right)^{-1} [\psi(s') - C(s') C^{-1}]^\top \\
&\quad + c(s, s' | \varphi) + (C(s))^\top C^{-1} C(s')
\end{aligned}$$

(d) It is, from Hint-2,

$$\begin{aligned}
\Pr(Z | \varphi) &= \int \Pr(Z | \beta, \varphi) \Pr(\beta) d\beta \\
&= \int \mathcal{N}(Z | \Psi \beta, C(S, S | \varphi)) \mathcal{N}(\beta | b, B) d\beta \\
&= \int \mathcal{N}(Z | \Psi \beta, C(S, S | \varphi)) \mathcal{N}(\Psi \beta | \Psi b, \Psi B \Psi^\top) d\beta \\
&= \mathcal{N}(Z | \Psi b, C(S, S | \varphi) + \Psi B \Psi^\top)
\end{aligned}$$

where by letting $C = C(S, S|\varphi)$ and using the Hint I get

$$\left(C + \Psi B \Psi^\top\right)^{-1} = C^{-1} - C^{-1} \Psi \left(B^{-1} + \Psi^\top C^{-1} \Psi\right)^{-1} \Psi^\top C^{-1}$$

(5) Denote $C = C(S, S|\varphi)$. It is

$$\begin{aligned} \lim_{\substack{B^{-1} \rightarrow 0 \\ b \rightarrow 0}} \Pr(Z|\varphi) &= \lim_{\substack{B^{-1} \rightarrow 0 \\ b \rightarrow 0}} \mathcal{N}\left(Z|\Psi b, C + \Psi B \Psi^\top\right) \\ &\propto \lim_{\substack{B^{-1} \rightarrow 0 \\ b \rightarrow 0}} \mathcal{N}\left(Z|0, \left(C^{-1} - C^{-1} \Psi \left(\Psi^\top C^{-1} \Psi\right)^{-1} \Psi^\top C^{-1}\right)^{-1}\right) \\ &< \infty \end{aligned}$$

namely the bottom part of the fraction of the posterior of $\beta|Z, \varphi$ is bounded (finite), which implies a proper posterior. It is

$$\Pr(\beta|Z, \varphi) \propto \exp\left(-\frac{1}{2}(\beta - b_n)^\top B_n^{-1}(\beta - b_n)\right)$$

$$\begin{aligned} \lim_{\substack{B^{-1} \rightarrow 0 \\ b \rightarrow 0}} \exp\left(-\frac{1}{2}(\beta - b_n)^\top B_n^{-1}(\beta - b_n)\right) &= \\ \exp\left(-\frac{1}{2}\left(\beta - \left(\Psi^\top C^{-1} \Psi\right)^{-1} \Psi^\top C^{-1} Z\right)^\top \left(\Psi^\top C^{-1} \Psi\right) \left(\beta - \left(\Psi^\top C^{-1} \Psi\right)^{-1} \Psi^\top C^{-1} Z\right)\right) \end{aligned}$$

hence the limiting case is

$$\beta|Z, \varphi \stackrel{\text{approx}}{\sim} \mathcal{N}\left(\left(\Psi^\top C^{-1} \Psi\right)^{-1} \Psi^\top C^{-1} Z, \left(\Psi^\top C^{-1} \Psi\right)^{-1}\right)$$

The predictive process becomes

$$\begin{aligned} Z(\cdot)|Z, \varphi &\stackrel{\text{approx}}{\sim} \text{GP}(\mu_3(\cdot|\varphi), c_3(\cdot, \cdot|\varphi)) \\ \mu_3(s|\varphi) &= \left[(\psi(s) - C(s)C^{-1})\left(\Psi^\top C^{-1} \Psi\right)^{-1} \Psi^\top + C(s)\right] C^{-1} Z \\ c_3(s, s'|\varphi) &= (\psi(s) - C(s)C^{-1})\left(\Psi^\top C^{-1} \Psi\right)^{-1} (\psi(s') - C(s')C^{-1})^\top \\ &\quad + c(s, s'|\varphi) + (C(s))^\top C^{-1} C(s') \end{aligned}$$

(Given as Problem class 3 material)

Exercise 21. (★) Show that the extension variance $\sigma_E^2(v, V)$ of a small volume v to a larger volume V is obtained by

$$\sigma_E^2(v, V) = 2\bar{\gamma}(v, V) - \bar{\gamma}(v, v) - \bar{\gamma}(V, V)$$

where

$$\bar{\gamma}(v, V) = \frac{1}{|v| |V|} \int_{s \in v} \int_{s' \in V} \gamma(s - s') \, ds \, ds'$$

Solution. Essentially I need to show that that

$$\begin{aligned} \text{Var}(Z(A) - Z(B)) &= \frac{1}{|A| |B|} \int_{x \in A} \int_{y \in B} \gamma(x - y) \, dx \, dy \\ &\quad - \frac{1}{|A| |A|} \int_{x \in A} \int_{y \in A} \gamma(x - y) \, dx \, dy \\ &\quad - \frac{1}{|B| |B|} \int_{x \in B} \int_{y \in B} \gamma(x - y) \, dx \, dy \end{aligned}$$

where I use A, B instead of v, V and x, y instead of s, s' for clarity on notation.

It is

$$\begin{aligned} \text{Var}(Z(A) - Z(B)) &= \text{Cov}(Z(A) - Z(B), Z(A) - Z(B)) \\ &= \text{Cov}(Z(A), Z(A)) + \text{Cov}(Z(B), Z(B)) - 2\text{Cov}(Z(A), Z(B)) \\ &= \frac{1}{|A| |A|} \int_{x \in A} \int_{y \in A} \text{Cov}(Z(x), Z(y)) \, dx \, dy \\ &\quad + \frac{1}{|B| |B|} \int_{x \in B} \int_{y \in B} \text{Cov}(Z(x), Z(y)) \, dx \, dy \\ &\quad - 2 \frac{1}{|A| |B|} \int_{x \in A} \int_{y \in B} \text{Cov}(Z(x), Z(y)) \, dx \, dy \end{aligned}$$

OK, now I need to write all these Cov as γ ; I know that

$$\begin{aligned} \gamma(x - y) &= \frac{1}{2} \text{Var}(Z(x) - Z(y)) \\ &= \frac{1}{2} \text{Var}(Z(x)) + \frac{1}{2} \text{Var}(Z(y)) - \text{Cov}(Z(x), Z(y)) \end{aligned}$$

that is

$$\text{Cov}(Z(x), Z(y)) = \frac{1}{2} \text{Var}(Z(x)) + \frac{1}{2} \text{Var}(Z(y)) - \gamma(x - y)$$

Now I'll gonna put all these in the quantity of interest, one by one

$$\begin{aligned} \frac{1}{|A| |A|} \int_{x \in A} \int_{y \in A} \text{Cov}(Z(x), Z(y)) \, dx \, dy &= \frac{1}{|A| |A|} \int_{x \in A} \int_{y \in A} \frac{1}{2} \text{Var}(Z(x)) \, dx \, dy \\ &\quad + \frac{1}{|A| |A|} \int_{x \in A} \int_{y \in A} \frac{1}{2} \text{Var}(Z(y)) \, dx \, dy \\ &\quad - \frac{1}{|A| |A|} \int_{x \in A} \int_{y \in A} \gamma(x - y) \, dx \, dy \\ &= \frac{1}{|A|} \int_{x \in A} \text{Var}(Z(x)) \, dx \\ &\quad - \frac{1}{|A|^2} \int_{x \in A} \int_{y \in A} \gamma(x - y) \, dx \, dy \end{aligned}$$

and by symmetry

$$\begin{aligned} \frac{1}{|B||B|} \int_{x \in B} \int_{y \in B} \text{Cov}(Z(x), Z(y)) \, dx dy &= \frac{1}{|B|} \int_{x \in B} \text{Var}(Z(x)) \, dx \\ &\quad - \frac{1}{|B|^2} \int_{x \in B} \int_{y \in B} \gamma(x-y) \, dx dy \end{aligned}$$

and finally,

$$\begin{aligned} \frac{1}{|A||B|} \int_{x \in A} \int_{y \in B} \text{Cov}(Z(x), Z(y)) \, dx dy &= \frac{1}{|A||B|} \int_{x \in A} \int_{y \in B} \frac{1}{2} \text{Var}(Z(x)) \, dx dy \\ &\quad + \frac{1}{|A||B|} \int_{x \in A} \int_{y \in B} \frac{1}{2} \text{Var}(Z(y)) \, dx dy \\ &\quad - \frac{1}{|A||B|} \int_{x \in A} \int_{y \in B} \gamma(x-y) \, dx dy \\ &= \frac{1}{2} \frac{1}{|A|} \int_{x \in A} \text{Var}(Z(x)) \, dx \\ &\quad + \frac{1}{2} \frac{1}{|B|} \int_{x \in B} \text{Var}(Z(x)) \, dx \\ &\quad - \frac{1}{|A||B|} \int_{x \in A} \int_{y \in B} \gamma(x-y) \, dx dy \end{aligned}$$

Putting all these together, we get the result.

(Given as Formative assessment 3)

Exercise 22. (★) Suppose a large volume V is partitioned into n smaller units v of equal size. Show that the dispersion variance $\sigma^2(u|V)$ can be written in term of variogram integrals

$$\bar{\gamma}(v, V) = \frac{1}{|v||V|} \int_{s \in v} \int_{s' \in V} \gamma(s-s') \, ds ds'$$

as

$$\sigma^2(v|V) = \bar{\gamma}(V, V) - \bar{\gamma}(v, v)$$

Solution.

(Given as Formative assessment 3)

Exercise 23. (★) Consider a statistical model which is a stochastic process $(Z_s)_{s \in \mathbb{R}}$ (so s has dimension 1), where $Z(\cdot) \sim \text{GP}(\mu(\cdot), c(\cdot, \cdot))$ with mean function $\mu(s) = 1$ and covariance function $c(s, t) = (s-t)^2$. Assume there is available a dataset $\{(Z_i, s_i)\}_{i=1}^n$ where $Z_i = Z(s_i)$ and $s_i \in \mathbb{R}$ are point sites.

- (1) Compute the length $|v|$ of the block $v = [a, b] \subset \mathbb{R}$.
- (2) Compute the block mean $\mu(v)$ for some block $v = [a, b] \subset \mathbb{R}$.
- (3) Compute the block covariance function $c(v, s)$ for some block $v = [a, b] \subset \mathbb{R}$.

- (4) Compute the block covariance function $c(v, v')$ for some blocks $v = [a, b] \subset \mathbb{R}$ and $v' = [a', b'] \subset \mathbb{R}$.
- (5) Denote $Z = (Z_1, \dots, Z_n)^\top$, and $S = \{s_1, \dots, s_n\}$. Let $v = [a, b] \subset \mathbb{R}$ and $v' = [a', b'] \subset \mathbb{R}$ be two intervals. Compute the joint distribution of $(Z(v), Z(v'), Z)^\top$ as a function of $c(\cdot, \cdot)$, S , v , v' , Z , and $\mu(\cdot)$. What is the name of the distribution and what are the parameter functions define them?
- (6) (Bayesian Kriging) Compute the predictive stochastic process at blocks v with $|v| > 0$.

Hint:: Let $x_1 \in \mathbb{R}^{d_1}$, and $x_2 \in \mathbb{R}^{d_2}$. If

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \sim N_{d_1+d_2} \left(\begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix}, \begin{bmatrix} \Sigma_1 & \Sigma_{21}^\top \\ \Sigma_{21} & \Sigma_2 \end{bmatrix} \right)$$

then it is

$$x_2|x_1 \sim N_{d_2}(\mu_{2|1}, \Sigma_{2|1})$$

where

$$\mu_{2|1} = \mu_2 + \Sigma_{21}\Sigma_1^{-1}(x_1 - \mu_1) \quad \text{and} \quad \Sigma_{2|1} = \Sigma_2 - \Sigma_{21}\Sigma_1^{-1}\Sigma_{21}^\top$$

Solution.
