

**PRIME MINISTER'S OFFICE
REGIONAL ADMINISTRATION AND LOCAL GOVERNMENT
ARUSHA REGION**



**FORM ONE TERMINAL EXAMINATION
PHYSICS**

CODE: 031

TIME: 2: 30 HRS

26th May, 2026

INSTRUCTIONS

1. This paper consists of sections A, B and B with a total of the (10) questions.
2. Answer all questions in each section.
3. All answers must be written in the spaces provided
4. All writing must be in blue or black in except drawings which must be in pencil.
5. Write your name at the top of each page
6. Where necessary the following constants may be used
 - (i) Acceleration due to gravity, $g = 10 \text{ m/s}$ or 10 N/kg
 - (ii) Density of water 1000 Kg/m^3 or 1 g/cm^3
 - (iii) $\Pi = 3.14$


GEPAM science hub

WhatsApp 0748978398

SECTION A:(15MARKS)

Answer **all** question from this section


1. For each of the items (i-x) choose the correct answer among the given alternatives and write its answer in the box provided
- (i) How does physics help us in understanding the natural world and the universe?
- A. Physics is not related to the study of natural world and universe
 - B. Physics help us to understand the behavior of matter and energy from subatomic particles to celestial bodies
 - C. Physics only deals with artificial and human made systems
 - D. Physics is not essential for understanding the natural world and the universe
- (ii) Both density and relative density are physical quantities; How do they differ?
- A. Density is the basic quantity while relative density is a derived quantity
 - B. Density has unit but relative density has no units
 - C. Density has no unit while relative density has unit
 - D. Density is derived quantity while relative density is a basic quantity
- (iii) During measurement of the length by using a metre rule, one can get a wrong answer due to wrong position of the eye, an error associated with this situation is called,
- A. Parallax error
 - B. Eye error
 - C. Zero error
 - D. metre rule error
- (iv) The mass of an object is increased if its
- A. State is changed
 - B. Surface is increased
 - C. Volume is reduced
 - D. Amount of matter is increased
- (v) A person measures the length, width, height and mass of a rectangular metal block. Which of these measurements must be used in order to calculate the density of the metal?
- A. Mass and volume
 - B. Height and mass only
 - C. Length, width and height only
 - D. Length, width, height and mass
- (vi) The ball in the figure 1 below have the same size which one will hit the ground first if they are dropped at the same time and the same height?



Wooden ball



Plastic ball



Metal ball

- A. Metal first, then wood followed by plastic ball
 B. Wood first, then plastic followed by metal ball
 C. They will hit the ground at the same time
 D. They will; hit the ground at different interval for the same acceleration due to gravity.
- (vii) The same body is immersed in liquid A then in liquid B. The extent which the body sinks in liquid B is less than in liquid A. What conclusion can be drawn from such an observation? The density of
- A. Liquid A is more than the density in liquid B
 B. The liquid B is more than that of liquid A
 C. The solid body is less than that of the liquid in both cases
 D. The liquid A is the same as that of liquid B
- (viii) A student wants to compare the densities of two different liquids. Which instrument should the student use for a direct comparison?
- A. Barometer
 B. Thermometer
 C. Hygrometer
 D. Hydrometer
- (ix) A body is pulling a box by a force of 100N, if the frictional force experienced by the box is 20N, what is the amount of causing the motion of the box?
- A. 100 N
 B. 120 N
 C. 80 N
 D. 5 N
- (x) If the relative density of a substance is 11.3, the density of a substance is
- A. 11.3 Kg/m³
 B. 11.3g/cm³
 C. 11.3×10⁻³g/m³
 D. 11.3×10³g/cm³
2. match each of the following physical quantities in List A with its corresponding units in List B by writing the letter of the correct answer below an item number in a table provided

LIST A	LIST B
(i) Area	A: Newton per square meter
(ii) Volume	B: has no unit
(iii) Weight	C: Newton
(iv) Density	D: length x width
(v) pressure	E: Kilogram per cubic meter
	F: cubic meter
	G: Mass and accelerations due to gravity
	H: Square meter

ANSWERS

Item number	i	ii	Iii	iv	v
Answer					

SECTION B: (70 Marks)

Answer **all** questions

3. (a) In the process of measuring the diameter of a ball bearing by using a micrometer screw gauge with a least count of 0.01mm, the sleeve shows that it has 1.6cm, and the thimble indicate 46 divisions. What is the actual diameter of the ball bearing?

- (b) (i) what are the significant of the two parties in measurement?

(a)

(b)

- (ii) With the aid of diagrams write down the procedures which must be followed during measurement of volume of irregular objects by using measuring cylinder

- (iv) Why a triple beam balance is so named?

.....
.....

4. (a)(i) the force of friction between the surface of a floor and the surface of a box in contact with the floor is 200N. John wishes to move the box on the floor with the constant velocity. How much force has to be applied on the box?

(ii) If a box has a mass of 100kg, calculate the coefficient of the friction force.

(b) Why car dashboards are fixed with air bags?

.....
.....

5. (a) (i) Explain how each of the following quantities differ from each other

(a) Density and relative density

.....
.....
.....
.....
.....

(b) Mass and Weight

.....
.....
.....
.....
.....

(ii) The density of water is 1000kg/m^3 . What does this imply?

.....
.....
.....

(b) In our daily life the application of physics is very important particularly in density; Mention five (5) application of density

- (i)
- (ii)
- (iii)
- (iv)
- (v)

6. (a) (i) energy can neither be created nor destroyed but transformed from one form to another form of energy, what is the name given to a device used to convert one form of energy to another form?

(ii) what are the energy changes when two cars are colliding head to on each other?

.....
.....
.....
.....
.....
.....

(b) Draw a diagram showing a simple pendulum swing freely from point A to point C through point B and show the energy changes in each point.

7. (a) A sample of milk diluted with water has a density of 1040 Kg/m^3 . If a pure milk has a density of 1080 Kg/m^3 . Find the percentage of water by volume in the diluted milk.

.....
.....
.....
.....
.....

-
.....
- (b)(i) A student place a hollow plastic ball and solid plastic ball of the same size into water. The hollow ball floats and the solid ball sinks. Explains the difference in their behavior using the Archimedes' Principle and the concept of density.

-
.....
.....
- (ii) State the Archimedes' Principle

-
.....
.....
8. (a) During experiment in laboratory or measurement in a school laboratory, someone may obtain an answer which is somehow deviate from the current answer due to errors. Mention four (4) reasons for arise of errors;

- (i)
(ii)
(iii)
(iv)

- (b) Rectangular block measures length 4.00cm, width 2.5cm and height 1.009cm.

- (i) Name the instrument used to measure the sides of the rectangular block

- (ii) Calculate the volume of the rectangular block

- (iii) If the mass of the rectangular block is 300g, find the relative density of the block.

9. (a) Laboratory is a special room or building designed specifically for carrying scientific experiments, therefore a laboratory must have good features for scientific experiments, mention five (5) features of a good laboratory.

(i)
(ii)
(iii)
(iv)
(v)

- (b)(i) What is the difference between first Aid and first Aid kit

.....

- (ii)Mention four importance of first Aid

[Faint blue watermark reading "AMM"]

SECTION C (15 Marks)

10. (a) During experiment of determine the relative density of ethanal in the laboratory, John has got the following data during the observation

Mass of empty bottle = 40g

Mass of empty bottle plus mass of a sand = 90g

Mass of a sand plus mass of an ethanal is 120g

If the bottle used has a capacity of 600.0cm^3 . find the following

- Mass of ethanal in kg
- density of a bottle used
- relative density of an ethanal
- which is denser between sand and ethanal? Why?

[The page contains horizontal dotted lines for writing.]

(b) A ship is sailing in sea enters a wide and deep river. The density of sea water is 1100kg/m^3 .

Determine a comparison of volume of object when it is in sea water to when it is in river water.

REGIONAL TERMINAL EXAMINATION

PRIME MINISTER'S OFFICE
REGIONAL ADMINISTRATION AND LOCAL GOVERNMENT
ARUSHA REGION FORM ONE TERMINAL EXAMINATION
031 PHYSICS

MARKING GUIDE: - 2026.

1:

i	ii	iii	iv	v	vi	vii	viii	ix	x
B	B	A	D	D	C	B	D	C	B

2:

①② = 10 marks

i	ii	iii	iv	v
H	F	C	E	A

3a

Data

Least count = 0.01 mm




Main scale = 1.6 cm
= 16 mm

Circular scale = 46 divisions
= $46 \times 0.01 \text{ mm}$
= 0.46 mm

Total reading = main scale + circular scale
= 16 mm + 0.46 mm
= 16.46 mm

∴ Diameter of a ball bearing is 16.46 mm.

- (b) (i) @ Number part, it tells about the magnitude of what to be measured.
(b) unit part, it tells about what is to be measured.
(ii) Consider the measuring cylinders below



GEPAM science hub

WhatsApp 0748978398

The volume of irregular object AB is determined by following the procedures below

- i) Fill an empty measuring cylinder with a liquid to a certain level (V_1)
- ii) Carefully measure the initial volume of liquid and name it as V_1
- iii) Gently lower the irregular object into a liquid as shown in figure above
- iv) Read the final volume of liquid and name it V_2
- v) Find the difference in volume

$$\text{difference} = V_2 - V_1 = \text{volume of an irregular object. } (0.02\frac{1}{2}) = 0.02\frac{1}{2}$$

- 3b) (ii) because it contains three beams $(0.01\frac{1}{2})$
100g graduations, 10g graduations, 1g graduation

- 4a) (i) Data:
frictional force = 200N
John must apply a force of magnitude greater or equal to the frictional force, therefore he must apply a force of 200N (0.2)

- (ii) Mass = 100kg
Required, Coefficient of frictional force (μ)
from:

$$\mu = \frac{F_f}{R} \text{ but } R = W = mg.$$
$$= 100\text{kg} \times 10\text{N/kg} \quad (0.3)$$
$$= 1000\text{N}$$

$$\mu = \frac{200\text{N}}{1000\text{N}} \quad (0.2)$$
$$= 0.2$$

$\therefore$ The coefficient of frictional force is 0.2N. (0.1)

- 4b) Cars are fixed with airbags on their dash board in order to reduce the effects of force toward the driver when an accident occurs (0.2)

(1) (a) Density is the mass of an object divide by volume of an object while Relative density is the ratio of density of object to density of water (02)
→ Density has SI-unit while relative density has no SI-unit.

(b) Mass is measured in kg while Weight is measured in Newton. (Any difference) (02)

(ii) The density of water is 1000 kg/m^3 . This means that a capacity of 1 m^3 can hold an amount of 1000 kg . Therefore for every 1 m^3 there are 1000 kg of a substance (water). (01)


5(b) Applications of density

- (i) used in identification of gemstones eg: diamond, gold.
- ii) used by geologists to determine mineral content
- iii) used in selection of building materials
- iv) used during designing of swimming and diving equipment
- v) used in designing Marine vehicles. 01 mark @ = 05

6(a) (i) Transducer (01)

(ii) Kinetic energy is converted to → Heat energy
→ Light energy 01 @ = 03
→ sound energy

6(b) A diagram of simple pendulum.



At point A there is maximum potential energy
At point B there is maximum kinetic energy
At point C there is maximum potential energy. (03)
At any point along AB ~ BC, the energy is conserved, $P.E + K.E = \text{Constant}$.

Q7a

Data.

Density of milk = 1080 kg/m^3 .

Density of diluted milk = 1040 kg/m^3

Required, Percentage of water in milk.

Mass of milk = M_m , volume of milk = V_m

Mass of water = M_w , volume of water = V_w

Density of mixture = $\frac{M_w + M_m}{V_w + V_m} = 1040 \text{ kg/m}^3$

$$\frac{M_w + M_m}{V_w + V_m} = 1040 \quad \text{--- (i)} \quad (01)$$

but

$$\frac{M_m}{V_m} = 1080 \quad (01)$$

$$M_m = 1080 V_m \quad \text{--- (ii)}$$

Subst. (ii) into (i)

$$\frac{1080 V_m + M_w}{V_m + V_w} = 1040 \quad (01)$$

$$1080 V_m + M_w = 1040 V_m + 1040 V_w$$

$$\text{but } M_w = 1000 V_w \quad (01)$$

$$1080 V_m + 1000 V_w = 1040 V_m + 1040 V_w$$

$$1080 V_m - 1040 V_m = 1040 V_w - 1000 V_w \quad (01)$$

$$40 V_m = 40 V_w$$

$$V_m = V_w = \text{volume of milk} = \text{volume of water.}$$

$$\text{total volume} = V_m + V_w = 2 V_m.$$

$$\begin{aligned} \text{Percentage of milk} &= \frac{V_m}{2 V_m} \times 100\%, \quad \frac{V_w}{2 V_w} \times 100\% \\ &= \frac{1}{2} \times 100\%, \quad \frac{1}{2} \times 100\% \quad (01) \\ &= 50\%, \quad 50\%. \end{aligned}$$

$\therefore$ - The volume percentage of water is 50%.

- (i) A hollow ball has less density than solid ball
 - A hollow ball has less weight than the upthrust compared to the weight of a solid ball. (02)

(ii) Any object when it is partially or totally immersed in a fluid it experiences an upthrust which is equal to the weight of the fluid displaced by the object. (02)

Q2 Reasons for arise of errors

- (i) instrument fault during manufacture.
- (ii) Damage during use (04)
- (iii) Poor storage of an instrument.
- (iv) Human factors (parallax error, zero error, instrument error).

(b) Data:

$$\text{Length} = 4.00 \text{ cm}$$

$$\text{Width} = 2.5 \text{ cm}$$

$$\text{Height} = 1.00 \text{ cm}$$

(i) The instrument used is meter rule (01)

$$\begin{aligned} \text{(ii) volume} &= L \times W \times H \\ &= 4.00 \text{ cm} \times 2.5 \text{ cm} \times 1.00 \text{ cm} \\ &= 10 \text{ cm}^3 \\ \text{volume is } 10 \text{ cm}^3 & \quad (01) \end{aligned}$$

(iii) Relative density of

$$\text{Density of rectangular block} = \frac{\text{Mass of block}}{\text{volume of block}} \quad (01)$$

$$\begin{aligned} \rho_b &= \frac{300 \text{ g}}{10 \text{ cm}^3} \\ &= 30 \text{ g/cm}^3 \quad (01) \end{aligned}$$

$$\begin{aligned} \text{Relative density} &= \frac{\rho_b}{\rho_w} \\ &= \frac{30 \text{ g/cm}^3}{1 \text{ g/cm}^3} \\ &= 30 \quad (02) \end{aligned}$$

$\therefore$ Relative density of rectangular block is 30

9@ A good laboratory must have the following features.

- (i) Physics Laboratory should be well ventilated and its doors should be open outwards.
- (ii) Fire extinguisher should be fitted in accessible positions.
- (iii) Laboratory floors should not be polished.
- (iv) First aid kit should be present.
- (v) Cabinets and drawers should be included during designing Laboratory room. (05)

9@ First Aid kit is a small box containing items that are used to give help to a sick or injured person while First Aid - is the first and immediate assistance or care given to a sick or injured person before getting a Professional Medical help. (05) (03)

- (i) $\rightarrow$ saves life
 $\rightarrow$ prevent victim's condition from becoming worse
 $\rightarrow$ it promotes hope and encouragement to a victim
 $\rightarrow$ it helps to reduce pain and suffering
 $\rightarrow$ it prevents infections. (05) (05) (00.5) = 2 marks

10@ Data:

$$M_e = 40g$$

$$M_e + M_s = 90g$$

$$M_s + M_{\text{ethano}} = 120g$$

$$90g - 40g = \text{Mass of sand} \quad (03)$$

$$M_s = 50g$$

$$M_{\text{ethano}} = 120g - 50g \\ = 70g.$$

(i) Mass of ethano = $70g = 0.07kg$.

(ii) Density of bottle = $\frac{\text{Mass of bottle}}{\text{Volume of bottle}} \quad (01)$

$$= \frac{40g}{600cm^3}$$
$$= 0.067g/cm^3$$

$\therefore$ Density of bottle = $0.067g/cm^3 \quad (01)$

iii) Relative density of an ethanol

$$\text{Density of ethanol} = \frac{\text{Mass of ethanol}}{\text{volume of bottle}} \quad (01)$$

$$= \frac{70g}{600\text{cm}^3}$$

$$= 0.12\text{g/cm}^3$$

$$\text{Relative density} = \frac{\text{Density of ethanol}}{\text{Density of water}} \quad (01)$$

$$= \frac{0.12\text{g/cm}^3}{4\text{g/cm}^3}$$

$$= 0.12$$

$\therefore$ Relative density of ethanol is 0.12. (01)

(iv) Comparison of densities of sand and ethanol.

$$\text{Density of sand} = \frac{\text{Mass of sand}}{\text{volume of bottle.}}$$

$$= \frac{50g}{600\text{cm}^3}$$

$$= 0.08\text{g/cm}^3 \quad (01)$$

$$\text{Density of ethanol} = 0.12\text{g/cm}^3 > 0.08\text{g/cm}^3$$

$\therefore$ Ethanol is more denser than sand. (01)

Ans)

Data:

$$\text{Density of sea water } (\rho_s) = 1100\text{kg/m}^3$$

$$\text{Volume of ship in sea water} = V_s$$

$$\text{Density of ship river water} = 1000\text{kg/m}^3 = \rho_r$$

$$\text{volume of ship in river} = V_r$$

Mass of ship is the same everywhere ($M_s = M_r$).

$$M_s = \rho_s V_s$$

$$M_r = \rho_r V_r$$

$$\text{But } M_s = M_r \quad (01)$$

$$f_s v_s = f_R v_R$$

Required: f_s ; f_R , v_s/v_R :

$$\frac{f_s}{f_R} = \frac{v_R}{v_s} \quad (01)$$

Not =

$$\frac{v_s}{v_R} = \frac{f_R}{f_s} \quad (01)$$

$$= \frac{1000 \text{ kg/m}^3}{1100 \text{ kg/m}^3}$$

$$= \frac{10}{11} \quad (01)$$

$$v_s : v_R = 10 : 11 \quad (01)$$