

# **Fundamentals of Machine Learning**


Yassine Laguel

Mail : [yassine.laguel@univ-cotedazur.fr](mailto:yassine.laguel@univ-cotedazur.fr)

# Limitations of linear models

- Linear methods are great for linear tasks

- In linear classification




$x \mapsto w^\top x + b$  linearly separates the two classes.

# Limitations of linear models


- Linear methods are great for linear tasks

- In linear classification



$x \mapsto w^\top x + b$  linearly separates the two classes.

- In linear regression




$x \mapsto w^\top x + b$  approximates the continuous response  $y$ .

# Limitations of linear models

- Linear methods are great for linear tasks


- In linear classification




$x \mapsto w^\top x + b$  linearly separates the two classes.

- What can we do when for non-linear tasks ?

- In  $\mathbb{R}^2$  no hyperplane separates the following two classes.



- In linear regression




$x \mapsto w^\top x + b$  approximates the continuous response  $y$ .

# Limitations of linear models


- Linear methods are great for linear tasks

- In linear classification



$x \mapsto w^\top x + b$  linearly separates the two classes.


- In linear regression



$x \mapsto w^\top x + b$  approximates the continuous response  $y$ .


- What can we do when for non-linear tasks ?

- In  $\mathbb{R}^2$  no hyperplane separates the following two classes.



- Yet, if we apply the transformation  $x \mapsto \varphi(x) = (x_1^2, \sqrt{2}x_1x_2, x_2^2)^\top$ .

We obtain linearly separable data points in  $\mathbb{R}^3$ .



# Positive Definite Kernels

## ■ A First definition

### Definition : Positive definite Kernel

A positive definite kernel on a set  $\mathcal{X}$  is a function  $K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$

which satisfies

$$(\text{symmetry}) \text{ (i)} \quad K(x, x') = K(x', x), \quad \forall x, x' \in \mathcal{X}^2$$

$$(\text{positive-definiteness}) \text{ (ii)} \quad \sum_{i=1}^N \sum_{j=1}^N a_i a_j K(\mathbf{x}_i, \mathbf{x}_j) \geq 0 \quad \forall (x_1, \dots, x_n) \in \mathcal{X}^n \\ \forall (a_1, \dots, a_n) \in \mathbb{R}^n$$

# Positive Definite Kernels

## ■ A First definition

### Definition : Positive definite Kernel

A positive definite kernel on a set  $\mathcal{X}$  is a function  $K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$

which satisfies

$$(\text{symmetry}) \text{ (i)} \quad K(x, x') = K(x', x), \quad \forall x, x' \in \mathcal{X}^2$$

$$(\text{positive-definiteness}) \text{ (ii)} \quad \sum_{i=1}^N \sum_{j=1}^N a_i a_j K(\mathbf{x}_i, \mathbf{x}_j) \geq 0 \quad \forall (x_1, \dots, x_n) \in \mathcal{X}^n \\ \forall (a_1, \dots, a_n) \in \mathbb{R}^n$$

### Remarks :

- In other words,  $K$  is a p.d. kernel if and only if, for any  $N \in \mathbb{N}$ , and any set of points  $(x_1, \dots, x_n) \in \mathcal{X}^N$ , the similarity matrix  $\mathbf{K}_{i,j} = K(x_i, x_j)$  is positive definite.

# Positive Definite Kernels

## ■ A First definition

### Definition : Positive definite Kernel

A positive definite kernel on a set  $\mathcal{X}$  is a function  $K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$

which satisfies

$$(\text{symmetry}) \text{ (i)} \quad K(x, x') = K(x', x), \quad \forall x, x' \in \mathcal{X}^2$$

$$(\text{positive-definiteness}) \text{ (ii)} \quad \sum_{i=1}^N \sum_{j=1}^N a_i a_j K(\mathbf{x}_i, \mathbf{x}_j) \geq 0 \quad \forall (x_1, \dots, x_n) \in \mathcal{X}^n \\ \forall (a_1, \dots, a_n) \in \mathbb{R}^n$$

### Remarks :

- In other words,  $K$  is a p.d. kernel if and only if, for any  $N \in \mathbb{N}$ , and any set of points  $(x_1, \dots, x_n) \in \mathcal{X}^N$ , the similarity matrix  $\mathbf{K}_{i,j} = K(x_i, x_j)$  is positive definite.
- We call **kernel methods**, algorithm that apply kernel transformations to input data.

# Positive Definite Kernels

## ■ A First definition

### Definition : Positive definite Kernel

A positive definite kernel on a set  $\mathcal{X}$  is a function  $K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  which satisfies

$$(\text{symmetry}) \text{ (i)} \quad K(x, x') = K(x', x), \quad \forall x, x' \in \mathcal{X}^2$$

$$(\text{positive-definiteness}) \text{ (ii)} \quad \sum_{i=1}^N \sum_{j=1}^N a_i a_j K(\mathbf{x}_i, \mathbf{x}_j) \geq 0 \quad \forall (x_1, \dots, x_n) \in \mathcal{X}^n \\ \forall (a_1, \dots, a_n) \in \mathbb{R}^n$$

### Remarks :

- In other words,  $K$  is a p.d. kernel if and only if, for any  $N \in \mathbb{N}$ , and any set of points  $(x_1, \dots, x_n) \in \mathcal{X}^N$ , the similarity matrix  $\mathbf{K}_{i,j} = K(x_i, x_j)$  is positive definite.
- We call **kernel methods**, algorithm that apply kernel transformations to input data.

## ■ Stability properties

### Proposition : Stability through elementary operations

Let  $K_1, K_2$  be two positive definite kernels. Let  $c > 0$ . Then

(i)  $K_1 + K_2$  is a positive definite kernel.

(ii)  $c \cdot K_1$  is a positive definite kernel.

(iii)  $K_1 \cdot K_2$  is a positive definite kernel.

**Proof :** on the blackboard.

# Positive Definite Kernels

## ■ A First definition

### Definition : Positive definite Kernel

A positive definite kernel on a set  $\mathcal{X}$  is a function  $K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  which satisfies

$$(\text{symmetry}) \text{ (i)} \quad K(x, x') = K(x', x), \quad \forall x, x' \in \mathcal{X}^2$$

$$(\text{positive-definiteness}) \text{ (ii)} \quad \sum_{i=1}^N \sum_{j=1}^N a_i a_j K(\mathbf{x}_i, \mathbf{x}_j) \geq 0 \quad \forall (x_1, \dots, x_n) \in \mathcal{X}^n \\ \forall (a_1, \dots, a_n) \in \mathbb{R}^n$$

### Remarks :

- In other words,  $K$  is a p.d. kernel if and only if, for any  $N \in \mathbb{N}$ , and any set of points  $(x_1, \dots, x_n) \in \mathcal{X}^N$ , the similarity matrix  $\mathbf{K}_{i,j} = K(x_i, x_j)$  is positive definite.
- We call **kernel methods**, algorithm that apply kernel transformations to input data.

## ■ Stability properties

### Proposition : Stability through elementary operations

Let  $K_1, K_2$  be two positive definite kernels. Let  $c > 0$ . Then

(i)  $K_1 + K_2$  is a positive definite kernel.

(ii)  $c \cdot K_1$  is a positive definite kernel.

(iii)  $K_1 \cdot K_2$  is a positive definite kernel.

**Proof :** on the blackboard.

### Proposition : Pointwise limit of p.d. kernels

Let  $(K_i)_{i \geq 0}$  be a sequence of p.d. kernels that converges pointwise to a function  $K$ . Then  $K$  is also a p.d. kernel.

**Proof :** on the blackboard.

# Positive Definite Kernels

## ■ A First definition

### Definition : Positive definite Kernel

A positive definite kernel on a set  $\mathcal{X}$  is a function  $K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  which satisfies

$$(\text{symmetry}) \text{ (i)} \quad K(x, x') = K(x', x), \quad \forall x, x' \in \mathcal{X}^2$$

$$(\text{positive-definiteness}) \text{ (ii)} \quad \sum_{i=1}^N \sum_{j=1}^N a_i a_j K(\mathbf{x}_i, \mathbf{x}_j) \geq 0 \quad \forall (x_1, \dots, x_n) \in \mathcal{X}^n \\ \forall (a_1, \dots, a_n) \in \mathbb{R}^n$$

### Remarks :

- In other words,  $K$  is a p.d. kernel if and only if, for any  $N \in \mathbb{N}$ , and any set of points  $(x_1, \dots, x_n) \in \mathcal{X}^N$ , the similarity matrix  $\mathbf{K}_{i,j} = K(x_i, x_j)$  is positive definite.
- We call **kernel methods**, algorithm that apply kernel transformations to input data.

## ■ Stability properties

### Proposition : Stability through elementary operations

Let  $K_1, K_2$  be two positive definite kernels. Let  $c > 0$ . Then

(i)  $K_1 + K_2$  is a positive definite kernel.

(ii)  $c \cdot K_1$  is a positive definite kernel.

(iii)  $K_1 \cdot K_2$  is a positive definite kernel.

**Proof :** on the blackboard.

### Proposition : Pointwise limit of p.d. kernels

Let  $(K_i)_{i \geq 0}$  be a sequence of p.d. kernels that converges pointwise to a function  $K$ . Then  $K$  is also a p.d. kernel.

**Proof :** on the blackboard.

### Remark :

- We deduce from the above properties that if  $K$  is a p.d. kernel, then
  - $K^n$  is a p.d. kernel for all  $n \in \mathbb{N}, n \geq 1$ .

# Positive Definite Kernels

## ■ A First definition

### Definition : Positive definite Kernel

A positive definite kernel on a set  $\mathcal{X}$  is a function  $K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  which satisfies

$$(\text{symmetry}) \text{ (i)} \quad K(x, x') = K(x', x), \quad \forall x, x' \in \mathcal{X}^2$$

$$(\text{positive-definiteness}) \text{ (ii)} \quad \sum_{i=1}^N \sum_{j=1}^N a_i a_j K(\mathbf{x}_i, \mathbf{x}_j) \geq 0 \quad \forall (x_1, \dots, x_n) \in \mathcal{X}^n \\ \forall (a_1, \dots, a_n) \in \mathbb{R}^n$$

### Remarks :

- In other words,  $K$  is a p.d. kernel if and only if, for any  $N \in \mathbb{N}$ , and any set of points  $(x_1, \dots, x_n) \in \mathcal{X}^N$ , the similarity matrix  $\mathbf{K}_{i,j} = K(x_i, x_j)$  is positive definite.
- We call **kernel methods**, algorithm that apply kernel transformations to input data.

## ■ Stability properties

### Proposition : Stability through elementary operations

Let  $K_1, K_2$  be two positive definite kernels. Let  $c > 0$ . Then

(i)  $K_1 + K_2$  is a positive definite kernel.

(ii)  $c \cdot K_1$  is a positive definite kernel.

(iii)  $K_1 \cdot K_2$  is a positive definite kernel.

**Proof :** on the blackboard.

### Proposition : Pointwise limit of p.d. kernels

Let  $(K_i)_{i \geq 0}$  be a sequence of p.d. kernels that converges pointwise to a function  $K$ . Then  $K$  is also a p.d. kernel.

**Proof :** on the blackboard.

### Remark :

- We deduce from the above properties that if  $K$  is a p.d. kernel, then
  - $K^n$  is a p.d. kernel for all  $n \in \mathbb{N}, n \geq 1$ .
  - $e^K$  is a p.d. kernel.

# **Implementation**



# Reproducing Kernel Hilbert space

## ■ Kernels as inner products

### Definition : Reproducing Kernel Hilbert space

Let  $\mathcal{X}$  be a set and  $\mathcal{H} \subset \mathbb{R}^{\mathcal{X}}$  be a class of functions forming a (real) Hilbert space with inner product  $\langle ., . \rangle_{\mathcal{H}}$ .

The function  $K : \mathcal{X}^2 \mapsto \mathbb{R}$  is called a **reproducing kernel** of  $\mathcal{H}$  if

- (i)  $\mathcal{H}$  contains all functions of the form  $K_x : u \mapsto K(x, u)$ , for  $x \in \mathcal{X}$ .
- (ii) For every  $x \in \mathcal{X}$  and  $f \in \mathcal{H}$ , we have  $f(x) = \langle f, K_x \rangle_{\mathcal{H}}$ .

# Reproducing Kernel Hilbert space

## ■ Kernels as inner products

### Definition : Reproducing Kernel Hilbert space

Let  $\mathcal{X}$  be a set and  $\mathcal{H} \subset \mathbb{R}^{\mathcal{X}}$  be a class of functions forming a (real) Hilbert space with inner product  $\langle ., . \rangle_{\mathcal{H}}$ .

The function  $K : \mathcal{X}^2 \mapsto \mathbb{R}$  is called a **reproducing kernel** of  $\mathcal{H}$  if

- (i)  $\mathcal{H}$  contains all functions of the form  $K_x : u \mapsto K(x, u)$ , for  $x \in \mathcal{X}$ .
- (ii) For every  $x \in \mathcal{X}$  and  $f \in \mathcal{H}$ , we have  $f(x) = \langle f, K_x \rangle_{\mathcal{H}}$ .

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.

# Reproducing Kernel Hilbert space

## ■ Kernels as inner products

### Definition : Reproducing Kernel Hilbert space

Let  $\mathcal{X}$  be a set and  $\mathcal{H} \subset \mathbb{R}^{\mathcal{X}}$  be a class of functions forming a (real) Hilbert space with inner product  $\langle ., . \rangle_{\mathcal{H}}$ .

The function  $K : \mathcal{X}^2 \mapsto \mathbb{R}$  is called a **reproducing kernel** of  $\mathcal{H}$  if

- (i)  $\mathcal{H}$  contains all functions of the form  $K_x : u \mapsto K(x, u)$ , for  $x \in \mathcal{X}$ .
- (ii) For every  $x \in \mathcal{X}$  and  $f \in \mathcal{H}$ , we have  $f(x) = \langle f, K_x \rangle_{\mathcal{H}}$ .

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- Property (ii) is often called the reproducing property.

# Reproducing Kernel Hilbert space

## ■ Kernels as inner products

### Definition : Reproducing Kernel Hilbert space

Let  $\mathcal{X}$  be a set and  $\mathcal{H} \subset \mathbb{R}^{\mathcal{X}}$  be a class of functions forming a (real) Hilbert space with inner product  $\langle ., . \rangle_{\mathcal{H}}$ .

The function  $K : \mathcal{X}^2 \mapsto \mathbb{R}$  is called a **reproducing kernel** of  $\mathcal{H}$  if

- (i)  $\mathcal{H}$  contains all functions of the form  $K_x : u \mapsto K(x, u)$ , for  $x \in \mathcal{X}$ .
- (ii) For every  $x \in \mathcal{X}$  and  $f \in \mathcal{H}$ , we have  $f(x) = \langle f, K_x \rangle_{\mathcal{H}}$ .

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- Property (ii) is often called the reproducing property.
- For  $\mathcal{H}$  given, if a reproducing kernel exists, then  $\mathcal{H}$  is called a **reproducing kernel Hilbert space**.

# Reproducing Kernel Hilbert space

## ■ Kernels as inner products

### Definition : Reproducing Kernel Hilbert space

Let  $\mathcal{X}$  be a set and  $\mathcal{H} \subset \mathbb{R}^{\mathcal{X}}$  be a class of functions forming a (real) Hilbert space with inner product  $\langle ., . \rangle_{\mathcal{H}}$ .

The function  $K : \mathcal{X}^2 \mapsto \mathbb{R}$  is called a **reproducing kernel** of  $\mathcal{H}$  if

- (i)  $\mathcal{H}$  contains all functions of the form  $K_x : u \mapsto K(x, u)$ , for  $x \in \mathcal{X}$ .
- (ii) For every  $x \in \mathcal{X}$  and  $f \in \mathcal{H}$ , we have  $f(x) = \langle f, K_x \rangle_{\mathcal{H}}$ .

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- Property (ii) is often called the reproducing property.
- For  $\mathcal{H}$  given, if a reproducing kernel exists, then  $\mathcal{H}$  is called a **reproducing kernel Hilbert space**.

### Proposition : Reproducing kernels are p.d. kernels

Reproducing kernels are positive definite kernels.

# Reproducing Kernel Hilbert space

## ■ Kernels as inner products

### Definition : Reproducing Kernel Hilbert space

Let  $\mathcal{X}$  be a set and  $\mathcal{H} \subset \mathbb{R}^{\mathcal{X}}$  be a class of functions forming a (real) Hilbert space with inner product  $\langle ., . \rangle_{\mathcal{H}}$ .

The function  $K : \mathcal{X}^2 \mapsto \mathbb{R}$  is called a **reproducing kernel** of  $\mathcal{H}$  if

- (i)  $\mathcal{H}$  contains all functions of the form  $K_x : u \mapsto K(x, u)$ , for  $x \in \mathcal{X}$ .
- (ii) For every  $x \in \mathcal{X}$  and  $f \in \mathcal{H}$ , we have  $f(x) = \langle f, K_x \rangle_{\mathcal{H}}$ .

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- Property (ii) is often called the reproducing property.
- For  $\mathcal{H}$  given, if a reproducing kernel exists, then  $\mathcal{H}$  is called a **reproducing kernel Hilbert space**.

### Proposition : Reproducing kernels are p.d. kernels

Reproducing kernels are positive definite kernels.

**Proof :** On the black board

# Reproducing Kernel Hilbert space

## ■ Kernels as inner products

### Definition : Reproducing Kernel Hilbert space

Let  $\mathcal{X}$  be a set and  $\mathcal{H} \subset \mathbb{R}^{\mathcal{X}}$  be a class of functions forming a (real) Hilbert space with inner product  $\langle ., . \rangle_{\mathcal{H}}$ .

The function  $K : \mathcal{X}^2 \mapsto \mathbb{R}$  is called a **reproducing kernel** of  $\mathcal{H}$  if

- (i)  $\mathcal{H}$  contains all functions of the form  $K_x : u \mapsto K(x, u)$ , for  $x \in \mathcal{X}$ .
- (ii) For every  $x \in \mathcal{X}$  and  $f \in \mathcal{H}$ , we have  $f(x) = \langle f, K_x \rangle_{\mathcal{H}}$ .

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- Property (ii) is often called the reproducing property.
- For  $\mathcal{H}$  given, if a reproducing kernel exists, then  $\mathcal{H}$  is called a **reproducing kernel Hilbert space**.

### Proposition : Reproducing kernels are p.d. kernels

Reproducing kernels are positive definite kernels.

### Proof : On the black board

## ■ Unicity results for the kernel and the RKHS

### Proposition : Continuity. of the evaluation map

A Hilbert space of functions  $H \subset \mathbb{R}^{\mathcal{X}}$  is a RKHS if and only if for any  $x \in \mathcal{X}$ ,  $f \mapsto f(x)$  from  $H$  to  $\mathbb{R}$  is continuous.

# Reproducing Kernel Hilbert space

## ■ Kernels as inner products

### Definition : Reproducing Kernel Hilbert space

Let  $\mathcal{X}$  be a set and  $\mathcal{H} \subset \mathbb{R}^{\mathcal{X}}$  be a class of functions forming a (real) Hilbert space with inner product  $\langle ., . \rangle_{\mathcal{H}}$ .

The function  $K : \mathcal{X}^2 \mapsto \mathbb{R}$  is called a **reproducing kernel** of  $\mathcal{H}$  if

- (i)  $\mathcal{H}$  contains all functions of the form  $K_x : u \mapsto K(x, u)$ , for  $x \in \mathcal{X}$ .
- (ii) For every  $x \in \mathcal{X}$  and  $f \in \mathcal{H}$ , we have  $f(x) = \langle f, K_x \rangle_{\mathcal{H}}$ .

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- Property (ii) is often called the reproducing property.
- For  $\mathcal{H}$  given, if a reproducing kernel exists, then  $\mathcal{H}$  is called a **reproducing kernel Hilbert space**.

### Proposition : Reproducing kernels are p.d. kernels

Reproducing kernels are positive definite kernels.

### Proof : On the black board

## ■ Unicity results for the kernel and the RKHS

### Proposition : Continuity. of the evaluation map

A Hilbert space of functions  $H \subset \mathbb{R}^{\mathcal{X}}$  is a RKHS if and only if for any  $x \in \mathcal{X}$ ,  $f \mapsto f(x)$  from  $H$  to  $\mathbb{R}$  is continuous.

### Proof : On the blackboard.

# Reproducing Kernel Hilbert space

## ■ Kernels as inner products

### Definition : Reproducing Kernel Hilbert space

Let  $\mathcal{X}$  be a set and  $\mathcal{H} \subset \mathbb{R}^{\mathcal{X}}$  be a class of functions forming a (real) Hilbert space with inner product  $\langle ., . \rangle_{\mathcal{H}}$ .

The function  $K : \mathcal{X}^2 \mapsto \mathbb{R}$  is called a **reproducing kernel** of  $\mathcal{H}$  if

- (i)  $\mathcal{H}$  contains all functions of the form  $K_x : u \mapsto K(x, u)$ , for  $x \in \mathcal{X}$ .
- (ii) For every  $x \in \mathcal{X}$  and  $f \in \mathcal{H}$ , we have  $f(x) = \langle f, K_x \rangle_{\mathcal{H}}$ .

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- Property (ii) is often called the reproducing property.
- For  $\mathcal{H}$  given, if a reproducing kernel exists, then  $\mathcal{H}$  is called a **reproducing kernel Hilbert space**.

### Proposition : Reproducing kernels are p.d. kernels

Reproducing kernels are positive definite kernels.

**Proof :** On the black board

## ■ Unicity results for the kernel and the RKHS

### Proposition : Continuity of the evaluation map

A Hilbert space of functions  $H \subset \mathbb{R}^{\mathcal{X}}$  is a RKHS if and only if for any  $x \in \mathcal{X}$ ,  $f \mapsto f(x)$  from  $H$  to  $\mathbb{R}$  is continuous.

**Proof :** On the blackboard.

### Proposition : Unicity of the reproducing kernel

If two reproducing kernel exist for a Hilbert space, then they are equal.

# Reproducing Kernel Hilbert space

## ■ Kernels as inner products

### Definition : Reproducing Kernel Hilbert space

Let  $\mathcal{X}$  be a set and  $\mathcal{H} \subset \mathbb{R}^{\mathcal{X}}$  be a class of functions forming a (real) Hilbert space with inner product  $\langle \cdot, \cdot \rangle_{\mathcal{H}}$ .

The function  $K : \mathcal{X}^2 \mapsto \mathbb{R}$  is called a **reproducing kernel** of  $\mathcal{H}$  if

- (i)  $\mathcal{H}$  contains all functions of the form  $K_x : u \mapsto K(x, u)$ , for  $x \in \mathcal{X}$ .
- (ii) For every  $x \in \mathcal{X}$  and  $f \in \mathcal{H}$ , we have  $f(x) = \langle f, K_x \rangle_{\mathcal{H}}$ .

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- Property (ii) is often called the reproducing property.
- For  $\mathcal{H}$  given, if a reproducing kernel exists, then  $\mathcal{H}$  is called a **reproducing kernel Hilbert space**.

### Proposition : Reproducing kernels are p.d. kernels

Reproducing kernels are positive definite kernels.

**Proof :** On the black board

## ■ Unicity results for the kernel and the RKHS

### Proposition : Continuity of the evaluation map

A Hilbert space of functions  $H \subset \mathbb{R}^{\mathcal{X}}$  is a RKHS if and only if for any  $x \in \mathcal{X}$ ,  $f \mapsto f(x)$  from  $H$  to  $\mathbb{R}$  is continuous.

**Proof :** On the blackboard.

### Proposition : Unicity of the reproducing kernel

If two reproducing kernel exist for a Hilbert space, then they are equal.

**Proof :** On the blackboard.

# Reproducing Kernel Hilbert space

## ■ Kernels as inner products

### Definition : Reproducing Kernel Hilbert space

Let  $\mathcal{X}$  be a set and  $\mathcal{H} \subset \mathbb{R}^{\mathcal{X}}$  be a class of functions forming a (real) Hilbert space with inner product  $\langle \cdot, \cdot \rangle_{\mathcal{H}}$ .

The function  $K : \mathcal{X}^2 \mapsto \mathbb{R}$  is called a **reproducing kernel** of  $\mathcal{H}$  if

- (i)  $\mathcal{H}$  contains all functions of the form  $K_x : u \mapsto K(x, u)$ , for  $x \in \mathcal{X}$ .
- (ii) For every  $x \in \mathcal{X}$  and  $f \in \mathcal{H}$ , we have  $f(x) = \langle f, K_x \rangle_{\mathcal{H}}$ .

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- Property (ii) is often called the reproducing property.
- For  $\mathcal{H}$  given, if a reproducing kernel exists, then  $\mathcal{H}$  is called a **reproducing kernel Hilbert space**.

### Proposition : Reproducing kernels are p.d. kernels

Reproducing kernels are positive definite kernels.

**Proof :** On the black board

## ■ Unicity results for the kernel and the RKHS

### Proposition : Continuity. of the evaluation map

A Hilbert space of functions  $H \subset \mathbb{R}^{\mathcal{X}}$  is a RKHS if and only if for any  $x \in \mathcal{X}$ ,  $f \mapsto f(x)$  from  $H$  to  $\mathbb{R}$  is continuous.

**Proof :** On the blackboard.

### Proposition : Unicity of the reproducing kernel

If two reproducing kernel exist for a Hilbert space, then they are equal.

**Proof :** On the blackboard.

### Proposition : Unicity of the. RKHS

If two RKHS have the same reproducing kernel, then they are equal.

# Reproducing Kernel Hilbert space

## ■ Kernels as inner products

### Definition : Reproducing Kernel Hilbert space

Let  $\mathcal{X}$  be a set and  $\mathcal{H} \subset \mathbb{R}^{\mathcal{X}}$  be a class of functions forming a (real) Hilbert space with inner product  $\langle \cdot, \cdot \rangle_{\mathcal{H}}$ .

The function  $K : \mathcal{X}^2 \mapsto \mathbb{R}$  is called a **reproducing kernel** of  $\mathcal{H}$  if

- (i)  $\mathcal{H}$  contains all functions of the form  $K_x : u \mapsto K(x, u)$ , for  $x \in \mathcal{X}$ .
- (ii) For every  $x \in \mathcal{X}$  and  $f \in \mathcal{H}$ , we have  $f(x) = \langle f, K_x \rangle_{\mathcal{H}}$ .

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- Property (ii) is often called the reproducing property.
- For  $\mathcal{H}$  given, if a reproducing kernel exists, then  $\mathcal{H}$  is called a **reproducing kernel Hilbert space**.

### Proposition : Reproducing kernels are p.d. kernels

Reproducing kernels are positive definite kernels.

**Proof :** On the black board

## ■ Unicity results for the kernel and the RKHS

### Proposition : Continuity. of the evaluation map

A Hilbert space of functions  $H \subset \mathbb{R}^{\mathcal{X}}$  is a RKHS if and only if for any  $x \in \mathcal{X}$ ,  $f \mapsto f(x)$  from  $H$  to  $\mathbb{R}$  is continuous.

**Proof :** On the blackboard.

### Proposition : Unicity of the reproducing kernel

If two reproducing kernel exist for a Hilbert space, then they are equal.

**Proof :** On the blackboard.

### Proposition : Unicity of the. RKHS

If two RKHS have the same reproducing kernel, then they are equal.

**Proof :** On the blackboard.

# Correspondence property and representation theorem

## ■ Aronszajn theorem

**Proposition : Aronszajn theorem**

$K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  is a p.d. kernel on  $\mathcal{X}$  if and only if there exists a Hilbert space  $\mathcal{H}$  and a mapping  $\varphi : \mathcal{X} \rightarrow \mathcal{H}$  such that

$$K(\mathbf{x}, \mathbf{x}') = \langle \Phi(\mathbf{x}), \Phi(\mathbf{x}') \rangle_{\mathcal{H}}, \quad \forall x, x' \in \mathcal{X}.$$

# Correspondence property and representation theorem

## ■ Aronszajn theorem

**Proposition : Aronszajn theorem**

$K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  is a p.d. kernel on  $\mathcal{X}$  if and only if there exists a Hilbert space  $\mathcal{H}$  and a mapping  $\varphi : \mathcal{X} \rightarrow \mathcal{H}$  such that

$$K(\mathbf{x}, \mathbf{x}') = \langle \Phi(\mathbf{x}), \Phi(\mathbf{x}') \rangle_{\mathcal{H}}, \quad \forall x, x' \in \mathcal{X}.$$

**Remarks :**

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.

# Correspondence property and representation theorem

## ■ Aronszajn theorem

**Proposition : Aronszajn theorem**

$K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  is a p.d. kernel on  $\mathcal{X}$  if and only if there exists a Hilbert space  $\mathcal{H}$  and a mapping  $\varphi : \mathcal{X} \rightarrow \mathcal{H}$  such that

$$K(\mathbf{x}, \mathbf{x}') = \langle \Phi(\mathbf{x}), \Phi(\mathbf{x}') \rangle_{\mathcal{H}}, \quad \forall x, x' \in \mathcal{X}.$$

**Remarks :**

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- In other words, the kernel operates a measure of similarity on data, made explicit as an inner product after application of  $\varphi$ .

# Correspondence property and representation theorem

## ■ Aronszajn theorem

### Proposition : Aronszajn theorem

$K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  is a p.d. kernel on  $\mathcal{X}$  if and only if there exists a Hilbert space  $\mathcal{H}$  and a mapping  $\varphi : \mathcal{X} \rightarrow \mathcal{H}$  such that

$$K(\mathbf{x}, \mathbf{x}') = \langle \Phi(\mathbf{x}), \Phi(\mathbf{x}') \rangle_{\mathcal{H}}, \quad \forall x, x' \in \mathcal{X}.$$

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- In other words, the kernel operates a measure of similarity on data, made explicit as an inner product after application of  $\varphi$ .
- With a good choice for  $\varphi$ , one may expect to transform their nonlinear prediction task into a linear one in the space  $\mathcal{H}$ .

# Correspondence property and representation theorem

## ■ Aronszajn theorem

### Proposition : Aronszajn theorem

$K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  is a p.d. kernel on  $\mathcal{X}$  if and only if there exists a Hilbert space  $\mathcal{H}$  and a mapping  $\varphi : \mathcal{X} \rightarrow \mathcal{H}$  such that

$$K(\mathbf{x}, \mathbf{x}') = \langle \Phi(\mathbf{x}), \Phi(\mathbf{x}') \rangle_{\mathcal{H}}, \quad \forall x, x' \in \mathcal{X}.$$

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- In other words, the kernel operates a measure of similarity on data, made explicit as an inner product after application of  $\varphi$ .
- With a good choice for  $\varphi$ , one may expect to transform their nonlinear prediction task into a linear one in the space  $\mathcal{H}$ .
- Any algorithm that solely involves dot products on  $\mathcal{X} \times \mathcal{X}$  may be generalized at a cheap cost by replacing the dot product with a p.d. kernel: this procedure is often called the **kernel trick**.

# Correspondence property and representation theorem

## ■ Aronszajn theorem

**Proposition : Aronszajn theorem**

$K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  is a p.d. kernel on  $\mathcal{X}$  if and only if there exists a Hilbert space  $\mathcal{H}$  and a mapping  $\varphi : \mathcal{X} \rightarrow \mathcal{H}$  such that

$$K(\mathbf{x}, \mathbf{x}') = \langle \Phi(\mathbf{x}), \Phi(\mathbf{x}') \rangle_{\mathcal{H}}, \quad \forall x, x' \in \mathcal{X}.$$

**Remarks :**

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- In other words, the kernel operates a measure of similarity on data, made explicit as an inner product after application of  $\varphi$ .
- With a good choice for  $\varphi$ , one may expect to transform their nonlinear prediction task into a linear one in the space  $\mathcal{H}$ .
- Any algorithm that solely involves dot products on  $\mathcal{X} \times \mathcal{X}$  may be generalized at a cheap cost by replacing the dot product with a p.d. kernel: this procedure is often called the **kernel trick**.

**Proof :** On the black board

# Correspondence property and representation theorem

## ■ Aronszajn theorem

### Proposition : Aronszajn theorem

$K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  is a p.d. kernel on  $\mathcal{X}$  if and only if there exists a Hilbert space  $\mathcal{H}$  and a mapping  $\varphi : \mathcal{X} \rightarrow \mathcal{H}$  such that

$$K(\mathbf{x}, \mathbf{x}') = \langle \Phi(\mathbf{x}), \Phi(\mathbf{x}') \rangle_{\mathcal{H}}, \quad \forall \mathbf{x}, \mathbf{x}' \in \mathcal{X}.$$

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- In other words, the kernel operates a measure of similarity on data, made explicit as an inner product after application of  $\varphi$ .
- With a good choice for  $\varphi$ , one may expect to transform their nonlinear prediction task into a linear one in the space  $\mathcal{H}$ .
- Any algorithm that solely involves dot products on  $\mathcal{X} \times \mathcal{X}$  may be generalized at a cheap cost by replacing the dot product with a p.d. kernel: this procedure is often called the **kernel trick**.

**Proof :** On the black board

## ■ Representation theorem

### Proposition : Representation theorem

Let  $\mathcal{X}$  be a set with a p.d. kernel  $K$ ,  $\mathcal{H}$  be the corresponding RKHS, and  $S = \{\mathbf{x}_1, \dots, \mathbf{x}_n\} \subset \mathcal{X}$  a finite set of points in  $\mathcal{X}$ .

Let  $\psi : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  be a function of  $n + 1$  variables strictly increasing w.r.t. the last variable. Then any solution of

$$\min_{f \in \mathcal{H}} \psi(f(\mathbf{x}_1), \dots, f(\mathbf{x}_n), \|f\|_{\mathcal{H}})$$

admits a representation of the form :

$$f(\mathbf{x}) = \sum_{i=1}^n \alpha_i K(\mathbf{x}_i, \mathbf{x}) = \sum_{i=1}^n \alpha_i K_{\mathbf{x}_i}(\mathbf{x}), \quad \forall \mathbf{x} \in \mathcal{X}.$$

# Correspondence property and representation theorem

## ■ Aronszajn theorem

### Proposition : Aronszajn theorem

$K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  is a p.d. kernel on  $\mathcal{X}$  if and only if there exists a Hilbert space  $\mathcal{H}$  and a mapping  $\varphi : \mathcal{X} \rightarrow \mathcal{H}$  such that

$$K(\mathbf{x}, \mathbf{x}') = \langle \Phi(\mathbf{x}), \Phi(\mathbf{x}') \rangle_{\mathcal{H}}, \quad \forall \mathbf{x}, \mathbf{x}' \in \mathcal{X}.$$

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- In other words, the kernel operates a measure of similarity on data, made explicit as an inner product after application of  $\varphi$ .
- With a good choice for  $\varphi$ , one may expect to transform their nonlinear prediction task into a linear one in the space  $\mathcal{H}$ .
- Any algorithm that solely involves dot products on  $\mathcal{X} \times \mathcal{X}$  may be generalized at a cheap cost by replacing the dot product with a p.d. kernel: this procedure is often called the **kernel trick**.

**Proof :** On the black board

## ■ Representation theorem

### Proposition : Representation theorem

Let  $\mathcal{X}$  be a set with a p.d. kernel  $K$ ,  $\mathcal{H}$  be the corresponding RKHS, and  $S = \{x_1, \dots, x_n\} \subset \mathcal{X}$  a finite set of points in  $\mathcal{X}$ .

Let  $\psi : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  be a function of  $n + 1$  variables strictly increasing w.r.t. the last variable. Then any solution of

$$\min_{f \in \mathcal{H}} \psi(f(\mathbf{x}_1), \dots, f(\mathbf{x}_n), \|f\|_{\mathcal{H}})$$

admits a representation of the form :

$$f(\mathbf{x}) = \sum_{i=1}^n \alpha_i K(\mathbf{x}_i, \mathbf{x}) = \sum_{i=1}^n \alpha_i K_{\mathbf{x}_i}(\mathbf{x}), \quad \forall \mathbf{x} \in \mathcal{X}.$$

### Remarks :

- In other words, the solution lives in the finite dimensional space  $\text{Span}(K_{x_1}, \dots, K_{x_n})$ .

# Correspondence property and representation theorem

## ■ Aronszajn theorem

### Proposition : Aronszajn theorem

$K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  is a p.d. kernel on  $\mathcal{X}$  if and only if there exists a Hilbert space  $\mathcal{H}$  and a mapping  $\varphi : \mathcal{X} \rightarrow \mathcal{H}$  such that

$$K(\mathbf{x}, \mathbf{x}') = \langle \Phi(\mathbf{x}), \Phi(\mathbf{x}') \rangle_{\mathcal{H}}, \quad \forall \mathbf{x}, \mathbf{x}' \in \mathcal{X}.$$

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- In other words, the kernel operates a measure of similarity on data, made explicit as an inner product after application of  $\varphi$ .
- With a good choice for  $\varphi$ , one may expect to transform their nonlinear prediction task into a linear one in the space  $\mathcal{H}$ .
- Any algorithm that solely involves dot products on  $\mathcal{X} \times \mathcal{X}$  may be generalized at a cheap cost by replacing the dot product with a p.d. kernel: this procedure is often called the **kernel trick**.

**Proof :** On the black board

## ■ Representation theorem

### Proposition : Representation theorem

Let  $\mathcal{X}$  be a set with a p.d. kernel  $K$ ,  $\mathcal{H}$  be the corresponding RKHS, and  $S = \{x_1, \dots, x_n\} \subset \mathcal{X}$  a finite set of points in  $\mathcal{X}$ .

Let  $\psi : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  be a function of  $n + 1$  variables strictly increasing w.r.t. the last variable. Then any solution of

$$\min_{f \in \mathcal{H}} \psi(f(x_1), \dots, f(x_n), \|f\|_{\mathcal{H}})$$

admits a representation of the form :

$$f(\mathbf{x}) = \sum_{i=1}^n \alpha_i K(\mathbf{x}_i, \mathbf{x}) = \sum_{i=1}^n \alpha_i K_{\mathbf{x}_i}(\mathbf{x}), \quad \forall \mathbf{x} \in \mathcal{X}.$$

### Remarks :

- In other words, the solution lives in the finite dimensional space  $\text{Span}(K_{x_1}, \dots, K_{x_n})$ .
- From a numerical point of view, this is a very powerful property!

# Correspondence property and representation theorem

## ■ Aronszajn theorem

### Proposition : Aronszajn theorem

$K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  is a p.d. kernel on  $\mathcal{X}$  if and only if there exists a Hilbert space  $\mathcal{H}$  and a mapping  $\varphi : \mathcal{X} \rightarrow \mathcal{H}$  such that

$$K(\mathbf{x}, \mathbf{x}') = \langle \Phi(\mathbf{x}), \Phi(\mathbf{x}') \rangle_{\mathcal{H}}, \quad \forall \mathbf{x}, \mathbf{x}' \in \mathcal{X}.$$

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- In other words, the kernel operates a measure of similarity on data, made explicit as an inner product after application of  $\varphi$ .
- With a good choice for  $\varphi$ , one may expect to transform their nonlinear prediction task into a linear one in the space  $\mathcal{H}$ .
- Any algorithm that solely involves dot products on  $\mathcal{X} \times \mathcal{X}$  may be generalized at a cheap cost by replacing the dot product with a p.d. kernel: this procedure is often called the **kernel trick**.

Proof : On the black board

## ■ Representation theorem

### Proposition : Representation theorem

Let  $\mathcal{X}$  be a set with a p.d. kernel  $K$ ,  $\mathcal{H}$  be the corresponding RKHS, and  $S = \{\mathbf{x}_1, \dots, \mathbf{x}_n\} \subset \mathcal{X}$  a finite set of points in  $\mathcal{X}$ .

Let  $\psi : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  be a function of  $n + 1$  variables strictly increasing w.r.t. the last variable. Then any solution of

$$\min_{f \in \mathcal{H}} \psi(f(\mathbf{x}_1), \dots, f(\mathbf{x}_n), \|f\|_{\mathcal{H}})$$

admits a representation of the form :

$$f(\mathbf{x}) = \sum_{i=1}^n \alpha_i K(\mathbf{x}_i, \mathbf{x}) = \sum_{i=1}^n \alpha_i K_{\mathbf{x}_i}(\mathbf{x}), \quad \forall \mathbf{x} \in \mathcal{X}.$$

### Remarks :

- In other words, the solution lives in the finite dimensional space  $\text{Span}(K_{\mathbf{x}_1}, \dots, K_{\mathbf{x}_n})$ .
- From a numerical point of view, this is a very powerful property!
- A typical form for  $\psi$  is :

$$\psi(f(\mathbf{x}_1), \dots, f(\mathbf{x}_n), \|f\|_{\mathcal{H}}) = c(f(\mathbf{x}_1), \dots, f(\mathbf{x}_n)) + \lambda \Omega(\|f\|_{\mathcal{H}})$$

**Enforces fit to data  
w.r.t. the kernel geometry**

**Regularizer**

# Correspondence property and representation theorem

## ■ Aronszajn theorem

### Proposition : Aronszajn theorem

$K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  is a p.d. kernel on  $\mathcal{X}$  if and only if there exists a Hilbert space  $\mathcal{H}$  and a mapping  $\varphi : \mathcal{X} \rightarrow \mathcal{H}$  such that

$$K(\mathbf{x}, \mathbf{x}') = \langle \Phi(\mathbf{x}), \Phi(\mathbf{x}') \rangle_{\mathcal{H}}, \quad \forall \mathbf{x}, \mathbf{x}' \in \mathcal{X}.$$

### Remarks :

- As a recall, a Hilbert space is a complete vector space endowed with an inner product.
- In other words, the kernel operates a measure of similarity on data, made explicit as an inner product after application of  $\varphi$ .
- With a good choice for  $\varphi$ , one may expect to transform their nonlinear prediction task into a linear one in the space  $\mathcal{H}$ .
- Any algorithm that solely involves dot products on  $\mathcal{X} \times \mathcal{X}$  may be generalized at a cheap cost by replacing the dot product with a p.d. kernel: this procedure is often called the **kernel trick**.

Proof : On the black board

## ■ Representer theorem

### Proposition : Representer theorem

Let  $\mathcal{X}$  be a set with a p.d. kernel  $K$ ,  $\mathcal{H}$  be the corresponding RKHS, and  $S = \{\mathbf{x}_1, \dots, \mathbf{x}_n\} \subset \mathcal{X}$  a finite set of points in  $\mathcal{X}$ .

Let  $\psi : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  be a function of  $n + 1$  variables strictly increasing w.r.t. the last variable. Then any solution of

$$\min_{f \in \mathcal{H}} \psi(f(\mathbf{x}_1), \dots, f(\mathbf{x}_n), \|f\|_{\mathcal{H}})$$

admits a representation of the form :

$$f(\mathbf{x}) = \sum_{i=1}^n \alpha_i K(\mathbf{x}_i, \mathbf{x}) = \sum_{i=1}^n \alpha_i K_{\mathbf{x}_i}(\mathbf{x}), \quad \forall \mathbf{x} \in \mathcal{X}.$$

### Remarks :

- In other words, the solution lives in the finite dimensional space  $\text{Span}(K_{\mathbf{x}_1}, \dots, K_{\mathbf{x}_n})$ .
- From a numerical point of view, this is a very powerful property!
- A typical form for  $\psi$  is :

$$\psi(f(\mathbf{x}_1), \dots, f(\mathbf{x}_n), \|f\|_{\mathcal{H}}) = c(f(\mathbf{x}_1), \dots, f(\mathbf{x}_n)) + \lambda \Omega(\|f\|_{\mathcal{H}})$$

**Enforces fit to data**

**w.r.t. the kernel geometry**

**Regularizer**

# **Implementation**

