

# Asymptotic Analysis

# Primality Testing

# Primality Testing

## Definition

# Primality Testing

Definition: Let  $num \geq 2$  be an integer.

# Primality Testing

Definition: Let  $num \geq 2$  be an integer. We say that *num is prime*, if its only divisors are 1 and *num*

# Primality Testing

Definition: Let  $num \geq 2$  be an integer. We say that *num is prime*, if its only divisors are 1 and *num*

Examples:

13 is prime

# Primality Testing

Definition: Let  $num \geq 2$  be an integer. We say that *num is prime*, if its only divisors are 1 and *num*

Examples:

13 is prime

12 is not prime

# Primality Testing

Definition: Let  $num \geq 2$  be an integer. We say that *num is prime*, if its only divisors are 1 and *num*

Examples:

13 is prime

12 is not prime

Definition:



# Primality Testing

Definition: Let  $num \geq 2$  be an integer. We say that *num is prime*, if its only divisors are 1 and *num*

Examples:

13 is prime

12 is not prime

Definition: Let  $num \geq 2$  be an integer

# Primality Testing

Definition: Let  $num \geq 2$  be an integer. We say that *num is prime*, if its only divisors are 1 and *num*

Examples:

13 is prime

12 is not prime

Definition: Let  $num \geq 2$  be an integer, and let  $d$  and  $k$  be two divisors of *num*.

# Primality Testing

Definition: Let  $num \geq 2$  be an integer. We say that *num is prime*, if its only divisors are 1 and  $num$

Examples:

13 is prime

12 is not prime

Definition: Let  $num \geq 2$  be an integer, and let  $d$  and  $k$  be two divisors of  $num$ . We say that *k and d are complementary divisors of num*

# Primality Testing

Definition: Let  $num \geq 2$  be an integer. We say that *num is prime*, if its only divisors are 1 and  $num$

Examples:

13 is prime

12 is not prime

Definition: Let  $num \geq 2$  be an integer, and let  $d$  and  $k$  be two divisors of  $num$ . We say that *k and d are complementary divisors of num*, if  $d \cdot k = num$

# Primality Testing

Definition: Let  $num \geq 2$  be an integer. We say that *num is prime*, if its only divisors are 1 and  $num$

Examples:

13 is prime

12 is not prime

Definition: Let  $num \geq 2$  be an integer, and let  $d$  and  $k$  be two divisors of  $num$ . We say that *k and d are complementary divisors of num*, if  $d \cdot k = num$

Examples:

4 and 25 are complementary divisors of 100

# Primality Testing

Definition: Let  $num \geq 2$  be an integer. We say that *num is prime*, if its only divisors are 1 and  $num$

Examples:

13 is prime

12 is not prime

Definition: Let  $num \geq 2$  be an integer, and let  $d$  and  $k$  be two divisors of  $num$ . We say that *k and d are complementary divisors of num*, if  $d \cdot k = num$


Examples:

4 and 25 are complementary divisors of 100

5 and 20 are complementary divisors of 100

# Primality Testing

Version I:  $1, 2, 3, \dots, num$



# Primality Testing

Version I: 1, 2, 3, . . . , *num*



Version II:




# Primality Testing

Version I:  $1, 2, 3, \dots, \dots, \dots, num$



Version II:  $1, 2, 3, \dots, \dots, \dots, \frac{num}{2}, \dots, \dots, num$




# Primality Testing

Version I:  $1, 2, 3, \dots, \dots, \dots, num$



Version II:  $1, 2, 3, \dots, \dots, \dots, \frac{num}{2}, \dots, \dots, num$




$num = 100$ :

# Primality Testing

Version I: 1, 2, 3, . . . ,  $num$



Version II: 1, 2, 3, . . . ,  $\frac{num}{2}$ , . . . ,  $num$




$num = 100$ : 1 2 4 5 10 20 25 50 100

# Primality Testing

Version I: 1, 2, 3, . . . ,  $num$




Version II: 1, 2, 3, . . . ,  $\frac{num}{2}$ , . . . ,  $num$



$num = 100$ : 1 2 4 5 10 20 25 50 100

$\frac{100}{2}$




# Primality Testing

$1, 2, 3, \dots, \frac{num}{2}, \dots, num$

# Primality Testing


$1, 2, 3, \dots, \frac{num}{2}, \dots, num$



The diagram illustrates a sequence of numbers from 1 to  $num$ . The numbers  $1, 2, 3, \dots, \frac{num}{2}, \dots, num$  are shown. The portion of the sequence from  $\frac{num}{2}$  to  $num$  is enclosed in an orange oval. An orange arrow labeled  $k$  points to the first number within this oval, representing the starting point for testing divisibility.

# Primality Testing


$1, 2, 3, \dots, \frac{num}{2}, \dots, num$



Let  $k$  be a divisor of  $num$  in the second half of the range.

# Primality Testing

$1, 2, 3, \dots, \frac{num}{2}, \dots, num$




*k*

Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$



# Primality Testing

$1, 2, 3, \dots, \frac{num}{2}, \dots, num$




*k*

Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$

Let  $d$  be  $k$ 's complementary divisor

# Primality Testing

$1, 2, 3, \dots, \frac{num}{2}, \dots, num$




$k$

Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$

Let  $d$  be  $k$ 's complementary divisor, therefore  $d = \frac{num}{k}$

# Primality Testing

$$1, 2, 3, \dots, \frac{num}{2}, \dots, num$$



Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$

Let  $d$  be  $k$ 's complementary divisor, therefore  $d = \frac{num}{k}$

We have:

# Primality Testing

$1, 2, 3, \dots, \frac{num}{2}, \dots, num$




Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$

Let  $d$  be  $k$ 's complementary divisor, therefore  $d = \frac{num}{k}$

We have:  $d = \frac{num}{k}$

# Primality Testing


$1, 2, 3, \dots, \frac{num}{2}, \dots, num$ 


Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$

Let  $d$  be  $k$ 's complementary divisor, therefore  $d = \frac{num}{k}$

We have:  $d = \frac{num}{k} \neq \frac{num}{\left(\frac{num}{2}\right)}$

# Primality Testing

$$1, 2, 3, \dots, \frac{num}{2}, \dots, num$$



*k*

Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$

Let  $d$  be  $k$ 's complementary divisor, therefore  $d = \frac{num}{k}$

We have:  $d = \frac{num}{k} < \frac{num}{\left(\frac{num}{2}\right)}$

# Primality Testing

$$1, 2, 3, \dots, \frac{num}{2}, \dots, num$$



*k*

Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$

Let  $d$  be  $k$ 's complementary divisor, therefore  $d = \frac{num}{k}$

We have:  $d = \frac{num}{k} < \frac{num}{\left(\frac{num}{2}\right)} = \frac{num}{1} \cdot \frac{2}{num}$

# Primality Testing

$$1, 2, 3, \dots, \frac{num}{2}, \dots, num$$



Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$

Let  $d$  be  $k$ 's complementary divisor, therefore  $d = \frac{num}{k}$

We have: 
$$d = \frac{num}{k} < \frac{num}{\left(\frac{num}{2}\right)} = \frac{num}{1} \cdot \frac{2}{num} = 2$$



# Primality Testing

$$1, 2, 3, \dots, \frac{num}{2}, \dots, num$$



Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$

Let  $d$  be  $k$ 's complementary divisor, therefore  $d = \frac{num}{k}$

We have:

$$d = \frac{num}{k} < \frac{num}{\left(\frac{num}{2}\right)} = \frac{num}{1} \cdot \frac{2}{num} = 2$$

# Primality Testing

$$1, 2, 3, \dots, \frac{num}{2}, \dots, num$$


Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$


Let  $d$  be  $k$ 's complementary divisor, therefore  $d = \frac{num}{k}$

We have:

$$d = \frac{num}{k} < \frac{num}{\left(\frac{num}{2}\right)} = \frac{num}{1} \cdot \frac{2}{num} = 2$$

Therefore  $d < 2$ .

# Primality Testing

$$1, 2, 3, \dots, \frac{num}{2}, \dots, num$$



Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$

Let  $d$  be  $k$ 's complementary divisor, therefore  $d = \frac{num}{k}$

We have: 
$$d = \frac{num}{k} < \frac{num}{\left(\frac{num}{2}\right)} = \frac{num}{1} \cdot \frac{2}{num} = 2$$

Therefore  $d < 2$ . Since  $d$  is a divisor

# Primality Testing

$$1, 2, 3, \dots, \frac{num}{2}, \dots, num$$



Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$

Let  $d$  be  $k$ 's complementary divisor, therefore  $d = \frac{num}{k}$

We have:  $d = \frac{num}{k} < \frac{num}{\left(\frac{num}{2}\right)} = \frac{num}{1} \cdot \frac{2}{num} = 2$

Therefore  $d < 2$ . Since  $d$  is a divisor

# Primality Testing

$$1, 2, 3, \dots, \frac{num}{2}, \dots, num$$



Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$

Let  $d$  be  $k$ 's complementary divisor, therefore  $d = \frac{num}{k}$

We have: 
$$d = \frac{num}{k} < \frac{num}{\left(\frac{num}{2}\right)} = \frac{num}{1} \cdot \frac{2}{num} = 2$$

Therefore  $d < 2$ . Since  $d$  is a divisor, we get that  $d = 1$ .

# Primality Testing

$1, 2, 3, \dots, \frac{num}{2}, \dots, num$ 


Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$


Let  $d$  be  $k$ 's complementary divisor, therefore  $d = \frac{num}{k}$

We have:  $d = \frac{num}{k} < \frac{num}{\left(\frac{num}{2}\right)} = \frac{num}{1} \cdot \frac{2}{num} = 2$

Therefore  $d < 2$ . Since  $d$  is a divisor, we get that  $d = 1$ .

# Primality Testing

$1, 2, 3, \dots, \frac{num}{2}, \dots, num$



Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$


Let  $d$  be  $k$ 's complementary divisor, therefore  $d = \frac{num}{k}$

We have:  $d = \frac{num}{k} < \frac{num}{\left(\frac{num}{2}\right)} = \frac{num}{1} \cdot \frac{2}{num} = 2$

Therefore  $d < 2$ . Since  $d$  is a divisor, we get that  $d = 1$ .

So:  $\frac{num}{k} = 1$

# Primality Testing

$1, 2, 3, \dots, \frac{num}{2}, \dots, num$ 


Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$

Let  $d$  be  $k$ 's complementary divisor, therefore  $d = \frac{num}{k}$

We have:  $d = \frac{num}{k} < \frac{num}{\left(\frac{num}{2}\right)} = \frac{num}{1} \cdot \frac{2}{num} = 2$


Therefore  $d < 2$ . Since  $d$  is a divisor, we get that  $d = 1$ .

So:  $\frac{num}{k} = 1$ , therefore  $k = num$ .



# Primality Testing

$1, 2, 3, \dots, \frac{num}{2}, \dots, num$



Let  $k$  be a divisor of  $num$  in the second half of the range. That is,  $k > \frac{num}{2}$

Let  $d$  be  $k$ 's complementary divisor, therefore  $d = \frac{num}{k}$

We have:  $d = \frac{num}{k} < \frac{num}{\left(\frac{num}{2}\right)} = \frac{num}{1} \cdot \frac{2}{num} = 2$

Therefore  $d < 2$ . Since  $d$  is a divisor, we get that  $d = 1$ .

So:  $\frac{num}{k} = 1$ , therefore  $k = num$ .


This shows that the only divisor in the second half of the range is  $num$  itself.

# Primality Testing

Version I: 1, 2, 3, . . . ,  $num$



Version II: 1, 2, 3, . . . ,  $\frac{num}{2}$ , . . . ,  $num$




# Primality Testing

Version I: 1, 2, 3, . . . ,  $num$



Version II: 1, 2, 3, . . . ,  $\frac{num}{2}$ , . . . ,  $num$



Version III:

# Primality Testing

Version I:  $1, 2, 3, \dots, \dots, \dots, num$

Version II:  $1, 2, 3, \dots, \dots, \dots, \frac{num}{2}, \dots, \dots, num$

Version III:  $1, 2, 3, \dots, \sqrt{num}, \dots, \dots, \dots, num$

# Primality Testing

Version I:  $1, 2, 3, \dots, num$

Version II:  $1, 2, 3, \dots, \frac{num}{2}, \dots, num$

Version III:  $1, 2, 3, \dots, \sqrt{num}, \dots, num$

$num = 100$ : 1 2 4 5 10 20 25 50 100


# Primality Testing

Version I:  $1, 2, 3, \dots, num$

Version II:  $1, 2, 3, \dots, \frac{num}{2}, \dots, num$

Version III:  $1, 2, 3, \dots, \sqrt{num}, \dots, num$

$num = 100$ : 1 2 4 5 10 20 25 50 100

$\frac{100}{2}$  

# Primality Testing

Version I:  $1, 2, 3, \dots, \dots, \dots, num$

Version II:  $1, 2, 3, \dots, \dots, \dots, \frac{num}{2}, \dots, \dots, num$

Version III:  $1, 2, 3, \dots, \sqrt{num}, \dots, \dots, num$

$num = 100$ : 1 2 4 5 10 20 25 50 100

$\swarrow$   
 $\sqrt{100}$

$\swarrow$   
 $\frac{100}{2}$

# Primality Testing

1 2 4 5 10 20 25 50 100




# Primality Testing


1 2 4 5 10 20 25 50 100

A vertical red line is drawn through the number 10, extending from above the text to below it.


# Primality Testing




# Primality Testing




# Primality Testing



# Primality Testing





# Primality Testing




$1, 2, 3, \dots, \sqrt{num}, \dots, num$


# Primality Testing



$1, 2, 3, \dots, \sqrt{num},$  ,  $num$

  $k, d$

# Primality Testing




$1, 2, 3, \dots, \sqrt{num},$   $\cdot \quad \cdot \quad \cdot$   $, num$


$\swarrow$   
 $k, d$


Let  $k$  and  $d$  be complementary divisors of  $num$



# Primality Testing





$1, 2, 3, \dots, \sqrt{num},$    $, num$

  $k, d$

Let  $k$  and  $d$  be complementary divisors of  $num$ , and assume that they are both greater than  $\sqrt{num}$

# Primality Testing




$1, 2, 3, \dots, \sqrt{num},$    $, num$


$k, d$

Let  $k$  and  $d$  be complementary divisors of  $num$ , and assume that they are both greater than  $\sqrt{num}$

We therefore have:

# Primality Testing




$1, 2, 3, \dots, \sqrt{num},$    $, num$


*k, d*


Let  $k$  and  $d$  be complementary divisors of  $num$ , and assume that they are both greater than  $\sqrt{num}$

We therefore have:  $num = k \cdot d$

# Primality Testing




$1, 2, 3, \dots, \sqrt{num},$    $, num$


  $k, d$


Let  $k$  and  $d$  be complementary divisors of  $num$ , and assume that they are both greater than  $\sqrt{num}$

We therefore have:  $num = k \cdot d > \sqrt{num} \cdot \sqrt{num}$

# Primality Testing




$1, 2, 3, \dots, \sqrt{num},$    $, num$


  $k, d$

Let  $k$  and  $d$  be complementary divisors of  $num$ , and assume that they are both greater than  $\sqrt{num}$

We therefore have:  $num = k \cdot d > \sqrt{num} \cdot \sqrt{num} = num$

# Primality Testing




$1, 2, 3, \dots, \sqrt{num},$ 

 $, num$


Let  $k$  and  $d$  be complementary divisors of  $num$ , and assume that they are both greater than  $\sqrt{num}$

We therefore have:  $num = k \cdot d > \sqrt{num} \cdot \sqrt{num} = num$

This implies that  $num > num$

# Primality Testing



$1, 2, 3, \dots, \sqrt{num},$ 
,
 $num$


$k, d$


Let  $k$  and  $d$  be complementary divisors of  $num$ , and assume that they are both greater than  $\sqrt{num}$

We therefore have:  $num = k \cdot d > \sqrt{num} \cdot \sqrt{num} = num$

This implies that  $num > num$ , which is a contradiction.

# Primality Testing



$1, 2, 3, \dots, \sqrt{num},$ 

 $, num$


Let  $k$  and  $d$  be complementary divisors of  $num$ , and assume that they are both greater than  $\sqrt{num}$


We therefore have:  $num = k \cdot d > \sqrt{num} \cdot \sqrt{num} = num$

This implies that  $num > num$ , which is a contradiction.



# Primality Testing



$1, 2, 3, \dots, \sqrt{num},$    $, num$   
*k, d*

Let  $k$  and  $d$  be complementary divisors of  $num$ , and assume that they are both greater than  $\sqrt{num}$

We therefore have:  $num = k \cdot d > \sqrt{num} \cdot \sqrt{num} = num$

This implies that  $num > num$ , which is a contradiction.

This shows that at least one in each pair of complementary divisors is less than or equal to  $\sqrt{num}$

# Primality Testing

Version I:  $1, 2, 3, \dots, \dots, \dots, num$

Version II:  $1, 2, 3, \dots, \dots, \dots, \frac{num}{2}, \dots, \dots, num$

Version III:  $1, 2, 3, \dots, \sqrt{num}, \dots, \dots, \dots, num$

# Runtime Analysis

# Runtime Analysis

- The running time depends on the size of the input

# Runtime Analysis

- The running time depends on the size of the input
- The running time depends on the operators we use, and on the types of the data they are applied on

# Runtime Analysis

- The running time depends on the size of the input
- The running time depends on the operators we use, and on the types of the data they are applied on
- The running time depends on the machine's hardware technology

# Runtime Analysis

- The running time depends on the size of the input
  - ✓ We parameterize running time by the size of the input
- The running time depends on the operators we use, and on the types of the data they are applied on
- The running time depends on the machine's hardware technology

# Runtime Analysis

- The running time depends on the size of the input
  - ✓ We parameterize running time by the size of the input
- The running time depends on the operators we use, and on the types of the data they are applied on
  - ✓ The abstract model we use, ignores machine-dependent constants.
    - We count each primitive operation as 1
- The running time depends on the machine's hardware technology



# Runtime Analysis

- The running time depends on the size of the input
  - ✓ We parameterize running time by the size of the input
- The running time depends on the operators we use, and on the types of the data they are applied on
  - ✓ The abstract model we use, ignores machine-dependent constants.  
We count each primitive operation as 1
- The running time depends on the machine's hardware technology
  - ✓ The abstract model we use, divides the algorithms to classes based on their "quality".  
We make asymptotic analysis: look at the order of growth of  $T(n)$

# Runtime Analysis

## Informal Criteria

# Runtime Analysis

## Informal Criteria

We compare the **asymptotic order** of the **number of primitive operations** executed by a process, **as a function of its input size**

# Runtime Analysis

## Informal Criteria

We compare the **asymptotic order** of the **number of primitive operations** executed by a process, **as a function of its input size**

$$T(n) = 3n^2 + 6n - 15$$

# Runtime Analysis

## Informal Criteria

We compare the **asymptotic order** of the **number of primitive operations** executed by a process, **as a function of its input size**

$$T(n) = 3n^2 + 6n - 15 = \theta(n^2)$$

# Runtime Analysis

## Informal Criteria

We compare the **asymptotic order** of the **number of primitive operations** executed by a process, **as a function of its input size**

$$T(n) = 3n^2 + 6n - 15 = \theta(n^2)$$

Rule of thumb to get the order of growth:

# Runtime Analysis

## Informal Criteria

We compare the **asymptotic order** of the **number of primitive operations** executed by a process, **as a function of its input size**

$$T(n) = 3n^2 + \cancel{6n - 15} = \theta(n^2)$$

Rule of thumb to get the order of growth:

- Drop low-order terms

# Runtime Analysis

## Informal Criteria

We compare the **asymptotic order** of the **number of primitive operations** executed by a process, **as a function of its input size**

$$T(n) = \cancel{3n^2} + \cancel{6n - 15} = \theta(n^2)$$

Rule of thumb to get the order of growth:

- Drop low-order terms
- Ignore leading constants



# Runtime Analysis

## Informal Criteria

We compare the **asymptotic order** of the **number of primitive operations** executed by a process, **as a function of its input size**

$$T(n) = \cancel{3n^2} + \cancel{6n - 15} = \theta(n^2)$$

Rule of thumb to get the order of growth:

- Drop low-order terms
- Ignore leading constants

More Formally . . .

# Asymptotic Analysis

$O$  definition

# Asymptotic Analysis

## O definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

# Asymptotic Analysis

## O definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = O(g(n))$

# Asymptotic Analysis

## O definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = O(g(n))$  if there exist positive real constant  $c$

# Asymptotic Analysis

## O definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = O(g(n))$  if there exist positive real constant  $c$  and a positive integer constant  $n_0$

# Asymptotic Analysis

## O definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = O(g(n))$  if there exist positive real constant  $c$  and a positive integer constant  $n_0$  such that  $f(n) \leq c \cdot g(n)$  for all  $n \geq n_0$


# Asymptotic Analysis

## $O$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = O(g(n))$  if there exist positive real constant  $c$  and a positive integer constant  $n_0$  such that  $f(n) \leq c \cdot g(n)$  for all  $n \geq n_0$






# Asymptotic Analysis

## O definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = O(g(n))$  if there exist positive real constant  $c$  and a positive integer constant  $n_0$  such that  $f(n) \leq c \cdot g(n)$  for all  $n \geq n_0$




# Asymptotic Analysis

## $O$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = O(g(n))$  if there exist positive real constant  $c$  and a positive integer constant  $n_0$  such that  $f(n) \leq c \cdot g(n)$  for all  $n \geq n_0$




# Asymptotic Analysis

## O definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = O(g(n))$  if there exist positive real constant  $c$  and a positive integer constant  $n_0$  such that  $f(n) \leq c \cdot g(n)$  for all  $n \geq n_0$




# Asymptotic Analysis

## $O$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = O(g(n))$  if there exist positive real constant  $c$  and a positive integer constant  $n_0$  such that  $f(n) \leq c \cdot g(n)$  for all  $n \geq n_0$



# Asymptotic Analysis

$\Omega$  definition

# Asymptotic Analysis

## $\Omega$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

# Asymptotic Analysis

## $\Omega$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Omega(g(n))$

# Asymptotic Analysis

## $\Omega$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Omega(g(n))$  if there exist positive real constant  $c$



# Asymptotic Analysis

## $\Omega$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Omega(g(n))$  if there exist positive real constant  $c$  and a positive integer constant  $n_0$

# Asymptotic Analysis

## $\Omega$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Omega(g(n))$  if there exist positive real constant  $c$  and a positive integer constant  $n_0$  such that  $f(n) \geq c \cdot g(n)$  for all  $n \geq n_0$


# Asymptotic Analysis

## $\Omega$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Omega(g(n))$  if there exist positive real constant  $c$  and a positive integer constant  $n_0$  such that  $f(n) \geq c \cdot g(n)$  for all  $n \geq n_0$




# Asymptotic Analysis

## $\Omega$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Omega(g(n))$  if there exist positive real constant  $c$  and a positive integer constant  $n_0$  such that  $f(n) \geq c \cdot g(n)$  for all  $n \geq n_0$




# Asymptotic Analysis

## $\Omega$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Omega(g(n))$  if there exist positive real constant  $c$  and a positive integer constant  $n_0$  such that  $f(n) \geq c \cdot g(n)$  for all  $n \geq n_0$




# Asymptotic Analysis

## $\Omega$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Omega(g(n))$  if there exist positive real constant  $c$  and a positive integer constant  $n_0$  such that  $f(n) \geq c \cdot g(n)$  for all  $n \geq n_0$




# Asymptotic Analysis

## $\Omega$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Omega(g(n))$  if there exist positive real constant  $c$  and a positive integer constant  $n_0$  such that  $f(n) \geq c \cdot g(n)$  for all  $n \geq n_0$



# Asymptotic Analysis

$\Theta$  definition



# Asymptotic Analysis

## $\Theta$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

# Asymptotic Analysis

## $\Theta$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \theta(g(n))$

# Asymptotic Analysis

## $\Theta$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Theta(g(n))$  if there exist positive real constants  $c_1, c_2$

# Asymptotic Analysis

## $\Theta$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Theta(g(n))$  if there exist positive real constants  $c_1, c_2$  and a positive integer constant  $n_0$

# Asymptotic Analysis

## $\Theta$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Theta(g(n))$  if there exist positive real constants  $c_1, c_2$  and a positive integer constant  $n_0$  such that  $c_2 g(n) \leq f(n) \leq c_1 g(n)$  for all  $n \geq n_0$


# Asymptotic Analysis

## $\Theta$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Theta(g(n))$  if there exist positive real constants  $c_1, c_2$  and a positive integer constant  $n_0$  such that  $c_2g(n) \leq f(n) \leq c_1g(n)$  for all  $n \geq n_0$




# Asymptotic Analysis

## $\Theta$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Theta(g(n))$  if there exist positive real constants  $c_1, c_2$  and a positive integer constant  $n_0$  such that  $c_2 g(n) \leq f(n) \leq c_1 g(n)$  for all  $n \geq n_0$




# Asymptotic Analysis

## $\Theta$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Theta(g(n))$  if there exist positive real constants  $c_1, c_2$  and a positive integer constant  $n_0$  such that  $c_2 g(n) \leq f(n) \leq c_1 g(n)$  for all  $n \geq n_0$






# Asymptotic Analysis

## $\Theta$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Theta(g(n))$  if there exist positive real constants  $c_1, c_2$  and a positive integer constant  $n_0$  such that  $c_2 g(n) \leq f(n) \leq c_1 g(n)$  for all  $n \geq n_0$



# Asymptotic Analysis

## $\Theta$ definition

### Definition


Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Theta(g(n))$  if there exist positive real constants  $c_1, c_2$  and a positive integer constant  $n_0$  such that  $c_2 g(n) \leq f(n) \leq c_1 g(n)$  for all  $n \geq n_0$

$$n_0 = 8$$

$$c_1 = 4$$

$$c_2 = \frac{1}{3}$$




# Asymptotic Analysis

## $\Theta$ definition

### Definition

Let  $f(n)$  and  $g(n)$  be two functions mapping positive integers to positive real numbers.

We say that  $f(n) = \Theta(g(n))$  if there exist positive real constants  $c_1, c_2$  and a positive integer constant  $n_0$  such that  $c_2 g(n) \leq f(n) \leq c_1 g(n)$  for all  $n \geq n_0$



# Asymptotic Analysis

$\Theta$  definition

Show that:  $3n^2 + 6n - 15 = \theta(n^2)$

# Asymptotic Analysis

$\Theta$  definition

Show that:

$$\underbrace{3n^2 + 6n - 15}_{f(n)} = \theta(\underbrace{n^2}_{g(n)})$$

# Asymptotic Analysis

$\Theta$  definition

Show that:

$$\underbrace{3n^2 + 6n - 15}_{f(n)} = \theta(\underbrace{n^2}_{g(n)})$$

Proof:

# Asymptotic Analysis

## $\Theta$ definition

Show that:

$$\underbrace{3n^2 + 6n - 15}_{f(n)} = \theta(\underbrace{n^2}_{g(n)})$$

Proof:

if we take

$$c_1 = \underline{\hspace{2cm}}$$

$$c_2 = \underline{\hspace{2cm}}$$

$$n_0 = \underline{\hspace{2cm}}$$

# Asymptotic Analysis

## $\Theta$ definition

Show that:  $\underbrace{3n^2 + 6n - 15}_{f(n)} = \theta(\underbrace{n^2}_{g(n)})$

Proof:

if we take  $c_1 = \underline{\hspace{2cm}}$

$c_2 = \underline{\hspace{2cm}}$

$n_0 = \underline{\hspace{2cm}}$

Then for all  $n \geq n_0$  we have:



# Asymptotic Analysis

## $\Theta$ definition

Show that:  $\underbrace{3n^2 + 6n - 15}_{f(n)} = \theta(\underbrace{n^2}_{g(n)})$

Proof:

if we take  $c_1 = \underline{\hspace{2cm}}$

$c_2 = \underline{\hspace{2cm}}$

$n_0 = \underline{\hspace{2cm}}$

Then for all  $n \geq n_0$  we have:

$$3n^2 + 6n - 15$$

# Asymptotic Analysis

## $\Theta$ definition

Show that:  $\underbrace{3n^2 + 6n - 15}_{f(n)} = \theta(\underbrace{n^2}_{g(n)})$

Proof:

if we take  $c_1 = \underline{\hspace{2cm}}$

$c_2 = \underline{\hspace{2cm}}$

$n_0 = \underline{\hspace{2cm}}$

Then for all  $n \geq n_0$  we have:

$$3n^2 + 6n - 15 \leq 3n^2 + 6n$$

# Asymptotic Analysis

## $\Theta$ definition

Show that:  $\underbrace{3n^2 + 6n - 15}_{f(n)} = \theta(\underbrace{n^2}_{g(n)})$

Proof:

if we take  $c_1 = \underline{\hspace{2cm}}$

$c_2 = \underline{\hspace{2cm}}$

$n_0 = \underline{\hspace{2cm}}$

Then for all  $n \geq n_0$  we have:

$$3n^2 + 6n - 15 \leq 3n^2 + 6n \leq 3n^2 + 6n^2$$

# Asymptotic Analysis

## $\Theta$ definition

Show that:  $\underbrace{3n^2 + 6n - 15}_{f(n)} = \theta(\underbrace{n^2}_{g(n)})$

Proof:

if we take  $c_1 = \underline{\hspace{2cm}}$

$c_2 = \underline{\hspace{2cm}}$

$n_0 = \underline{\hspace{2cm}}$

Then for all  $n \geq n_0$  we have:

$$3n^2 + 6n - 15 \leq 3n^2 + 6n \leq 3n^2 + 6n^2 = 9n^2$$

# Asymptotic Analysis

## $\Theta$ definition

Show that:  $\underbrace{3n^2 + 6n - 15}_{f(n)} = \theta(\underbrace{n^2}_{g(n)})$

Proof:

if we take  $c_1 = \underline{9}$

$c_2 = \underline{\hspace{2cm}}$

$n_0 = \underline{\hspace{2cm}}$

Then for all  $n \geq n_0$  we have:

$$3n^2 + 6n - 15 \leq 3n^2 + 6n \leq 3n^2 + 6n^2 = 9n^2$$

# Asymptotic Analysis

## $\Theta$ definition

Show that:  $\underbrace{3n^2 + 6n - 15}_{f(n)} = \theta(\underbrace{n^2}_{g(n)})$

Proof:

if we take  $c_1 = \underline{9}$

$c_2 = \underline{\hspace{2cm}}$

$n_0 = \underline{\hspace{2cm}}$

Then for all  $n \geq n_0$  we have:

$$3n^2 \leq 3n^2 + 6n - 15 \leq 3n^2 + 6n \leq 3n^2 + 6n^2 = 9n^2$$

# Asymptotic Analysis

## $\Theta$ definition

Show that:  $\underbrace{3n^2 + 6n - 15}_{f(n)} = \theta(\underbrace{n^2}_{g(n)})$

Proof:

if we take  $c_1 = \underline{9}$

$c_2 = \underline{\hspace{2cm}}$

$n_0 = \underline{\hspace{2cm}}$

Then for all  $n \geq n_0$  we have:

$$3n^2 \leq 3n^2 + 6n - 15 \leq 3n^2 + 6n \leq 3n^2 + 6n^2 = 9n^2$$


$$6n - 15 \geq 0$$

# Asymptotic Analysis

## $\Theta$ definition

Show that:  $\underbrace{3n^2 + 6n - 15}_{f(n)} = \theta(\underbrace{n^2}_{g(n)})$

Proof:

if we take  $c_1 = \underline{9}$

$c_2 = \underline{\hspace{2cm}}$

$n_0 = \underline{\hspace{2cm}}$

Then for all  $n \geq n_0$  we have:

$$3n^2 \leq 3n^2 + 6n - 15 \leq 3n^2 + 6n \leq 3n^2 + 6n^2 = 9n^2$$


$$6n - 15 \geq 0$$

$$6n \geq 15$$



# Asymptotic Analysis

## $\Theta$ definition

Show that:  $\underbrace{3n^2 + 6n - 15}_{f(n)} = \theta(\underbrace{n^2}_{g(n)})$

Proof:

if we take  $c_1 = \underline{9}$

$c_2 = \underline{\hspace{2cm}}$

$n_0 = \underline{\hspace{2cm}}$

Then for all  $n \geq n_0$  we have:

$$3n^2 \leq 3n^2 + 6n - 15 \leq 3n^2 + 6n \leq 3n^2 + 6n^2 = 9n^2$$

$$\begin{array}{c} 6n - 15 \geq 0 \\ \updownarrow \\ 6n \geq 15 \\ \updownarrow \\ n \geq 2.5 \end{array}$$

# Asymptotic Analysis

## $\Theta$ definition

Show that:  $\underbrace{3n^2 + 6n - 15}_{f(n)} = \theta(\underbrace{n^2}_{g(n)})$

Proof:

if we take  $c_1 = \underline{9}$

$c_2 = \underline{\hspace{1cm}}$

$n_0 = \underline{3}$

Then for all  $n \geq n_0$  we have:

$$3n^2 \leq 3n^2 + 6n - 15 \leq 3n^2 + 6n \leq 3n^2 + 6n^2 = 9n^2$$

$$\begin{array}{c} 6n - 15 \geq 0 \\ \updownarrow \\ 6n \geq 15 \\ \updownarrow \\ n \geq 2.5 \end{array}$$

# Asymptotic Analysis

## $\Theta$ definition

Show that:  $\underbrace{3n^2 + 6n - 15}_{f(n)} = \theta(\underbrace{n^2}_{g(n)})$

Proof:

if we take  $c_1 = \underline{9}$

$$c_2 = \underline{3}$$

$$n_0 = \underline{3}$$

Then for all  $n \geq n_0$  we have:

$$3n^2 \leq 3n^2 + 6n - 15 \leq 3n^2 + 6n \leq 3n^2 + 6n^2 = 9n^2$$

$$\begin{array}{c} 6n - 15 \geq 0 \\ \updownarrow \\ 6n \geq 15 \\ \updownarrow \\ n \geq 2.5 \end{array}$$

# Asymptotic Analysis

## $\Theta$ definition

Show that:  $\underbrace{3n^2 + 6n - 15}_{f(n)} = \theta(\underbrace{n^2}_{g(n)})$

Proof:

if we take  $c_1 = \underline{9}$

$$c_2 = \underline{3}$$

$$n_0 = \underline{3}$$

Then for all  $n \geq n_0$  we have:

$$3n^2 \leq 3n^2 + 6n - 15 \leq 3n^2 + 6n \leq 3n^2 + 6n^2 = 9n^2$$

$\Downarrow$

$$3n^2 \leq 3n^2 + 6n - 15 \leq 9n^2$$

$$\begin{array}{c} 6n - 15 \geq 0 \\ \updownarrow \\ 6n \geq 15 \\ \updownarrow \\ n \geq 2.5 \end{array}$$

# Asymptotic Analysis

## $\Theta$ definition

Show that:  $\underbrace{3n^2 + 6n - 15}_{f(n)} = \theta(\underbrace{n^2}_{g(n)})$

Proof:

if we take  $c_1 = \underline{9}$

$$c_2 = \underline{3}$$


$$n_0 = \underline{3}$$

Then for all  $n \geq n_0$  we have:

$$3n^2 \leq 3n^2 + 6n - 15 \leq 3n^2 + 6n \leq 3n^2 + 6n^2 = 9n^2$$



$$3n^2 \leq 3n^2 + 6n - 15 \leq 9n^2$$


$$\begin{aligned} 6n - 15 &\geq 0 \\ \updownarrow \\ 6n &\geq 15 \\ \updownarrow \\ n &\geq 2.5 \end{aligned}$$

Therefore:  $3n^2 + 6n - 15 = \theta(n^2)$

# The Searching Problem

# The Searching Problem

## Problem

Implement the following function:

```
def linear_search(lst, val)
```

# The Searching Problem

## Problem

Implement the following function:

```
def linear_search(lst, val)
```

The function should return an index in lst, where val appears first



# The Searching Problem

## Problem

Implement the following function:

```
def linear_search(lst, val)
```

The function should return an index in lst, where val appears first, or **None** if val is not one of lst's elements.

# The Searching Problem

## Problem

Implement the following function:

```
def linear_search(lst, val)
```

The function should return an index in lst, where val appears first, or **None** if val is not one of lst's elements.

## Examples

If lst is: [5, 8, 12, 7, 8, 10]

# The Searching Problem

## Problem

Implement the following function:

```
def linear_search(lst, val)
```

The function should return an index in lst, where val appears first, or **None** if val is not one of lst's elements.

## Examples

If lst is: [5, 8, 12, 7, 8, 10]

- The call: linear\_search(lst, 8)

# The Searching Problem

## Problem

Implement the following function:

```
def linear_search(lst, val)
```

The function should return an index in lst, where val appears first, or **None** if val is not one of lst's elements.

## Examples

If lst is: [5, 8, 12, 7, 8, 10]

- The call: linear\_search(lst, 8) should return 1

# The Searching Problem

## Problem

Implement the following function:

```
def linear_search(lst, val)
```

The function should return an index in lst, where val appears first, or **None** if val is not one of lst's elements.

## Examples

If lst is: [5, 8, 12, 7, 8, 10]

- The call: linear\_search(lst, 8) should return 1
- The call: linear\_search (lst, 4)

# The Searching Problem

## Problem

Implement the following function:

```
def linear_search(lst, val)
```

The function should return an index in lst, where val appears first, or **None** if val is not one of lst's elements.

## Examples

If lst is: [5, 8, 12, 7, 8, 10]

- The call: linear\_search(lst, 8) should return 1
- The call: linear\_search (lst, 4) should return **None**

# Linear Search

```
def linear_search(lst, val):
```

# Linear Search

```
def linear_search(lst, val):  
    for __ in _____:
```



# Linear Search

```
def linear_search(lst, val):  
    for i in range(len(lst)):
```

# Linear Search

```
def linear_search(lst, val):  
    for i in range(len(lst)):  
        if (lst[i] == val):  
            return i
```

# Linear Search

```
def linear_search(lst, val):  
    for i in range(len(lst)):  
        if (lst[i] == val):  
            return i  
    return None
```

# Linear Search

```
def linear_search(lst, val):  
    for i in range(len(lst)):  
        if (lst[i] == val):  
            return i  
 $\theta(1)$  return None
```

# Linear Search

```
def linear_search(lst, val):  
    for i in range(len(lst)):  
        if (lst[i] == val):  
            return i  
 $\Theta(1)$  return None
```

# Linear Search

```
def linear_search(lst, val):  
    for i in range(len(lst)):  
        if (lst[i] == val):  
            return i  
 $\Theta(1)$  return None
```

# Linear Search

```
def linear_search(lst, val):  
    for i in range(len(lst)):  
        if (lst[i] == val):  
            return i  
    return None
```

$\Theta(1)$

$\Theta(1)$

# Linear Search

```
def linear_search(lst, val):  
    for i in range(len(lst)):  
        if (lst[i] == val):  
            return i  
    return None
```

$\Theta(\# \text{ of iterations})$

$\Theta(1)$

$\Theta(1)$



# Linear Search

```
def linear_search(lst, val):  
    for i in range(len(lst)):  
        if (lst[i] == val):  
            return i  
    return None
```

$\Theta(\# \text{ of iterations})$

$\Theta(1)$

$\Theta(1)$

- $T(n) = \Theta(\# \text{ of iterations})$

# Linear Search

```
def linear_search(lst, val):  
    for i in range(len(lst)):  
        if (lst[i] == val):  
            return i  
    return None
```

$\Theta(\# \text{ of iterations})$  [bracketed next to the for loop]  
 $\Theta(1)$  [bracketed next to the if statement]  
 $\Theta(1)$  [bracketed next to the return statement]

- $T(n) = \Theta(\# \text{ of iterations})$
- *In worst-case: ( $\# \text{ of iterations}$ ) =  $n$*

# Linear Search

```
def linear_search(lst, val):  
    for i in range(len(lst)):  
        if (lst[i] == val):  
            return i  
    return None
```

$\Theta(\# \text{ of iterations})$  [bracketed next to the for loop]  
 $\Theta(1)$  [bracketed next to the if statement]  
 $\Theta(1)$  [bracketed next to the return None statement]

- $T(n) = \Theta(\# \text{ of iterations})$
- *In worst-case: (# of iterations) =  $n$*



$$T_{\text{worst}}(n) = \Theta(n)$$

# The Sorted-Search Problem

# The Sorted-Search Problem

## Problem

Implement the following function:

```
def sorted_search(srt_lst, val)
```

# The Sorted-Search Problem

## Problem

Implement the following function:

```
def sorted_search(srt_lst, val)
```

The function is given a **sorted** list `srt_lst`, and `val` to search for

# The Sorted-Search Problem

## Problem

Implement the following function:

```
def sorted_search(srt_lst, val)
```

The function is given a **sorted** list `srt_lst`, and `val` to search for. It should return an index, where `val` appears, or **None** if `val` is not one of `srt_lst`'s elements.

# The Sorted-Search Problem

## Problem

Implement the following function:

```
def sorted_search(srt_lst, val)
```

The function is given a **sorted** list `srt_lst`, and `val` to search for. It should return an index, where `val` appears, or **None** if `val` is not one of `srt_lst`'s elements.

## Example

If `srt_lst` is: [5, 7, 8, 8, 10, 12]



# The Sorted-Search Problem

## Problem

Implement the following function:

```
def sorted_search(srt_lst, val)
```

The function is given a **sorted** list `srt_lst`, and `val` to search for. It should return an index, where `val` appears, or **None** if `val` is not one of `srt_lst`'s elements.

## Example

If `srt_lst` is: [5, 7, 8, 8, 10, 12]

The call `sortedSearch(srt_lst, 8)`

# The Sorted-Search Problem

## Problem

Implement the following function:

```
def sorted_search(srt_lst, val)
```

The function is given a **sorted** list `srt_lst`, and `val` to search for. It should return an index, where `val` appears, or **None** if `val` is not one of `srt_lst`'s elements.

## Example

If `srt_lst` is: [5, 7, 8, 8, 10, 12]

The call `sortedSearch(srt_lst, 8)` could return 3


# The Sorted-Search Problem

val = 79

1	3	4	8	...
---	---	---	---	-----

# The Sorted-Search Problem

```
val = 79
```



# The Sorted-Search Problem

val = 79



# The Sorted-Search Problem

val = 79



# The Sorted-Search Problem

val = 79



# The Sorted-Search Problem

val = 79





# The Sorted-Search Problem

val = 79



$$T(n) = \Theta(n)$$


# The Sorted-Search Problem

val = 79




# The Sorted-Search Problem

val = 79



# The Sorted-Search Problem


val = 79



↑  
mid


# The Sorted-Search Problem

val = 79




# The Sorted-Search Problem

val = 79




# The Sorted-Search Problem

val = 79



# The Sorted-Search Problem


val = 79






# The Sorted-Search Problem

val = 79




# The Sorted-Search Problem

val = 79




# The Sorted-Search Problem

val = 79




# The Sorted-Search Problem

val = 79




# The Sorted-Search Problem

val = 79




# The Sorted-Search Problem

val = 79




# The Sorted-Search Problem

val = 79



# The Sorted-Search Problem


val = 79






# The Sorted-Search Problem

val = 79



# The Sorted-Search Problem

val = 79



```
def binary_search(srt_lst, val):  
    left = 0  
    right = len(srt_lst) - 1  
    ind = None  
    found = False  
    while ((found == False) and (left <= right)):  
        mid = (left + right) // 2  
        if (srt_lst[mid] == val):  
            ind = mid  
            found = True  
        elif (val < srt_lst[mid]):  
            right = mid - 1  
        else: # val > srt_lst[mid]  
            left = mid + 1  
    return ind
```

```
def binary_search(srt_lst, val):  
    left = 0  
    right = len(srt_lst) - 1  
     $\Theta(1)$  ind = None  
    found = False  
    while ((found == False) and (left <= right)):  
        mid = (left + right) // 2  
        if (srt_lst[mid] == val):  
            ind = mid  
            found = True  
        elif (val < srt_lst[mid]):  
            right = mid - 1  
        else: # val > srt_lst[mid]  
            left = mid + 1  
    return ind
```

```

def binary_search(srt_lst, val):
    left = 0
    right = len(srt_lst) - 1
    ind = None
    found = False
    while ((found == False) and (left <= right)):
        mid = (left + right) // 2
        if (srt_lst[mid] == val):
            ind = mid
            found = True
        elif (val < srt_lst[mid]):
            right = mid - 1
        else: # val > srt_lst[mid]
            left = mid + 1
    return ind

```

```

def binary_search(srt_lst, val):
    left = 0
    right = len(srt_lst) - 1
    ind = None
    found = False
    while ((found == False) and (left <= right)):
        mid = (left + right) // 2
        if (srt_lst[mid] == val):
            ind = mid
            found = True
        elif (val < srt_lst[mid]):
            right = mid - 1
        else: # val > srt_lst[mid]
            left = mid + 1
    return ind

```

```

def binary_search(srt_lst, val):
    left = 0
    right = len(srt_lst) - 1
    ind = None
    found = False
    while ((found == False) and (left <= right)):
        mid = (left + right) // 2
        if (srt_lst[mid] == val):
            ind = mid
            found = True
        elif (val < srt_lst[mid]):
            right = mid - 1
        else: # val > srt_lst[mid]
            left = mid + 1
    return ind

```

$\Theta(1)$  (for the first four lines)  
 $\Theta(1)$  (for the while loop body)  
 $\Theta(1)$  (for the return statement)

```

def binary_search(srt_lst, val):
    left = 0
    right = len(srt_lst) - 1
    ind = None
    found = False
    while ((found == False) and (left <= right)):
        mid = (left + right) // 2
        if (srt_lst[mid] == val):
            ind = mid
            found = True
        elif (val < srt_lst[mid]):
            right = mid - 1
        else: # val > srt_lst[mid]
            left = mid + 1
    return ind

```

$\Theta(1)$  (for the first four lines)  
 $\Theta(1)$  (for the while loop body)  
 $\Theta(1)$  (for the return statement)



```

def binary_search(srt_lst, val):
    left = 0
    right = len(srt_lst) - 1
    ind = None
    found = False
    while ((found == False) and (left <= right)):
        mid = (left + right) // 2
        if (srt_lst[mid] == val):
            ind = mid
            found = True
        elif (val < srt_lst[mid]):
            right = mid - 1
        else: # val > srt_lst[mid]
            left = mid + 1
    return ind

```

$\Theta(1)$  (for the first four lines)  
 $\Theta(\# \text{ of iterations})$  (for the while loop)  
 $\Theta(1)$  (for the if/elif/else block)  
 $\Theta(1)$  (for the return statement)

# The Sorted-Search Problem

[illegible]

# The Sorted-Search Problem

[illegible]

# The Sorted-Search Problem

[illegible]

# The Sorted-Search Problem

[illegible]

# The Sorted-Search Problem

[illegible]

# The Sorted-Search Problem

[illegible]

# The Sorted-Search Problem

[illegible]



# The Sorted-Search Problem

[illegible]

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
<i>1</i>	<i><math>n</math></i>
<i>2</i>	$\frac{n}{2}$
<i>3</i>	$\frac{n}{4}$

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
<i>1</i>	<i><b><math>n</math></b></i>
<i>2</i>	$\frac{n}{2}$
<i>3</i>	$\frac{n}{4}$
<i>4</i>	

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
<i>1</i>	<i><math>n</math></i>
<i>2</i>	$\frac{n}{2}$
<i>3</i>	$\frac{n}{4}$
<i>4</i>	$\frac{n}{8}$

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
<i>1</i>	<i><math>n</math></i>
<i>2</i>	$\frac{n}{2}$
<i>3</i>	$\frac{n}{4}$
<i>4</i>	$\frac{n}{8}$
$\vdots$	$\vdots$

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
$1$	$n$
$2$	$\frac{n}{2}$
$3$	$\frac{n}{4}$
$4$	$\frac{n}{8}$
$\vdots$	$\vdots$
$k$	

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
$1$	$n$
$2$	$\frac{n}{2}$
$3$	$\frac{n}{4}$
$4$	$\frac{n}{8} = \frac{n}{2^3}$
$\vdots$	$\vdots$
$k$	

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
$1$	$n$
$2$	$\frac{n}{2}$
$3$	$\frac{n}{4} = \frac{n}{2^2}$
$4$	$\frac{n}{8} = \frac{n}{2^3}$
$\vdots$	$\vdots$
$k$	



# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
$1$	$n$
$2$	$\frac{n}{2} = \frac{n}{2^1}$
$3$	$\frac{n}{4} = \frac{n}{2^2}$
$4$	$\frac{n}{8} = \frac{n}{2^3}$
$\vdots$	$\vdots$
$k$	

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
$1$	$\mathbf{n} = \frac{n}{2^0}$
$2$	$\frac{n}{2} = \frac{n}{2^1}$
$3$	$\frac{n}{4} = \frac{n}{2^2}$
$4$	$\frac{n}{8} = \frac{n}{2^3}$
$\vdots$	$\vdots$
$k$	

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
$1$	$\boldsymbol{n} = \frac{n}{2^0}$
$2$	$\frac{n}{2} = \frac{n}{2^1}$
$3$	$\frac{n}{4} = \frac{n}{2^2}$
$4$	$\frac{n}{8} = \frac{n}{2^3}$
$\vdots$	$\vdots$
$k$	$\frac{n}{2^{k-1}}$

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
$1$	$\mathbf{n} = \frac{n}{2^0}$
$2$	$\frac{n}{2} = \frac{n}{2^1}$
$3$	$\frac{n}{4} = \frac{n}{2^2}$
$4$	$\frac{n}{8} = \frac{n}{2^3}$
$\vdots$	$\vdots$
$k$	$\frac{n}{2^{k-1}}$
$\vdots$	$\vdots$

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
$1$	$\mathbf{n} = \frac{n}{2^0}$
$2$	$\frac{n}{2} = \frac{n}{2^1}$
$3$	$\frac{n}{4} = \frac{n}{2^2}$
$4$	$\frac{n}{8} = \frac{n}{2^3}$
$\vdots$	$\vdots$
$k$	$\frac{n}{2^{k-1}}$
$\vdots$	$\vdots$
	$1$

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
$1$	$\mathbf{n} = \frac{n}{2^0}$
$2$	$\frac{n}{2} = \frac{n}{2^1}$
$3$	$\frac{n}{4} = \frac{n}{2^2}$
$4$	$\frac{n}{8} = \frac{n}{2^3}$
$\vdots$	$\vdots$
$k$	$\frac{n}{2^{k-1}}$
$\vdots$	$\vdots$
$?$	$1$

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
$1$	$n = \frac{n}{2^0}$
$2$	$\frac{n}{2} = \frac{n}{2^1}$
$3$	$\frac{n}{4} = \frac{n}{2^2}$
$4$	$\frac{n}{8} = \frac{n}{2^3}$
$\vdots$	$\vdots$
$k$	$\frac{n}{2^{k-1}}$
$\vdots$	$\vdots$
$?$	$1$

$$\frac{n}{2^{k-1}} = 1$$

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
$1$	$n = \frac{n}{2^0}$
$2$	$\frac{n}{2} = \frac{n}{2^1}$
$3$	$\frac{n}{4} = \frac{n}{2^2}$
$4$	$\frac{n}{8} = \frac{n}{2^3}$
$\vdots$	$\vdots$
$k$	$\frac{n}{2^{k-1}}$
$\vdots$	$\vdots$
$?$	$1$

$$\frac{n}{2^{k-1}} = 1$$

$$\Downarrow$$

$$n = 2^{k-1}$$



# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
1	$n = \frac{n}{2^0}$
2	$\frac{n}{2} = \frac{n}{2^1}$
3	$\frac{n}{4} = \frac{n}{2^2}$
4	$\frac{n}{8} = \frac{n}{2^3}$
$\vdots$	$\vdots$
$k$	$\frac{n}{2^{k-1}}$
$\vdots$	$\vdots$
?	1

$$\begin{aligned}
 \frac{n}{2^{k-1}} &= 1 \\
 \Downarrow \\
 n &= 2^{k-1} \\
 \Downarrow \\
 \log_2(n) &= \log_2(2^{k-1})
 \end{aligned}$$

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
1	$n = \frac{n}{2^0}$
2	$\frac{n}{2} = \frac{n}{2^1}$
3	$\frac{n}{4} = \frac{n}{2^2}$
4	$\frac{n}{8} = \frac{n}{2^3}$
$\vdots$	$\vdots$
$k$	$\frac{n}{2^{k-1}}$
$\vdots$	$\vdots$
?	1

$$\frac{n}{2^{k-1}} = 1$$

$$\Downarrow$$

$$n = 2^{k-1}$$

$$\Downarrow$$

$$\log_2(n) = \log_2(2^{k-1})$$

$$\Downarrow$$

$$\log_2(n) = (k-1)\log_2(2)$$

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
1	$n = \frac{n}{2^0}$
2	$\frac{n}{2} = \frac{n}{2^1}$
3	$\frac{n}{4} = \frac{n}{2^2}$
4	$\frac{n}{8} = \frac{n}{2^3}$
$\vdots$	$\vdots$
$k$	$\frac{n}{2^{k-1}}$
$\vdots$	$\vdots$
?	1

$$\begin{aligned}
 \frac{n}{2^{k-1}} &= 1 \\
 \Downarrow \\
 n &= 2^{k-1} \\
 \Downarrow \\
 \log_2(n) &= \log_2(2^{k-1}) \\
 \Downarrow \\
 \log_2(n) &= (k-1) \underbrace{\log_2(2)}_1
 \end{aligned}$$

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
1	$n = \frac{n}{2^0}$
2	$\frac{n}{2} = \frac{n}{2^1}$
3	$\frac{n}{4} = \frac{n}{2^2}$
4	$\frac{n}{8} = \frac{n}{2^3}$
$\vdots$	$\vdots$
$k$	$\frac{n}{2^{k-1}}$
$\vdots$	$\vdots$
?	1

$$\begin{aligned}
 \frac{n}{2^{k-1}} &= 1 \\
 \Downarrow \\
 n &= 2^{k-1} \\
 \Downarrow \\
 \log_2(n) &= \log_2(2^{k-1}) \\
 \Downarrow \\
 \log_2(n) &= (k-1) \underbrace{\log_2(2)}_1 \\
 \Downarrow \\
 \log_2(n) &= (k-1)
 \end{aligned}$$

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
1	$n = \frac{n}{2^0}$
2	$\frac{n}{2} = \frac{n}{2^1}$
3	$\frac{n}{4} = \frac{n}{2^2}$
4	$\frac{n}{8} = \frac{n}{2^3}$
$\vdots$	$\vdots$
$k$	$\frac{n}{2^{k-1}}$
$\vdots$	$\vdots$
?	1

$$\frac{n}{2^{k-1}} = 1$$

$$\Downarrow$$

$$n = 2^{k-1}$$

$$\Downarrow$$

$$\log_2(n) = \log_2(2^{k-1})$$

$$\Downarrow$$

$$\log_2(n) = (k-1)\log_2(2)$$

$$\Downarrow$$

$$\log_2(n) = (k-1)$$

$$\Downarrow$$

$$k = 1 + \log_2(n)$$

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
1	$n = \frac{n}{2^0}$
2	$\frac{n}{2} = \frac{n}{2^1}$
3	$\frac{n}{4} = \frac{n}{2^2}$
4	$\frac{n}{8} = \frac{n}{2^3}$
$\vdots$	$\vdots$
$k$	$\frac{n}{2^{k-1}}$
$\vdots$	$\vdots$
?	1

$$\frac{n}{2^{k-1}} = 1$$

$$\Downarrow$$

$$n = 2^{k-1}$$

$$\Downarrow$$

$$\log_2(n) = \log_2(2^{k-1})$$

$$\Downarrow$$

$$\log_2(n) = (k-1)\log_2(2)$$

$$\Downarrow$$

$$\log_2(n) = (k-1)$$

$$\Downarrow$$

$$k = 1 + \log_2(n) = \theta(\log_2(n))$$

# The Sorted-Search Problem

Iteration Number	Size of Searching-Range
1	$n = \frac{n}{2^0}$
2	$\frac{n}{2} = \frac{n}{2^1}$
3	$\frac{n}{4} = \frac{n}{2^2}$
4	$\frac{n}{8} = \frac{n}{2^3}$
$\vdots$	$\vdots$
$k$	$\frac{n}{2^{k-1}}$
$\vdots$	$\vdots$
?	1

$$\frac{n}{2^{k-1}} = 1$$

$$\Downarrow$$

$$n = 2^{k-1}$$

$$\Downarrow$$

$$\log_2(n) = \log_2(2^{k-1})$$

$$\Downarrow$$

$$\log_2(n) = (k-1)\log_2(2)$$

$$\Downarrow$$

$$\log_2(n) = (k-1)$$

$$\Downarrow$$

$$k = 1 + \log_2(n) = \theta(\log_2(n))$$

$$\Downarrow$$

$$\left( \begin{array}{c} \# \text{ of} \\ \text{iterations} \end{array} \right) = \theta(\log_2(n))$$

```

def binary_search(srt_lst, val):
    left = 0
    right = len(srt_lst) - 1
    ind = None
    found = False
    while ((found == False) and (left <= right)):
        mid = (left + right) // 2
        if (srt_lst[mid] == val):
            ind = mid
            found = True
        elif (val < srt_lst[mid]):
            right = mid - 1
        else: # val > srt_lst[mid]
            left = mid + 1
    return ind

```

$\Theta(1)$  (for the first four lines)  
 $\Theta(\# \text{ of iterations})$  (for the while loop)  
 $\Theta(1)$  (for the if/elif/else block)  
 $\Theta(1)$  (for the return statement)



```

def binary_search(srt_lst, val):
    left = 0
    right = len(srt_lst) - 1
    ind = None
    found = False
    while ((found == False) and (left <= right)):
        mid = (left + right) // 2
        if (srt_lst[mid] == val):
            ind = mid
            found = True
        elif (val < srt_lst[mid]):
            right = mid - 1
        else: # val > srt_lst[mid]
            left = mid + 1
    return ind

```

$\Theta(1)$  (for the first four lines)  
 $\Theta(\# \text{ of iterations})$  (for the while loop)  
 $\Theta(1)$  (for the if/elif/else block)  
 $\Theta(1)$  (for the return statement)

$$T_{\text{worst}}(n) = \Theta(\log_2 n)$$

# Linear vs. Logarithmic

# Linear vs. Logarithmic

[illegible]

# Linear vs. Logarithmic

[illegible]

# Linear vs. Logarithmic

[illegible]

# Linear vs. Logarithmic

[illegible]

# Linear vs. Logarithmic

[illegible]

# Linear vs. Logarithmic

[illegible]



# Linear vs. Logarithmic

[illegible]

# Linear vs. Logarithmic

[illegible]

# Linear vs. Logarithmic

[illegible]

# Linear vs. Logarithmic

[illegible]

# Linear vs. Logarithmic

$n$	$\log_2(n)$
2	$\log_2(2) = 1$
$\vdots$	$\vdots$
4	$\log_2(2^2) = 2$
$\vdots$	$\vdots$
$2^3$	$\log_2(2^3) = 3$
$\vdots$	$\vdots$
$2^{10}$	

# Linear vs. Logarithmic

$n$	$\log_2(n)$
2	$\log_2(2) = 1$
$\vdots$	$\vdots$
4	$\log_2(2^2) = 2$
$\vdots$	$\vdots$
$2^3$	$\log_2(2^3) = 3$
$\vdots$	$\vdots$
$2^{10}$	$\log_2(2^{10})$

# Linear vs. Logarithmic

$n$	$\log_2(n)$
2	$\log_2(2) = 1$
$\vdots$	$\vdots$
4	$\log_2(2^2) = 2$
$\vdots$	$\vdots$
$2^3$	$\log_2(2^3) = 3$
$\vdots$	$\vdots$
$2^{10}$	$\log_2(2^{10}) = 10$

# Linear vs. Logarithmic

$n$	$\log_2(n)$
2	$\log_2(2) = 1$
$\vdots$	$\vdots$
4	$\log_2(2^2) = 2$
$\vdots$	$\vdots$
$2^3$	$\log_2(2^3) = 3$
$\vdots$	$\vdots$
$2^{10}$	$\log_2(2^{10}) = 10$
$\vdots$	$\vdots$
$2^{32}$	



# Linear vs. Logarithmic

$n$	$\log_2(n)$
2	$\log_2(2) = 1$
$\vdots$	$\vdots$
4	$\log_2(2^2) = 2$
$\vdots$	$\vdots$
$2^3$	$\log_2(2^3) = 3$
$\vdots$	$\vdots$
$2^{10}$	$\log_2(2^{10}) = 10$
$\vdots$	$\vdots$
$2^{32}$	$\log_2(2^{32})$

# Linear vs. Logarithmic

$n$	$\log_2(n)$
2	$\log_2(2) = 1$
$\vdots$	$\vdots$
4	$\log_2(2^2) = 2$
$\vdots$	$\vdots$
$2^3$	$\log_2(2^3) = 3$
$\vdots$	$\vdots$
$2^{10}$	$\log_2(2^{10}) = 10$
$\vdots$	$\vdots$
$2^{32}$	$\log_2(2^{32}) = 32$

# Linear vs. Logarithmic

$n$	$\log_2(n)$
2	$\log_2(2) = 1$
$\vdots$	$\vdots$
4	$\log_2(2^2) = 2$
$\vdots$	$\vdots$
$2^3$	$\log_2(2^3) = 3$
$\vdots$	$\vdots$
$2^{10}$	$\log_2(2^{10}) = 10$
$\vdots$	$\vdots$
$2^{32}$	$\log_2(2^{32}) = 32$
$\vdots$	$\vdots$
$2^{1000}$	

# Linear vs. Logarithmic


$n$	$\log_2(n)$
2	$\log_2(2) = 1$
$\vdots$	$\vdots$
4	$\log_2(2^2) = 2$
$\vdots$	$\vdots$
$2^3$	$\log_2(2^3) = 3$
$\vdots$	$\vdots$
$2^{10}$	$\log_2(2^{10}) = 10$
$\vdots$	$\vdots$
$2^{32}$	$\log_2(2^{32}) = 32$
$\vdots$	$\vdots$
$2^{1000}$	$\log_2(2^{1000})$

# Linear vs. Logarithmic

$n$	$\log_2(n)$
2	$\log_2(2) = 1$
$\vdots$	$\vdots$
4	$\log_2(2^2) = 2$
$\vdots$	$\vdots$
$2^3$	$\log_2(2^3) = 3$
$\vdots$	$\vdots$
$2^{10}$	$\log_2(2^{10}) = 10$
$\vdots$	$\vdots$
$2^{32}$	$\log_2(2^{32}) = 32$
$\vdots$	$\vdots$
$2^{1000}$	$\log_2(2^{1000}) = 1000$
$\vdots$	$\vdots$
$\vdots$	$\vdots$

# Linear vs. Logarithmic

$n$	$\log_2(n)$
2	$\log_2(2) = 1$
$\vdots$	$\vdots$
4	$\log_2(2^2) = 2$
$\vdots$	$\vdots$
$2^3$	$\log_2(2^3) = 3$
$\vdots$	$\vdots$
$2^{10}$	$\log_2(2^{10}) = 10$
$\vdots$	$\vdots$
$2^{32}$	$\log_2(2^{32}) = 32$
$\vdots$	$\vdots$
$2^{1000}$	$\log_2(2^{1000}) = 1000$
$\vdots$	$\vdots$
$\vdots$	$\vdots$



# Linear vs. Logarithmic

$n$	$\log_2(n)$
2	$\log_2(2) = 1$
$\vdots$	$\vdots$
4	$\log_2(2^2) = 2$
$\vdots$	$\vdots$
$2^3$	$\log_2(2^3) = 3$
$\vdots$	$\vdots$
$2^{10}$	$\log_2(2^{10}) = 10$
$\vdots$	$\vdots$
$2^{32}$	$\log_2(2^{32}) = 32$
$\vdots$	$\vdots$
$2^{1000}$	$\log_2(2^{1000}) = 1000$
$\vdots$	$\vdots$
$\vdots$	$\vdots$

